

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications: Department of  
Entomology

Entomology, Department of

---

9-24-1999

## Tissue polyunsaturated fatty acids and a digestive phospholipase A<sub>3</sub> in the primary screwworm, *Cochliomyia hominivorax*

A.R. Nor Aliza  
*University of Nebraska-Lincoln*

R.L. Rana  
*University of Nebraska-Lincoln*

Steven R. Skoda  
*University of Nebraska-Lincoln*, sskoda1@unl.edu

Dennis R. Berkebile  
*University of Nebraska-Lincoln*, DENNIS.BERKEBILE@ars.usda.gov

David W. Stanley  
*University of Nebraska-Lincoln*

Follow this and additional works at: <https://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

---

Nor Aliza, A.R.; Rana, R.L.; Skoda, Steven R.; Berkebile, Dennis R.; and Stanley, David W., "Tissue polyunsaturated fatty acids and a digestive phospholipase A<sub>3</sub> in the primary screwworm, *Cochliomyia hominivorax*" (1999). *Faculty Publications: Department of Entomology*. 1.  
<https://digitalcommons.unl.edu/entomologyfacpub/1>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Pergamon

Insect Biochemistry and Molecular Biology 29 (1999) 1029–1038

*Insect  
Biochemistry  
and  
Molecular  
Biology*

www.elsevier.com/locate/ibmb

## Tissue polyunsaturated fatty acids and a digestive phospholipase $A_2$ in the primary screwworm, *Cochliomyia hominivorax*

A.R. Nor Aliza <sup>a</sup>, R.L. Rana <sup>a</sup>, S.R. Skoda <sup>b</sup>, D.R. Berkebile <sup>b</sup>, David W. Stanley <sup>a,\*</sup>

<sup>a</sup> *Insect Biochemical Physiology Laboratory, University of Nebraska, Lincoln, NE 68583-0816, USA*

<sup>b</sup> *USDA/ARS, Midwest Livestock Insects Research Unit, Department of Entomology, University of Nebraska, Lincoln, NE 68583-0938, USA*

Received 29 March 1999; received in revised form 24 June 1999; accepted 8 July 1999

### Abstract

We report on the presence of arachidonic acid in larval and adult tissues of the primary screwworm, *Cochliomyia hominivorax* and of the secondary screwworm, *C. macellaria*. Arachidonic acid is present in the phospholipids of whole animal extracts of both species. This fatty acid appears to be accumulated during the larval stages, because proportions of arachidonic acid were higher in adults than in larvae. These insects probably obtain the arachidonic acid from dietary phospholipids. We also report on a phospholipase  $A_2$  activity in midgut preparations from third instars of the primary screwworm. Phospholipase  $A_2$  is responsible for hydrolyzing fatty acids from the *sn*-2 position of dietary phospholipids to release essential fatty acids. The screwworm enzyme is similar to mammalian digestive phospholipase  $A_2$ s because it depends on calcium for high catalytic activity, it is sensitive to the site-specific inhibitor oleyloxyethylphosphorylcholine, and it interacts with heparin. We further characterized the screwworm midgut phospholipase  $A_2$  by altering the reaction conditions, including reaction time, radioactive substrate concentration, protein concentration, pH and temperature. We speculate that the biological significance of this enzyme relates to acquiring essential fatty acids, including arachidonic acid, from dietary phospholipids. © 1999 Elsevier Science Ltd. All rights reserved.

**Keywords:** Screwworm; *Cochliomyia hominivorax*; Polyunsaturated fatty acids; Phospholipids; Phospholipase  $A_2$ ; Insect digestion

### 1. Introduction

The significance of C20 polyunsaturated fatty acids (PUFAs) and eicosanoids in the biology of invertebrates is gaining considerable attention (Stanley and Howard, 1998; Stanley, 1999). Eicosanoid is a general term for all oxygenated metabolites of arachidonic acid and two other C20 PUFAs. The structures and biosynthetic pathways of these molecules are illustrated in several reviews (Stanley-Samuelson, 1994; Stanley, 1999). Most of our understanding of eicosanoids is linked to their clinical significance in human and veterinary medicine, and recognition of the broader significance of these compounds represents a substantial shift in appreciation. Stanley and Howard (1998) proposed that eicosanoids should be regarded under a biological, rather than the more widely

held medical paradigm. The main tenet of the biological paradigm lies in recognition that eicosanoids act as lipid mediators in the earliest forms of animal life, and throughout ensuing evolution eicosanoids were recruited in a bewildering array of functions. We imagine that eicosanoids act in all animals, as detailed by Stanley (1999).

The point might be more easily embraced if eicosanoids and C20 PUFAs were present in high quantities in insects, the largest assemblage of invertebrates. Contrarily, most analyses of insect tissue lipids reveal that arachidonic acid, the principle substrate for eicosanoid biosynthesis, and other C20 PUFAs, are present in very low abundance in most terrestrial insect species. These components are often overlooked in routine analyses. This is particularly so for many dipterans. For example, in their very careful analysis of houseflies, *Musca domestica*, Wakayama et al. (1985) recorded arachidonic acid at about 0.04% of adult phospholipid (PL) fatty acids. Similarly, Rapport et al. (1984) were unable to detect any C20 PUFAs in their analyses of fruit fly, *Dro-*

\* Corresponding author. Tel.: +1-402-472-8710; fax: +1-402-472-4687.

E-mail address: dstanley@unlnotes.unl.edu (D.W. Stanley)

*sophila melanogaster*, tissue PL fatty acids. It would appear that C20 PUFAs occur in only vanishing traces in dipteran tissue lipids.

The situation is rather different for at least some of the hematophagous insects. In his lengthy research program on mosquito nutritional requirements, Dadd (1983) found that mosquitoes differ from all known animals, except a tick species, in that they require dietary arachidonic acid, or certain other C20 PUFAs, during their larval stages to develop into normal adulthood. Stanley-Samuelson and Dadd (1981) and Dadd et al. (1987, 1988) later determined that arachidonic acid is present in substantial abundance in mosquito tissue PLs. Similarly, Gadelhak et al. (1995) recorded substantial proportions of arachidonic acid in PLs prepared from stable flies, *Stomoxys calcitrans*. While arachidonic acid is accumulated from blood meals by adult stable flies, the authors concluded that, as in mosquitoes, the larval stages of stable flies are more important for accumulation of C20 PUFAs. Adult female ticks, *Amblyomma americanum*, accumulate substantial proportions of arachidonic acid into salivary gland phospholipids during their lengthy feeding periods (Shipley et al., 1993). Like mosquitoes, *A. americanum* is regarded as unable to biosynthesize arachidonic acid from C18 PUFAs (Bowman et al., 1995).

More recently, we have turned our attention to the primary screwworm, *Cochliomyia hominivorax*, a parasitic insect which feeds on living mammalian tissue, rather than blood per se. Primary screwworms infest the bodies of mammals, usually through wound sites or body openings, producing a disease state known as myiasis. Females of this species deposit their eggs in rows near the edges of wounds. Larvae emerge from eggs after 12–24 h, and make their way into a wound, where they remain in characteristic head-down postures. The insects feed on tissues within the wound. The screwworms develop to the third and last larval stadium, then crawl out of the wound and drop to the ground, burrow under the surface layer and pupate. Adults emerge from the ground, and feed on nectar while they develop to the reproductive phases of adulthood (Laake et al., 1936; Thomas, 1993).

Mammalian tissue lipids are rich in arachidonic acid, and screwworms have the potential to accumulate substantial amounts of this and other C20 PUFAs during the parasitic phase of their lives. To determine whether this is so, we analyzed the tissue lipids prepared from whole larvae, pupae and adults of the primary screwworm. For comparison, we also analyzed similar preparations from the secondary screwworm, *C. macellaria*, a sibling species which feeds on carrion, rather than living tissue. Here we report substantial proportions of arachidonic acid in primary screwworm tissue lipids, and smaller proportions in secondary screwworms.

Most of the arachidonic acid associated with mam-

malian tissue lipids is linked to cellular PLs, and digestive phospholipases are essential enzymes in processing dietary PLs. C18 and C20 PUFAs are typically associated with the *sn*-2 position of PLs, and phospholipase A<sub>2</sub> (PLA<sub>2</sub>) is the enzyme that hydrolyzes fatty acids from the *sn*-2 position of dietary PLs (Dennis, 1994; Waite, 1987). While well understood in mammalian physiology, we have relatively little information on the presence and characteristics of insect digestive PLA<sub>2</sub>s. Recognizing the relatively high proportions of arachidonic acid in primary screwworm tissue lipids, we considered a digestive PLA<sub>2</sub> in screwworm midgut contents. In this paper we also report on the partial enrichment and characterization of a digestive PLA<sub>2</sub> in the midguts of third instars of the primary screwworm.

## 2. Materials and methods

### 2.1. Chemicals

Radioactive phosphatidylcholine (1-palmitoyl, 2-arachidonoyl[arachidonoyl-1-<sup>14</sup>C], 1.85 GBq/mmol) and oleyloxyethylphosphorylcholine (OOPC) were purchased from New England Nuclear (Boston, MA) and BioMol Inc. (Plymouth Meeting, PA), respectively. The same buffer (0.1 M Tris[hydroxymethyl]aminomethane, pH 9; Sigma Chemical Co., St Louis, MO) was used for dissections, homogenizations and incubations, except in studies on Ca<sup>2+</sup>, as described in Section 3.

For the analysis of tissue fatty acids, fatty acid methyl esters (FAMES) were purchased from Sigma. All solvents were reagents grade from Fisher Scientific or Mallinckrodt.

### 2.2. Insects

Primary screwworms, *C. hominivorax*, (Costa Rican strain CR 92) were reared in a gelled diet in the Screwworm Biocontainment Facility, USDA/ARS, Lincoln, NE. The diet was composed of 70 g dried whole bovine blood, 30 g dried whole egg, 30 g dried nonfat milk, and 12 g acrylamide–acrylate polymer gelling agent per liter of water. Secondary screwworms, *C. macellaria*, were collected in Nebraska, and maintained in routine culture on ground beef (90% lean) in the USDA/ARS facility.

### 2.3. Preparing fatty acid methyl esters

Lipid extraction, separation and analysis followed our routine protocols (Howard and Stanley-Samuelson, 1996). Individual insects were suspended in 2 ml of chloroform:methanol (2:1, v/v) amended with 50 µl of 2% butylated hydroxytoluene to minimize autoxidation of PUFAs. Total lipids were extracted three times as

described. The total lipid extracts were applied to thin-layer chromatography (TLC) plates (20×20 cm, 0.25 mm silica gel G, from Aldrich). The plates were developed in petroleum ether:diethyl ether:acetic acid (80:20:1, v/v). Bands corresponding to TGs and PLs were scraped from the plates and transferred to 15-ml screw cap reaction tubes with Teflon-lined screw caps. The lipid fractions were transmethylated by refluxing in 3 ml of acidified methanol for 90 min. After cooling, the FAMES were extracted three times in petroleum ether. The samples were dried by evaporating the petroleum ether, then added to 25–100 µl of isooctane for gas chromatographic analysis.

#### 2.4. Gas chromatography and gas chromatography–mass spectrometry

The FAMES were analyzed on a Hewlett-Packard 5890 gas chromatograph (San Fernando, CA) equipped with a Supelcowax 10 capillary column (30 m by 0.25 mm, 0.25 µm film thickness, Supelco, Inc., Bellefonte, PA), a flame ionization detector, and a Hewlett-Packard 3390A recording integrator. Injections were conducted in split mode (45:1), and the FAMES were separated by temperature programming from 150 to 230°C at 2°C/min. The carrier gas was helium, flowing at 0.6 ml/min. The components were identified by comparing their chromatographic behaviors to the behaviors of authentic standards (Supelco, Inc.).

The fatty acid identifications were confirmed by gas chromatography–mass spectrometry. The FAME samples were analyzed using a Hewlett-Packard 6890 gas chromatograph (injections done in split mode, 50:1) interfaced with a Hewlett-Packard 5973 mass selective detector. The detector was operated in electron impact mode at 70 eV. The chromatograph was equipped with a column identical to the column just described, and separations were conducted with temperature programming from 160 to 230°C at 2°C/min. The carrier gas was ultra-pure helium at 1 ml/min. Identifications were confirmed by analyzing total ion mass spectra of the FAMES (McCloskey, 1970) and by comparing chromatographic retention times and mass spectra with authentic standards from Supelco.

#### 2.5. PLA<sub>2</sub> source preparation

The alimentary canals were dissected from third instars by making a small cut in the caudal end of the larvae, then gently pulling the entire alimentary canal from the larvae. The tissues were collected into tubes containing Tris buffer amended with 10 µl of phenylthiourea (0.02 mM) to prevent melanization. Samples were homogenized with approximately 500 µl buffer in a glass tissue grinder. The homogenates were centrifuged

at 735 g for 3 min, then at 11,750 g for 10 min. The resulting supernatants were used as enzyme sources.

#### 2.6. PLA<sub>2</sub> activity assay

Radioactive PL substrate was prepared in the form of vesicles to access enzyme activity, following the methods of Reynolds et al. (1991). Substrate (0.05 µCi/reaction) was dispensed into each Eppendorf tube, solvents were removed under N<sub>2</sub> and 80 µl of 0.1 M Tris buffer was added to each Eppendorf tube. For studies on Ca<sup>2+</sup>, Tris buffers were amended with either calcium chloride (5 mM) or EGTA (5 mM) as indicated in Section 3. The tubes were vortexed for 15 s to form vesicles. Substrate vesicle tubes were stored at 4°C for 1–2 h before use.

The PLA<sub>2</sub> assays were initiated by adding the enzyme source to the Eppendorf tubes containing substrate vesicles, and vortexing the tubes for 15 s. In our standard assay conditions, we used 250 µg of protein incubated in 250 µl of total volume for 30 min at 28°C. The effects of Ca<sup>2+</sup> dependency, incubation time, substrate concentration, protein concentration, pH, and temperature were investigated by varying each parameter, as indicated in Section 3.

The reactions were terminated by adding extraction solvent (500 µl of chloroform:methanol; 2:1, v/v) acidified with 100 µl 1.0 N HCl. Arachidonic acid (20 nmol in 10 µl chloroform) was added to each tube as a carrier and free fatty acid standard. Each tube was vortexed for 15 s and centrifuged at 735 g for 2 min. The lower organic phase was transferred to another Eppendorf tube and two more extraction steps followed with 500 µl chloroform.

#### 2.7. Chromatography and liquid scintillation counting

The samples were dried under N<sub>2</sub>, dissolved in 100 µl chloroform, and applied to TLC plates (20×20 cm silica gel G, 0.25 mm; Sigma Chemical Co., St Louis, MO). The TLC plates were developed in petroleum ether:diethyl ether:glacial acetic acid (80:20:1, v/v) and exposed to iodine vapors to visualize the fractions that corresponded to free fatty acids. Peaks of radioactivity associated with PL and free fatty acids were detected on a Bioscan 200 Imaging Scanner (Bioscan Inc., Washington, DC). The PL and free fatty acid fractions were transferred into liquid scintillation vials. The radioactivity in each fraction was estimated by adding Ecolite scintillation medium (ICN Biomedicals Inc., Irvine, CA) then counting on a LKB Wallac 1209 Rackbeta Liquid Scintillation Counter (Pharmacia, Turku, Finland) at 96% counting efficiency for <sup>14</sup>C.

## 2.8. Statistical analyses

Data were analyzed using the analysis of variance in the General Linear Models procedure, and mean comparisons were made using Least Significant Difference test (SAS Institute, 1989).

## 3. Results

### 3.1. Fatty acid compositions of screwworm tissue lipids

The fatty acid compositions of PLs and TGs prepared from third-stage larvae, pupae and adult males and females of the primary screwworms are presented in Tables 1 and 2. The fatty acid profiles of PLs and TGs prepared from the secondary screwworms collected in Jamaica and in Nebraska are fairly similar to the profiles from the primary screwworm, and these data are not repeated. Several comments are appropriate to all data. First, as seen in most analyses of animal tissue lipids, the quantitatively major fatty acids were 16:0, 16:1, 18:1, and 18:2n-6. Smaller proportions of other components were also detected. Second, the screwworms are similar to other dipteran species with relative high proportions of 16:1 in all analyses (12–25%). Third, while 18:2n-6 is a major component of most insect lipids, often comprising more than 50% of PL fatty acids, this compound made up less than 7% of the larval and pupal PLs

in both species. Fourth, more components were recorded from PLs than TGs, and higher proportions of PUFAs were present in PLs than TGs.

Arachidonic acid was present in the tissue lipids of both species, albeit in different proportions. For primary screwworms, this component was present at about 1.5–1.8% of larval and pupal PLs and about 3% of adult PLs. Arachidonic acid was also present in TGs, at about 1% of larval and pupal TGs, about 0.6% of male TGs and nearly 2% of female TGs. The secondary screwworms yielded considerably less arachidonic acid. Arachidonic acid accounted for less than 1% of larval PL fatty acids, and was not detected in pupal lipids. The adults maintained higher arachidonic acid proportions, about 2% of male and female PLs. Arachidonic acid was not present in TGs from secondary screwworms.

Aside from the differences just mentioned, the overall fatty acid profiles from larval and pupal lipids from both species were fairly similar. We note another difference between the two species. One component, 20:2n-6, accounted for about 10% of the TG fatty acids from adults of secondary screwworms. This component was not detected in either lipid fraction prepared from primary screwworms.

### 3.2. Initial experiments and partial enrichment of the digestive PLA<sub>2</sub>

Our initial experiment was designed to assess the presence of a digestive PLA<sub>2</sub> in the primary screwworm,

Table 1

Fatty acid composition, as proportions of total fatty acids ( $\pm$ SEM), in phospholipid and triacylglycerol fractions prepared from total lipid extracts from immature stages of the primary screwworm, *C. hominivorax* (Coq.)<sup>a</sup>

Fatty acids	Larva		Pupa					
	PLs	TGs	PLs	TGs	PLs	TGs		
12:0	–	–	0.13	(0.02)	–	–	0.40	(0.18)
14:0	0.79	(0.04)	2.64	(0.04)	0.90	(0.12)	2.26	(0.14)
15:0	0.13	(0.01)	0.16	(0.01)	–	–	–	–
16:0	22.58	(0.39)	28.28	(0.22)	31.56	(0.59)	25.87	(0.07)
16:1n-6	25.83	(0.43)	20.78	(0.83)	18.22	(1.58)	18.58	(1.88)
17:0	–	–	–	–	–	–	0.26	(0.02)
17:1	–	–	–	–	–	–	0.34	(0.02)
18:0	1.41	(0.03)	4.20	(0.09)	1.58	(0.03)	4.46	(0.03)
18:1	38.57	(0.37)	36.68	(0.65)	34.29	(0.82)	35.15	(0.89)
18:2	6.65	(0.09)	5.86	(0.15)	6.84	(0.79)	6.51	(0.73)
18:3n-6	0.14	(0.01)	–	–	–	–	–	–
18:3n-3	0.12	(0.03)	0.12	(0.01)	–	–	–	–
20:0	0.70	(0.03)	–	–	3.49	(1.69)	–	–
20:1	0.53	(0.02)	–	–	–	–	0.54	(0.10)
20:2	–	–	0.15	(0.02)	–	–	3.44	(0.62)
20:4n-6	1.53	(0.09)	1.01	(0.06)	1.85	(0.12)	1.40	(0.10)
20:5n-3	0.33	(0.02)	–	–	–	–	–	–
22:0	0.82	(0.01)	–	–	1.58	(0.04)	0.54	(0.12)

<sup>a</sup> Data based on three separate gas–liquid chromatographic analyses. Fatty acid identity was confirmed by gas liquid chromatography–mass spectrometry. (PLs=phospholipids; TGs=triacylglycerols.)

Table 2

Fatty acid composition, as proportions of total fatty acids ( $\pm$ SEM), in phospholipid and triacylglycerol fractions prepared from total lipid extracts from adults of the primary screwworm, *C. hominivorax* (Coq.)<sup>a</sup>

Fatty acids	Male		Female	
	PLs	TGs	PLs	TGs
12:0	–	–	–	–
14:0	0.45	(0.05)	2.70	(0.24)
15:0	–	–	–	–
16:0	24.05	(0.60)	27.37	(1.20)
16:1	13.86	(0.88)	17.82	(1.18)
17:0	–	–	–	–
17:1	–	–	–	–
18:0	2.62	(0.68)	5.21	(0.30)
18:1	34.56	(0.59)	39.04	(2.11)
18:2	20.62	(0.66)	7.28	(0.49)
18:3n-6	–	–	–	–
18:3n-3	0.75	(0.14)	–	–
20:0	–	–	–	–
20:1	–	–	–	0.49
20:2	–	–	–	–
20:4n-6	3.07	(0.09)	0.57	(0.06)
20:5n-3	–	–	–	–
22:0	–	–	–	–

<sup>a</sup> Data based on three separate gas–liquid chromatographic analyses. Fatty acid identity was confirmed by gas liquid chromatography–mass spectrometry. (PLs=phospholipids; TGs=triacylglycerols.)

using the reaction conditions we had established for tobacco hornworms, *Manduca sexta*, and larval mosquitoes, *Aedes aegypti*. Unfractionated homogenates of midguts and associated midgut contents yielded very low PLA<sub>2</sub> activity, less than 2 pmol/mg protein/min, compared to about 80 pmol/mg protein/min with the mosquito PLA<sub>2</sub> (Nor Aliza and Stanley, 1998). This finding supported the idea that screwworms express a digestive PLA<sub>2</sub>, however, the low specific activity indicated an enrichment step would be required before attempting further characterization.

Because we had achieved a partial enrichment of the tobacco hornworm digestive PLA<sub>2</sub> on ammonium sulfate precipitation (Rana et al., 1998), we subjected screwworm midgut homogenates to the same treatment. We did not recover PLA<sub>2</sub> activity, and concluded ammonium sulfate precipitation was not an effective first step. Many PLA<sub>2</sub>s from mammalian sources interact with heparin, which can be used to partially enrich the PLA<sub>2</sub> by affinity chromatography. We loaded 0.5-ml samples of filtered midgut homogenates (using a 0.45- $\mu$ m membrane) onto a heparin affinity column (1 ml HiTrap Heparin, Pharmacia Biotech AB, purchased from Sigma Chemical Co., St Louis, MO). After a 30-min incubation period, the column was sequentially washed with four buffers, first with Tris homogenization buffer, second with 0.15 M NaCl, third with 1.05 M NaCl and finally with 1.5 M NaCl. Highest specific activity was recovered in the 1.05 M NaCl fraction (about 8 pmol/mg protein/min), indicating this step effectively enriched the target PLA<sub>2</sub> to allow further characterization.

Enrichment on the affinity column may contribute to the analysis of this enzyme in two ways. First, the enrichment step effectively eliminates some of the protein provided in protein-rich diets from the midgut contents. Second, we had noted that PLA<sub>2</sub> activity in the midgut homogenates was not stable to overnight storage in the refrigerator nor at  $-20^{\circ}\text{C}$ . On the other hand, after the affinity chromatography step, the 1.05 M NaCl fractions were stable to long term storage, which is typical of secretory PLA<sub>2</sub>s. We suggest that the enrichment step separates the PLA<sub>2</sub> from digestive proteases, which degrade the PLA<sub>2</sub> during storage. The high salt concentration of the elution buffer may also act to stabilize the protein. For all subsequent work with the screwworm digestive PLA<sub>2</sub>, freshly prepared midgut homogenates were immediately processed through the heparin columns.

The digestive PLA<sub>2</sub>s associated with preparations from mosquito larvae and tobacco hornworms differ from all known secretory PLA<sub>2</sub>s because these two enzymes do not require the presence of calcium for full catalytic activity. In contrast, reactions performed in the presence of calcium yielded lower PLA<sub>2</sub> activity than reactions in the presence of a calcium chelator (Nor Aliza and Stanley, 1998; Rana et al., 1998). On the basis of these findings, we assessed the influence of calcium on the screwworm preparations. We found that added calcium (5 mM) did not yield significantly increased PLA<sub>2</sub> activity in crude fractions (Fig. 1). Alternatively, reactions with crude homogenates in the presence of 5 mM EGTA yielded virtually no PLA<sub>2</sub> activity




Fig. 1. The influence of Ca<sup>2+</sup> and the Ca<sup>2+</sup>-chelator EGTA on PLA<sub>2</sub> activity in third instar primary screwworm midgut preparations. Enzyme activities were determined in unfractionated samples prepared in three separate buffers, as indicated. The histogram bars represent means of three replicates, and the error bars represent 1 SEM.

(significantly different from other treatments at  $P < 0.05$ ), suggesting the screwworm digestive PLA<sub>2</sub> requires calcium for catalytic activity, as seen in all mammalian and venom-associated secretory PLA<sub>2</sub>s. The screwworm culture medium provides copious milk solids, which provide abundant calcium. We suggest that the added calcium did not enhance the screwworm PLA<sub>2</sub> activity because a sufficiency of calcium was already present. All subsequent PLA<sub>2</sub> reactions were conducted in Tris buffer without added calcium.

### 3.3. A characterization of the primary screwworm PLA<sub>2</sub> activity

The accumulation of free arachidonic acid, the reaction product, increased with reaction time from 10 to 60 min (Fig. 2). As seen in the other insect digestive PLA<sub>2</sub>s, the screwworm PLA<sub>2</sub> was not inhibited by the accumulated reaction product during the 1-h incubations. We used 45-min incubations in subsequent experiments.

The radioactive substrate concentration influenced the PLA<sub>2</sub> activity (Fig. 3). At 0.5 and 1.0 nmol/reaction, we obtained approximately 7 pmol/mg/min of enzyme activity, which was roughly doubled at 2.0 nmol/reaction. Although reactions in the presence of higher radioactive substrate concentration yielded significantly more enzyme activity, we used 1.0 nmol/reaction in subsequent work.

The influence of protein concentration on PLA<sub>2</sub> activity is shown in Fig. 4. PLA<sub>2</sub> activity increased in a linear way up to 250 μg protein/reaction, then plateaued. We used 250 μg protein/reaction in all subsequent experiments.

The influence of pH on the screwworm digestive PLA<sub>2</sub> activity is displayed in Fig. 5. Over the range of




Fig. 2. Progress of the primary screwworm digestive PLA<sub>2</sub> reaction. PLA<sub>2</sub> activity in third instar midgut preparations enriched on affinity column chromatography was assessed as described in Section 2. Each point represents the mean of three replicates, and the error bars indicate 1 SEM.




Fig. 3. The influence of radioactive substrate concentration on PLA<sub>2</sub> activity in third instar primary screwworm midgut preparations. PLA<sub>2</sub> activity in third instar midgut preparations enriched on affinity column chromatography was assessed as described in Section 2. Each point represents the mean of three replicates, and the error bars indicate 1 SEM.

pH 5–10, enzyme activity at pH 9 was significantly higher than all other pH values (LSD,  $P < 0.0005$ ). In particular, PLA<sub>2</sub> activity was substantially reduced at pH 8 and below. We used pH 9 for all subsequent experiments.

The screwworm digestive PLA<sub>2</sub> was sensitive to temperature (Fig. 6). Highest activity, approximately 12 pmol/mg/min, obtained at 30°C. PLA<sub>2</sub> activity was significantly reduced at 20 and 40°C, and virtually abolished at higher temperatures.




Fig. 4. The influence of midgut protein concentration on primary screwworm digestive PLA<sub>2</sub> activity. PLA<sub>2</sub> activity in third instar midgut preparations enriched on affinity column chromatography was assessed as described in Section 2. Each point represents the mean of three replicates, and the error bars indicate 1 SEM.




Fig. 5. The influence of pH on primary screwworm digestive PLA<sub>2</sub> activity. PLA<sub>2</sub> activity in third instar midgut preparations enriched on affinity column chromatography was assessed as described in Section 2. Each point represents the mean of three replicates, and the error bars indicate 1 SEM.

Mammalian digestive PLA<sub>2</sub>s are inhibited in the presence of some phospholipid analogues, including OOPC. By contrast, the digestive PLA<sub>2</sub>s associated with mosquito and tobacco hornworm midgut contents are not influenced by OOPC, even at very high concentrations (Rana et al., 1997; Nor Aliza and Stanley, 1998). PLA<sub>2</sub>




Fig. 6. The influence of temperature on primary screwworm digestive PLA<sub>2</sub> activity. PLA<sub>2</sub> activity in larval midgut preparations enriched on affinity column chromatography was assessed as described in Section 2. Each point represents the mean of three replicates, and the error bars indicate 1 SEM.

activity in the screwworm preparations was sensitive to OOPC. As seen in preparations from mammalian sources, PLA<sub>2</sub> activity decreased in the presence of increasing concentration of OOPC in the range of 5–500 µM (Fig. 7).

We have noted that an apparent PLA<sub>2</sub> activity can follow from other enzymatic pathways (Nor Aliza and Stanley, 1998). For example, phospholipase C could hydrolyze the choline base and its associated phosphate




Fig. 7. The influence of the site-specific inhibitor OOPC on primary screwworm PLA<sub>2</sub> activity. PLA<sub>2</sub> activity in third instar midgut preparations enriched on affinity column chromatography was assessed as described in Section 2. Each point represents the mean of three replicates, and the error bars indicate 1 SEM.

group from the PL backbone, leaving a diacylglycerol with a radioactive fatty acid associated with the *sn*-2 position. A subsequent lipase or esterase step could hydrolyze the radioactive fatty acid, yielding free radioactive fatty acid product that is indistinguishable from the product of a single PLA<sub>2</sub> step. Our routine thin-layer separations of products from PLA<sub>2</sub> reactions did not reveal radioactivity in the diacylglycerol fraction, indicating the activity we observed was due to a PLA<sub>2</sub> step and not to an alternative pathway. The data are similar to published radiochromatograms (Nor Aliza and Stanley, 1998), and are not repeated here.

#### 4. Discussion

The information in this report documents the fatty acid compositions of PLs and TGs prepared from larvae, pupae, and adults of the primary and secondary screwworms. Dipterans differ from other insect groups in their fatty acid compositions with characteristically high proportions of 16:1 in their tissue lipids (Stanley-Samuels and Dadd, 1983). Some features of the fatty acid profiles of juvenile and adult screwworms are in line with the general dipteran pattern. We recorded 16:1 in relatively high proportions of screwworm fatty acids, ranging from 13 to 25% of the PL and TG fatty acids of both species. For larvae and pupae of both screwworm species, we determined 18:2n-6 at about 3–6% of total PL and TG fatty acids. This differs from the background of animal lipids and other insects, in which 18:2n-6 is more abundant in PLs. In contrast to larvae and pupae, 18:2n-6 is considerably more abundant in PLs prepared from adults of both screwworm species. Stanley-Samuels et al. (1988) suggested that fatty acid patterns represent the outcomes of complex metabolic events. We infer from the differing patterns recorded for juvenile and adult screwworms that screwworms undergo considerable changes in lipid metabolism in the pupal–adult transition.

Most terrestrial insects maintain very low proportions of 20:4n-6 in their tissue lipids, often seen only as traces (Stanley-Samuels, 1994). For example, we recorded less than 1% of 20:4n-6 in PLs prepared from Malpighian tubules from larval, pupal and adults of the tenebrionid beetle, *Zophobas atratus* (Howard and Stanley-Samuels, 1996). In their work on houseflies, Wakayama et al. (1985) recorded 20:4n-6 at 0.04% of total fatty acids associated with PLs from newly emerged males. The accumulation of tissue arachidonic acid is otherwise for screwworms and certain other dipterans. Arachidonic acid was present in PLs and TGs of primary screwworm larvae and pupae at about 1–2% of total fatty acids. Again, the adults were different from the juveniles. Arachidonic acid was enriched in PLs from adults (about 3–4% of total fatty acids). Compared to primary

screwworms, lower proportions of arachidonic acid were present in tissue lipids of secondary screwworms. In larvae, arachidonic acid made up 0.4% of PL fatty acids, and was not detected in TGs. No fatty acid components beyond 18:3n-3 were present in pupal PLs or TGs.

We considered the possibility that geographical races of secondary screwworms would generate differing fatty acid compositions. Secondary screwworms collected from the field in Jamaica yielded fatty acid patterns quite similar to the patterns registered from secondary screwworms collected in Nebraska. Hence, environmental and geographical differences between Nebraska and Jamaica did not influence secondary screwworm fatty acid metabolism.

The relatively high abundance of arachidonic acid in screwworm lipids could, of course, be simple reflections of dietary inputs. However, comparisons of the fatty acid compositions of insect tissue lipids and the corresponding diets of the insects suggest that insects express fatty acid incorporation systems that are selective. The selectivity produces tissue lipid fatty acid patterns that are quite different from the diets. For example, in work on fatty acid compositions in tissues of the tobacco hornworm, we found higher proportions of arachidonic acid in the hornworm diet and in hornworm hemolymph than in the tissue lipids (Ogg and Stanley-Samuels, 1992). We investigated this point in more detail by tracing the incorporation of radioactive fatty acids into hornworm hemocytes (Gadelhak and Stanley-Samuels, 1994). When present in substantial quantities, free arachidonic acid can be deleterious to cells, and the hemocytes rapidly incorporated radioactive arachidonic acid into all cellular lipid pools. Soon after the initial, rapid incorporation, however, the arachidonic acid was selectively removed from hemocyte PLs, apparently to maintain very low levels of this component in PLs. We infer from these observations that high levels of arachidonic acid in some diets, such as mammalian blood or tissues, does not necessarily result in accumulation in tissue lipids. In particular, the relatively high proportions of arachidonic acid in PLs from primary screwworms, stable flies and mosquitoes are not passive consequences of diet.

We take an alternative view that some dipterans accumulate PUFAs for special purposes. For example, mosquitoes require dietary arachidonic acid during their larval phases to complete development to normal adults (Dadd, 1983). Part, but not all, of the arachidonic acid requirements could be spared by dietary PGs, from which Dadd inferred that arachidonic acid served multiple roles in the physiology of mosquitoes, some involving PG biosynthesis and others relating to structural components of cellular membranes (Dadd and Kleinjan, 1988). Similarly, adult stable flies maintain substantial proportions (1–5% of total fatty acids) of arachidonic acid in their tissue PLs (Gadelhak et al., 1995). As in

mosquitoes, it appears that the larval stages are most important for accumulating this component. The biological significance of the arachidonic acid may relate to nutritional interactions between blood-feeding insects and their hosts. Certain PGs act as anti-hemostatic factors in vertebrates. Adult female mosquitoes and adult stable flies of both sexes are hematophagous; perhaps the arachidonic acid is converted into PGs which facilitate blood feeding in these insects, as suggested for ticks (Sauer et al., 1993).

By comparison to the hematophagous and parasitic dipterans, arachidonic acid is generally not recorded in fatty acid analyses of other dipterans. This is so for the olive fruit fly, *Dacus oleae* (Madariaga et al., 1974), the Mediterranean fruit fly, *Ceratitis capitata* (Madariaga et al., 1972), and the fruit fly *Drosophila melanogaster* (Rapport et al., 1984). The results with *D. melanogaster* remain unclear, however, because Pages et al. (1986) recorded PG biosynthesis, which depends on the presence of arachidonic acid, in preparations from *D. melanogaster*. Stanley-Samuelson (1994) suggested that C20 PUFAs occur in these insects in very low levels, below the ranges of detectability on normal gas chromatography. These low catalytic levels would support PG biosynthesis for local physiological purposes.

Here, we also report on a digestive PLA<sub>2</sub> in midgut preparations from third instar screwworms. We now have basic descriptive information on digestive PLA<sub>2</sub>s from six insect species, including two predacious insects, the tiger beetle, *C. circumpecta*, and a robber fly, *Asilis* sp. (Uscian et al., 1995), the burying beetle, *N. marginatus* (Rana et al., 1997), a lepidopteran, *M. sexta* (Rana et al., 1998), a mosquito, *A. aegypti* (Nor Aliza and Stanley, 1998) and now the primary screwworm. We perceive fundamental differences among the PLA<sub>2</sub>s from these sources. The PLA<sub>2</sub>s from the tiger beetle, robber fly and screwworms require calcium for full catalytic activity and they are sensitive to the site-specific PLA<sub>2</sub> inhibitor, OOPC. We gather there are structural similarities between these insect enzymes and their mammalian counterparts, which also require calcium (Waite, 1987). The PLA<sub>2</sub> associated with oral secretions of adult burying beetles was sensitive to the presence of the calcium chelator EGTA (Rana et al., 1997), albeit not to the extent seen in the tiger beetle, robber fly and screwworms. The PLA<sub>2</sub>s from mosquito larvae and tobacco hornworms differed considerably from the others because enzyme activity increased, rather than decreased, in the presence of EGTA and the enzymes were not sensitive to OOPC. Moreover, the PLA<sub>2</sub> from screwworms interacted with heparin while the digestive PLA<sub>2</sub> from the tobacco hornworms did not (Rana, unpublished observation).

As to the usual biophysical parameters, the digestive PLA<sub>2</sub>s from all insect sources we have examined so far require fairly high pH for optimal catalytic activity. This

is in accord with expectation, because most insect midguts maintain alkaline conditions. The digestive PLA<sub>2</sub>s from most insect sources are fairly stable to temperature, however, the screwworm enzyme exhibited a relatively narrow temperature optimum. This may relate to the feeding ecology of screwworms, which live under stable temperature conditions within mammalian tissues.

We also note considerable differences in specific activities among these preparations under similar reaction conditions. Unfractionated larval mosquito preparations yielded approximately 80 pmol/mg protein/min of PLA<sub>2</sub> activity (Nor Aliza and Stanley, 1998), compared to about 40 pmol/mg protein/min with similar preparations of burying beetle oral secretions (Rana et al., 1997) and of tobacco hornworm midgut contents (Rana et al., 1998). The PLA<sub>2</sub>-enriched preparations from primary screwworm midguts yielded approximately 8 pmol/mg protein/min of PLA<sub>2</sub> activity, which could be pushed higher with increased substrate concentrations. These enzyme activity values are normalized to protein concentrations in midgut contents, which undoubtedly varies with dietary inputs. Hence, these differences in specific activities may be more apparent than real. We are currently working to purify a digestive PLA<sub>2</sub> from tobacco hornworms, after which we anticipate learning more on this point.

## Acknowledgements

We thank Drs Jack Campbell and Xinzhi Ni for reading a draft of this paper. This is paper no. 12,611, Nebraska Agricultural Research Division, and contribution no. 1026 of the Department of Entomology. This work is supported by the Agricultural Research Division, UNL (Project NEB-17-054) and by a scholarship to Nor Aliza from the University of Malaysia at Sarawak.

## References

- Bowman, A.S., Sauer, J.R., Neese, P.A., Dillwith, J.W., 1995. Origin of arachidonic acid in the salivary glands of the lone star tick, *Amblyomma americanum*. *Insect Biochem. Molec. Biol.* 25, 225–233.
- Dadd, R.H., 1983. Essential fatty acids: insects and vertebrates compared. In: Mittler, T.E., Dadd, R.H. (Eds.), *Metabolic Aspects of Lipid Nutrition in Insects*. Westview Press, Boulder, CO, pp. 107–147.
- Dadd, R.H., Kleinjan, J.E., 1988. Prostaglandin sparing of dietary arachidonic acid in the mosquito, *Culex pipiens*. *J. Insect Physiol.* 34, 779–786.
- Dadd, R.H., Kleinjan, J.E., Asman, S.M., 1988. Eicosapentaenoic acid in mosquito tissues: differences between wild and laboratory-reared adults. *Environ. Ent.* 17, 172–180.
- Dadd, R.H., Kleinjan, J.E., Stanley-Samuelson, D.W., 1987. Polyunsaturated fatty acids of mosquitoes reared with single dietary polyunsaturates. *Insect Biochem.* 17, 7–16.

- Dennis, E.A., 1994. Diversity of group types, regulation, and function of phospholipase A<sub>2</sub>. *J. Biol. Chem.* 269, 13057–13060.
- Gadelhak, G.G., Pedibhotla, V.K., Rosario, R.M.T., Thomas, G.T., Stanley-Samuelson, D.W., 1995. The influence of blood meals on accumulation of arachidonic acid by adult stable flies. *Comp. Biochem. Physiol.* 110B, 613–621.
- Gadelhak, G.G., Stanley-Samuelson, D.W., 1994. Incorporation of polyunsaturated fatty acids into PLs of hemocytes from the tobacco hornworm, *Manduca sexta*. *Insect Biochem. Molec. Biol.* 24, 775–785.
- Howard, R.W., Stanley-Samuelson, D.W., 1996. Fatty acid composition of fat body and Malpighian tubules of the tenebrionid beetle, *Zophobas atratus*: significance in eicosanoid-mediated physiology. *Comp. Biochem. Physiol.* 115B, 429–437.
- Laake, E.W., Cushing, E.C., Parish, H.E., 1936. Biology of the primary screwworm fly, *Cochliomyia americana*, and a comparison of its stages with those of *C. macellaria*. U.S. Dep. Agric. Tech. Bull. 500 (24 pp).
- Madariaga, M.A., Mata, F., Municio, A.M., Ribera, A., 1972. Effect of the lipid composition of the larval diet on the fatty acid composition during development of *Ceratitis capitata*. *Insect Biochem.* 2, 249–256.
- Madariaga, M.A., Mata, F., Municio, A.M., Ribera, A., 1974. Changes in the fatty acid pattern of glycerolipids of *Dacus oleae* during metamorphosis and development. *Insect Biochem.* 4, 151–160.
- McCloskey, J.M., 1970. Mass spectrometry of fatty acid derivatives. In: Gunstone, F.E. (Ed.) *Topics in Lipid Chemistry*, vol. I. Wiley, Inc., New York, pp. 369–440.
- Nor Aliza, A.R., Stanley, D.W., 1998. A digestive phospholipase A<sub>2</sub> in larval mosquitoes, *Aedes aegypti*. *Insect Biochem. Molec. Biol.* 28, 561–569.
- Ogg, C.L., Stanley-Samuelson, D.W., 1992. Phospholipid and triacylglycerol fatty acid compositions of the major life stages and selected tissues of the tobacco hornworm *Manduca sexta*. *Comp. Biochem. Physiol.* 101B, 345–351.
- Pages, M., Rosello, J., Casas, J., Gelpi, E., Gualde, N., Rigaud, M., 1986. Cyclooxygenase and lipoxygenase-like activity in *Drosophila melanogaster*. *Prostaglandins* 32, 729–740.
- Rana, R.L., Hoback, W.W., Nor Aliza, A.R., Bedick, J., Stanley, D.W., 1997. Pre-oral digestion: a phospholipase A<sub>2</sub> associated with oral secretions in adult burying beetles, *Nicrophorus marginatus*. *Comp. Biochem. Physiol.* 118B, 375–380.
- Rana, R.L., Sarath, G., Stanley, D.W., 1998. A digestive phospholipase A<sub>2</sub> in midguts of tobacco hornworm, *Manduca sexta* L. *J. of Insect Physiol.* 44, 297–303.
- Rapport, E.W., Stanley-Samuelson, D.W., Dadd, R.H., 1984. Ten generations of *Drosophila melanogaster* reared axenically on a fatty acid-free holidic diet. *Arch. Insect Biochem. Physiol.* 1, 243–250.
- Reynolds, L.J., Washburns, W.N., Deems, R.A., Dennis, E.A., 1991. Assay strategies and methods for phospholipases. In: Dennis, E.A. (Ed.) *Methods in Enzymology, Phospholipases*, vol. 197. Academic Press, San Diego, CA, pp. 3–23.
- SAS Institute Inc., 1989. *SAS/STAT® User's Guide*, Version 6, vol. 2. 4th ed. SAS Institute Inc. Cary, NC.
- Sauer, J.R., Bowman, A.S., Shipley, M.M., Gengler, C.L., Surdick, M.R., McSwain, J.L., Luo, C., Essenberg, R.C., Dillwith, J.W., 1993. Arachidonate metabolism in tick salivary glands. In: Stanley-Samuelson, D.W., Nelson, D.R. (Eds.), *Insect Lipids: Chemistry, Biochemistry and Biology*. University of Nebraska Press, Lincoln, pp. 99–138.
- Shipley, M.M., Dillwith, J.R., Bowman, A.S., Essenberg, R.C., Sauer, J.R., 1993. Changes in lipids of the salivary glands of the lone star tick, *Amblyomma americanum*, during feeding. *J. Parasitol.* 79, 834–842.
- Stanley, D.W., 1999. *Eicosanoids in Invertebrate Signal Transduction Systems*. Princeton University Press, Princeton, NJ.
- Stanley, D.W., Howard, R.W., 1998. The biology of prostaglandins and related eicosanoids in invertebrates: cellular, organismal and ecological actions. *Amer. Zool.* 38, 369–381.
- Stanley-Samuelson, D.W., 1994. Prostaglandins and related eicosanoids in insects. *Adv. Insect Physiol.* 24, 115–212.
- Stanley-Samuelson, D.W., Dadd, R.H., 1981. Arachidonic and other tissue fatty acids of *Culex pipiens* reared with various concentrations of dietary arachidonic acid. *J. Insect Physiol.* 27, 571–578.
- Stanley-Samuelson, D.W., Dadd, R.H., 1983. Long-chain polyunsaturated fatty acids: patterns of occurrence in insects. *Insect Biochem.* 13, 549–558.
- Stanley-Samuelson, D.W., Jurenka, R.A., Cripps, C., Blomquist, G.J., De Renobales, M., 1988. Fatty acids in insects: composition, metabolism, and biological significance. *Arch. Insect Biochem. Physiol.* 9, 1–33.
- Thomas, D.B., 1993. Behavioral aspects of screwworm ecology. *J. Kansas Entomol. Soc.* 66, 13–30.
- Uscian, J.M., Miller, J.S., Sarath, G., Stanley-Samuelson, D.W., 1995. A digestive phospholipase A<sub>2</sub> in the tiger beetle *Cicindella circum-picta*. *J. Insect Physiol.* 41, 135–141.
- Waite, M., 1987. *Handbook of Lipid Research: The phospholipases*. Plenum Press, New York.
- Wakayama, E.J., Dillwith, J.W., Blomquist, G.J., 1985. Occurrence and metabolism of arachidonic acid in the housefly, *Musca domestica* (L.). *Insect Biochem.* 15, 367–374.