

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from the Harold W. Manter
Laboratory of Parasitology

Parasitology, Harold W. Manter Laboratory of

January 2000

Bibliography of Parasites and Vertebrate Hosts in Arizona, New Mexico and Texas (1893–1984)

Lillian F. Mayberry

The University of Texas at El Paso, mayberry@utep.edu

Albert G. Canaris

The University of Texas at El Paso, agcanaris@utep.edu

John R. Bristol

The University of Texas at El Paso, jbristol@utep.edu

Scott Lyell Gardner

University of Nebraska - Lincoln, slg@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/parasitologyfacpubs>



Part of the [Parasitology Commons](#)

Mayberry, Lillian F.; Canaris, Albert G.; Bristol, John R.; and Gardner, Scott Lyell, "Bibliography of Parasites and Vertebrate Hosts in Arizona, New Mexico and Texas (1893–1984)" (2000). *Faculty Publications from the Harold W. Manter Laboratory of Parasitology*. 2.

<https://digitalcommons.unl.edu/parasitologyfacpubs/2>

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Harold W. Manter Laboratory of Parasitology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

**BIBLIOGRAPHY OF PARASITES AND
VERTEBRATE HOSTS
IN ARIZONA, NEW MEXICO
AND TEXAS (1893 - 1984)**

**Lillian F. Mayberry, Albert G. Canaris and John R. Bristol
Department of Biological Sciences
The University of Texas at El Paso
El Paso, Texas 79968**

and

**Scott L. Gardner
Associate Professor and Curator of Parasites
The Harold W. Manter Laboratory of Parasitology
W-529 Nebraska Hall
University of Nebraska-Lincoln
Lincoln, Nebraska 68588-0514
e-mail: slg@unl.edu
Tel: 402-472-3334
FAX: 402-472-8949**

**Published on the world-wide-web 11 January 2000 on the University of Nebraska Harold
W. Manter Laboratory of Parasitology Web Server.**

TABLE OF CONTENTS

Introduction

Tables

Parasite-Host Bibliography

Protozoa

Cestoda

Trematoda

Nemata

Aanthocephala

Arthropoda

 Acarina-Mite

 Acarina-Tick

 Copepoda-Crustacea

 Insecta-Diptera

 Insecta-Hemiptera

 Insecta-Mallophaga and Anoplura

 Insecta-Siphonaptera

 Pentastomida

 Class Unknown

Host-Parasite Bibliography

 Scientific Name

 Common Name

Literature Cited

Theses and Dissertations not Referenced

Acknowledgments

Introduction:

This bibliographic listing of parasites and hosts reported from Arizona, New Mexico and Texas has been prepared to facilitate literature searches by parasitologists and others needing such information. In the Parasite-Host Bibliography, parasites have been categorized into major taxonomic groups, assigned a number (left column) and listed in alphabetical order for the three states. Those from Arizona have reference numbers (right column) followed by "A", those from New Mexico are followed by "N" with all undesignated being from Texas.

In the Host-Parasite Bibliography, hosts have been listed in alphabetical order by their scientific name followed by parasite numbers which can be used to accession the reference by

referring to the Parasite-Host section. In some cases, however, the host common names were used in the original publication. These are listed alphabetically in a separate section of the Host-Parasite Bibliography.

We sincerely hope there are no errors or omissions. A total of 509 citations are included and 782 hosts and 1,528 parasites are recorded. In addition, two tables summarizing information on helminth parasite geographic distribution and frequency among the major vertebrate hosts are included. In a work of this scope, perfection is difficult, thus, we would appreciate being informed of errors and/or omissions.

TABLES

Table 1. Helminth Parasites Distributed by State in the South Western United States.

| | Parasites Found in Vertebrates in the Southwest | | | | |
|--------------|--|----------------|---------------|-----------------------|--------------|
| STATE | Trematoda | Cestoda | Nemata | Acanthocephala | Total |
| Arizona | 5 | 11 | 23 | 0 | 39 |
| New Mexico | 7 | 20 | 55 | 3 | 85 |
| Texas | 296 | 146 | 271 | 38 | 751 |
| TOTAL | 308 | 177 | 349 | 41 | 875 |

Table 2. Frequency of helminths among vertebrate hosts in the South Western United States.

| | Frequency of Parasites Found Vertebrates in the Southwest | | | | |
|-------------------|--|----------------|---------------|-----------------------|--------------|
| HOST GROUP | Trematoda | Cestoda | Nemata | Acanthocephala | Total |
| Pices | 33 | 11 | 33 | 1 | 78 |
| Amphibia | 35 | 20 | 81 | 4 | 140 |
| Aves | 83 | 102 | 63 | 26 | 274 |
| Mammalia | 67 | 118 | 344 | 24 | 553 |
| TOTAL | 573 | 320 | 689 | 119 | 1701 |

| | | | |
|----|--------------------------------|-------------------------------------|-------------|
| 24 | <i>Henneguya texana</i> | <i>Pogonias cromis</i> | 260 |
| 25 | <i>Isospora californica</i> | <i>Peromyscus californicus</i> | 300N |
| 26 | <i>Isospora rivolta</i> | <i>Canis latrans</i> | 464 |
| 27 | <i>Klossia perplexens</i> | <i>Peromyscus maniculatus</i> | 286A |
| 28 | <i>Klossia variabilis</i> | <i>Plecotus rafinesqii</i> | 410 |
| 29 | <i>Kudoa branchiata</i> | <i>Leiostomus xanthurus</i> | 261 |
| 30 | <i>Lankesterella</i> | <i>Agelaius phoeniceus</i> | 51 |
| | | <i>Cassidix mexicanus</i> | 51 |
| | | <i>Passer domesticus domesticus</i> | 52 |
| | | <i>Sturnella magna</i> | 51 |
| 31 | <i>Lankesterella garnhami</i> | <i>Passer domesticus domesticus</i> | 51 |
| 32 | <i>Leucocytozoon</i> | <i>Hesperiphona vespertina</i> | 506A |
| | | <i>Peucedramus olivaceus</i> | 506A |
| 33 | <i>Myxidium serotinum</i> | <i>Acris crepitans</i> | 49 |
| 34 | <i>Myxosporidia</i> | <i>Cyprinodon variegatus</i> | 396 |
| | | <i>Fundulus similis</i> | 396 |
| | | <i>Mugil cephalus</i> | 396 |
| 35 | <i>Nyctotherus cordiformis</i> | <i>Acris crepitans</i> | 49 |
| 36 | <i>Opalina cf. virguloidea</i> | <i>Acris crepitans</i> | 49 |
| 37 | <i>Plasmodium</i> | <i>Antrozous pallidus</i> | 507 |
| | | <i>Melopelia asiatica mearnsi</i> | 506A |
| | | <i>Pipistrellus hesperus</i> | 507 |
| | | <i>Sceloporus poinsetti</i> | 461 |
| | | <i>Urosaurus ornatus</i> | 461 |
| 38 | <i>Plasmodium chiricahuae</i> | <i>Sceloporus clarki</i> | 400A |
| | | <i>Sceloporus jarrovi</i> | 460A |
| 39 | <i>Plasmodium elongatum</i> | <i>Sturnella magna</i> | 51 |
| 40 | <i>Plasmodium hexamerium</i> | <i>Sturnella magna</i> | 51 |
| 41 | <i>Plasmodium relictum</i> | <i>Passer domesticus domesticus</i> | 51 |
| | | <i>Sturnella magna</i> | 51 |
| 42 | <i>Sarcocystis</i> | <i>Sus scrofa</i> | 113 |
| 43 | <i>Sarcocystis rileyi</i> | <i>Anas clypeata</i> | 58 |
| 44 | <i>Sarcosporidia</i> | <i>Columba livia</i> | 351 |
| 45 | <i>Theileria</i> | <i>Deer</i> | 429 |
| | | <i>Odocoileus virginianus</i> | 183,208,398 |
| 46 | <i>Theileria cervi</i> | <i>Odocoileus virginianus</i> | 407 |
| 47 | <i>Toxoplasma gondii</i> | <i>Canis familiaris</i> | 297N,A |
| | | <i>Canis latrans</i> | 297N,A |
| | | <i>Conepatus mesoleucus</i> | 297N,A |
| | | <i>Erethizon dorsatum</i> | 297N,A |
| | | <i>Felis domesticus</i> | 297N,A |
| | | <i>Felis rufus</i> | 297N,A |
| | | <i>Lepus californicus</i> | 297N,A |
| | | <i>Mephitis mephitis</i> | 297N,A |
| | | <i>Sylvilagus</i> | 297N,A |
| | | <i>Taxidea taxus</i> | 297N,A |
| | | <i>Urocyon cinereargenteus</i> | 297N,A |
| | | <i>Vulpes macrotis</i> | 297N,A |
| 48 | <i>Trichomonad canker</i> | <i>Scardafella inca</i> | 417 |
| 49 | <i>Trichomonas</i> | <i>Phrynosoma cornutum</i> | 482 |
| 50 | <i>Trichomonas gallinae</i> | <i>Columba livia</i> | 351 |
| | | <i>White-winged Dove</i> | 468A |
| 51 | <i>Trypanosoma</i> | <i>Cervus elaphus</i> | 124N |
| | | <i>Hesperiphona vespertina</i> | 506A |
| | | <i>Melopelia asiatica mearnsi</i> | 506A |
| | | <i>Odocoileus virginianus</i> | 407 |

| | | | |
|----|-----------------------------------|---|---------|
| 52 | <i>Trypanosoma cruzi</i> | <i>Canis latrans</i> | 210 |
| | | <i>Dasypus novemcinctus</i> | 350 |
| | | <i>Didelphis virginiana</i> | 350 |
| | | <i>Mus musculus</i> | 350 |
| | | <i>Neotoma albigula</i> | 507A |
| | | <i>Neotoma micropus</i> | 172,350 |
| | | <i>Peromyscus boylii</i> | 507A |
| | | <i>Triatoma ambigua</i> | 126 |
| | | <i>Triatoma gerstreckeri</i> | 349,490 |
| | | <i>Triatoma longipes</i> | 414A |
| | | <i>Triatoma rubida</i> | 414A |
| | | <i>Triatoma sanguisuga</i> | 126 |
| 53 | <i>Trypanosoma lewisi</i> | <i>Neotoma micropus</i> | 172 |
| 54 | <i>Trypanosoma peromysci</i> | <i>Peromyscus boylii</i> | 507A |
| | | <i>Peromyscus truei</i> | 507A |
| 55 | <i>Trypanosoma vespertilionis</i> | <i>Myotis velifer</i> | 507A |
| | | <i>Pipistrellus hesperus</i> | 507 |
| 56 | <i>Tyzzeria pellerdyi</i> | <i>Anas acuta</i> | 57 |
| | | <i>Anas americana</i> | 57 |
| | | <i>Anas creca</i> | 57 |
| | | <i>Anas platyrhynchos platyrhynchos</i> | 57 |
| | | <i>Anas strepera</i> | 57 |

Cestoda

| | | | |
|----|-----------------------------------|-------------------------------|------------|
| 57 | <i>Amoebotaenia fuhrmanni</i> | <i>Capella gallinago</i> | 287 |
| 58 | <i>Anochotaenia globata</i> | <i>Sturnella magna magna</i> | 284 |
| 59 | <i>Anomotaenia</i> | <i>Dendrocygna autumnalis</i> | 205 |
| | | <i>Grus canadensis</i> | 198 |
| 60 | <i>Anomotaenia ciliata</i> | Duck | 68 |
| 61 | <i>Anomotaenia constricta</i> | <i>Corvus brachyrhynchos</i> | 341 |
| | | <i>Corvus cryptoleucus</i> | 371N |
| 62 | <i>Anoplocephala magna</i> | Horse | 378 |
| | | Mule | 378 |
| 63 | <i>Anoplocephala mamillana</i> | Horse | 378 |
| | | Mule | 378 |
| 64 | <i>Aploparaksis occidentalis</i> | <i>Limnodromus griseus</i> | 383 |
| 65 | <i>Aploparaksis veitchi</i> | Duck | 68 |
| 66 | <i>Bakererpes addisi</i> | <i>Chordeiles minor minor</i> | 495 |
| 67 | <i>Biacetabulum meridianum</i> | <i>Carpiodes carpio</i> | 420 |
| 68 | <i>Bothriocephalidae</i> | <i>Lepomis auritus</i> | 479 |
| | | <i>Lepomis macrochirus</i> | 479 |
| 69 | <i>Bothriocephalus</i> | <i>Gambusia affinis</i> | 40,127,128 |
| | | <i>Gambusia geiseri</i> | 127 |
| | | <i>Ictalurus punctatus</i> | 40 |
| | | <i>Lepomis auritus</i> | 479 |
| | | <i>Lepomis gulosus</i> | 479 |
| | | <i>Lepomis punctatus</i> | 479 |
| | | <i>Micropterus salmoides</i> | 211,479 |
| 70 | <i>Bothriocephalus claviceps</i> | <i>Lepomis cyanellus</i> | 308 |
| | | <i>Lepomis humilis</i> | 308 |
| | | <i>Lepomis macrochirus</i> | 308 |
| | | <i>Lepomis megalotis</i> | 308 |
| 71 | <i>Bothriocephalus texomensis</i> | <i>Hiodon alosoides</i> | 419 |
| 72 | <i>Capigens singularis</i> | <i>Ictiobus bubalus</i> | 421 |

| | | | |
|-----|-----------------------------------|----------------------------------|---------------------|
| 73 | <i>Catenotaenia peromysci</i> | Peromyscus maniculatus | 428N |
| 74 | <i>Cestode larva</i> | Gambusia affinis | 128 |
| 75 | <i>Cestode</i> | Myotis velifer | 254 |
| 76 | <i>Choanotaenia</i> | Sturnus vulgaris | 75 |
| 77 | <i>Choanotaenia infundibulum</i> | Phasianus colchicus | 142,366 |
| 78 | <i>Choanotaenia iola</i> | Sturnella magna magna | 284 |
| 79 | <i>Choanotaenia numenii</i> | Numenius americanus | 70N |
| 80 | <i>Cittotaenia variabilis</i> | Rabbit | 378 |
| | | Sylvilagus floridanus | 201 |
| 81 | <i>Cloacotaenia megalops</i> | Anas clypeata | 58 |
| | | Anas creca | 74 |
| | | Anas cyanoptera septentrionalium | 499 |
| | | Dendrocygna autumnalis | 205 |
| 82 | <i>Corallobothrium fimbriatum</i> | Ictalurus nebulosus | 479 |
| | | Ictalurus punctatus | 211,280 |
| 83 | <i>Corallobothrium giganteum</i> | Ictalurus punctatus | 211,280 |
| 84 | <i>Cyclophyllidean larva</i> | Cyprinodon variegatus | 24 |
| 85 | <i>Cycloskrjabina taborensis</i> | Lasiurus cinereus | 73A |
| 86 | <i>Cyclustera capito</i> | Rallus elegans | 37 |
| | | Rallus longirostris | 37 |
| 87 | <i>Clyindrotaenia</i> | Rana pipiens complex | 425 |
| 88 | <i>Cylindrotaenia americana</i> | Acris creptians | 49 |
| | | Acris gryllus | 216 |
| | | Hyla squirella | 216 |
| | | Leiopisma laterale | 216 |
| | | Pseudacris triseriata | 216 |
| 89 | <i>Cysticercoides</i> | Adinia xenica | 24 |
| | | Cyprinodon variegatus | 24 |
| | | Fundulus grandis | 24 |
| | | Mollienisia latipinna | 24 |
| 90 | <i>Cysticercus</i> | Cnemidophorus tessellatus | 206N |
| | | Scincella lateralis | 216 |
| 91 | <i>Cysticercus bovis</i> | Cattle | 378 |
| 92 | <i>Cysticercus cellulosae</i> | Domestic Pig | 377 |
| | | Pig | 378 |
| 93 | <i>Cysticercus fasciolaris</i> | Rattus rattus | 250 |
| | | Sigmodon hispidus | 250 |
| 94 | <i>Cysticercus pisiformis</i> | Rabbit | 378 |
| 95 | <i>Cysticercus tenuicollis</i> | Antilope cervicapra | 463 |
| | | Cattle | 378 |
| | | Goat | 378 |
| | | Odocoileus hemionus | 457 |
| | | Ovis canadensis | 6N,11N,19A, 401A |
| | | Sciurus niger | 91 |
| | | Sheep | 378 |
| 96 | <i>Davainea salmoni</i> | Rabbit | 378 |
| 97 | <i>Dicranotaenia</i> | Dendrocygna autumnalis | 205 |
| 98 | <i>Diochetos phrynosomatis</i> | Phrynosoma cornutum | 216,482 |
| 99 | <i>Diorchis americana</i> | Anas clypeata | 58 |
| 100 | <i>Diorchis bulbodes</i> | Duck | 68 |
| | | Pintail | 69 |
| 101 | <i>Diorchis nyrocae</i> | Anas clypeata | 58 |
| | | Anas cyanoptera septentrionalium | 499 |
| | | Duck | 68 |
| | | Pintail | 69 |

| | | |
|--|----------------------------------|-------------|
| | Shoveler | 69 |
| 102 <i>Diorchis spinata</i> | Anas clypeata | 58 |
| | Duck | 68 |
| | Gadwall | 69 |
| 103 <i>Dipyllobothrium mansonoides</i> | Felis domesticus | 197 |
| 104 <i>Dipylidium caninum</i> | Cat | 378 |
| | Dog | 378,430,438 |
| | Man | 338 |
| 105 <i>Dipylidium sexcoronatum</i> | Cat | 378 |
| | Dog | 378 |
| 106 <i>Echinococcus granulosus</i> | Pig | 378 |
| 107 <i>Echinocotyle</i> | Anas cyanoptera septentrionalium | 499 |
| 108 <i>Echinocotyle rosseteri</i> | Anas clypeata | 58 |
| | Anas creca | 74 |
| 109 <i>Filaroides plerocercoid</i> | Ambystoma tigrinum | 505N |
| 110 <i>Gastrotaenia cygni</i> | Anas clypeata | 58 |
| | Anas creca | 74 |
| 111 <i>Glaradacris bothrium texomensis</i> | Ictiobus bubalus | 421 |
| 112 <i>Glaradacris confusus</i> | Carpiodes carpio | 420 |
| | Ictiobus bubalus | 22A |
| | Ictiobus cyprinellus | 22A |
| | Ictiobus niger | 22A,421 |
| 113 <i>Glossocercus</i> | Cyprinodon variegatus | 24 |
| | Fundulus grandis | 24 |
| | Fundulus similis | 24 |
| 114 <i>Glossocercus cyprinodontis</i> | Cyprinodon variegatus | 24 |
| 115 <i>Gryporhynchus tetrorchis</i> | Ardea herodias herodias | 257 |
| 116 <i>Haploparaxis</i> | Capella gallinago | 287 |
| 117 <i>Haploparaxis brachyphallos</i> | Capella gallinago | 287 |
| 118 <i>Haploparaxis crassirostris</i> | Capella gallinago | 287 |
| 119 <i>Haploparaxis echinovatum</i> | Capella gallinago | 287 |
| 120 <i>Hydatid cyst</i> | Cow | 424 |
| 121 <i>Hydatigera taeniaeformis</i> | Sigmodon hispidus | 298,316 |
| 122 <i>Hymenolepis</i> | Capella gallinago | 287 |
| | Geomys bursarius | 140,184 |
| | Geomys personatus | 140 |
| 123 <i>Hymenolepis calumnacantha</i> | Capella gallinago | 287 |
| 124 <i>Hymenolepis capellae</i> | Limnodromus griseus | 383 |
| 125 <i>Hymenolepis corvi</i> | Corvus brachyrhynchos | 341 |
| | Corvus cryptoleucus | 371N |
| 126 <i>Hymenolepis diminuta</i> | Mus musculus | 78 |
| | Perognathus hispidus | 316 |
| | Rattus rattus | 250 |
| | | 11 |
| | Sigmodon hispidus | 316,328,427 |
| 127 <i>Hymenolepis farciminosa</i> | Sturnus vulgaris | 43 |
| 128 <i>Hymenolepis gertschi</i> | Tadarida brasiliensis | 71N |
| 129 <i>Hymenolepis gracilis</i> | Duck | 68 |
| | Pintail | 69 |
| 130 <i>Hymenolepis hopkinsi</i> | Duck | 68 |
| | Gadwall | 69 |
| | Pintail | 69 |
| 131 <i>Hymenolepis limnodromi</i> | Limnodromus griseus | 383 |

| | | | |
|-----|---|----------------------------------|-------------------------|
| 132 | <i>Hymenolepis megalops</i> | Duck | 68 |
| | | Gadwall | 69 |
| | | Pintail | 69 |
| | | Shoveler | 69 |
| 133 | <i>Hymenolepis murina</i> | Rattus norvegicus | 78 |
| 134 | <i>Hymenolepis stolli</i> | Duck | 68 |
| | | Pintail | 69 |
| 135 | <i>Hymenolepis teresoides</i> | Duck | 68 |
| | | Gadwall | 69 |
| 136 | <i>Isoglaridacris hexacotyle</i> | Catostomus clarki | 497N |
| | | Catostomus insignis | 497N |
| 137 | <i>Liga braziliensis</i> | Meleagris gallopavo intermedia | 361 |
| 138 | <i>Longistriata vexillata</i> | Perognathus hispidus | 316 |
| 139 | <i>Mathevotaenia surinamensis</i> | Dasypus novemcinctus | 62 |
| 140 | <i>Mediorhynchus grande</i> | Sturnus vulgaris | 75 |
| 141 | <i>Mediorhynchus robustum</i> | Sturnus vulgaris | 75 |
| 142 | <i>Mesocestoides</i> | Basariscus astutus | 362 |
| | | Conepatus mesoleucus | 466 |
| | | Mephitis mephitis | 466 |
| | | Spilogale putorius | 466 |
| 143 | <i>Mesocestoides corti</i> | Canis latrans | 364,365 |
| | | Felis rufus | 365,440 |
| | | Wild Canids | 122 |
| 144 | <i>Mesocestoides latus</i> | Didelphis virginiana | 97 |
| | | Opossum | 92 |
| 145 | <i>Mesocestoides lineatus</i> | Canis latrans | 370 |
| 146 | <i>Mesocestoides litteratus</i> | Urocyon cinereoargenteus | 65 |
| 147 | <i>Mesocestoides variabilis</i> | Procyon lotor | 93 |
| | | Raccoon | 92 |
| 148 | <i>Mesocoelium americanum</i> | Eumeces fasciatus | 216 |
| 149 | <i>Metroliasthes lucida</i> | Meleagris gallopavo intermedia | 361 |
| 150 | <i>Microsomacanthus hopkinsi</i> | Anas clypeata | 58 |
| | | Anas cyanoptera septentrionalium | 499 |
| 151 | <i>Moniezia</i> | Cattle | 138A |
| 152 | <i>Moniezia benedeni</i> | Peccary | 2 |
| | | Tayassu tajacu | 409 |
| | | White-collared Peccary | 415 |
| 153 | <i>Moniezia expansa</i> | Antilocapra americana | 207N |
| | | Sheep | 378 |
| 154 | <i>Moniezia planissima</i> | Cattle | 378 |
| 155 | <i>Monoecocestus sigmodontis</i> | Sigmodon hispidus | 250,298,316, 319,328 |
| 156 | <i>Mosgovoyia pectinata</i> | Lepus californicus | 404N |
| 157 | <i>Multiceps serialis</i> | Dog | 378 |
| | | Rabbit | 378 |
| 158 | <i>Oochoristica</i> | Conepatus mesoleucus | 466 |
| | | Cophosaurus texanus | 486A |
| | | Holbrookia maculata | 486A |
| | | Mephitis mephitis | 466 |
| | | Spilogale putorius | 466 |
| 159 | <i>Oochoristica americana</i> | Farancia abacura | 216 |
| 160 | <i>Oochoristica anolis</i> | Anolis carolinensis | 216 |
| 161 | <i>Oochoristica</i> <i>bivitellolobata</i> | Cnemidophorus tessellatus | 206N |
| 162 | <i>Oochoristica elaphis</i> | Elaphe obsoleta lindheimerii | 216 |
| 163 | <i>Oochoristica eumecis</i> | Eumeces fasciatus | 216 |

| | | | |
|-----|--------------------------------------|------------------------------------|----------------|
| 164 | <i>Oochoristica loeweni</i> | <i>Cnemidophorus tessellatus</i> | 206N |
| 165 | <i>Oochoristica mesillensis</i> | <i>Cophosaurus texanus</i> | 206N |
| 166 | <i>Oochoristica natricis</i> | <i>Nerodia rhombifera</i> | 216 |
| 167 | <i>Oochoristica neo-mexicana</i> | <i>Cnemidophorus tessellatus</i> | 206N |
| | | <i>Cophosaurus texanus</i> | 206N |
| 168 | <i>Oochoristica parvovaria</i> | <i>Phrynosoma douglassi</i> | 486A |
| 169 | <i>Oochoristica procyonis</i> | <i>Procyon lotor</i> | 93 |
| 170 | <i>Oochoristica rattii</i> | <i>Rattus rattus</i> | 250 |
| 171 | <i>Oochoristica scelopori</i> | <i>Sceloporus magister</i> | 486A |
| 172 | <i>Ophiotaenia frigidus</i> | <i>Ptychocheilus lucius</i> | 64A |
| 173 | <i>Ophiotaenia magna</i> | <i>Rana catesbeiana</i> | 233 |
| | | <i>Rana pipiens complex</i> | 425 |
| 174 | <i>Ophiovalipora</i> | <i>Gambusia affinis</i> | 128 |
| 175 | <i>Ophryocotyle proteus</i> | <i>Rallus longirostris</i> | 37 |
| 176 | <i>Oxyspirura petrowi</i> | <i>Sturnus vulgaris</i> | 75 |
| 177 | <i>Paricterotaenia</i> | <i>Colinus virginianus texanus</i> | 491 |
| | | <i>Rallus longirostris</i> | 37 |
| 178 | <i>Paricterotaenia parina</i> | <i>Sturnus vulgaris</i> | 43 |
| 179 | <i>Paricterotaenia pauciannulata</i> | <i>Chordeiles minor minor</i> | 495 |
| 180 | <i>Philotaenia filaroides</i> | <i>Ambystoma tigrinum</i> | 505N |
| 181 | Plerocercoid larva | <i>Fundulus grandis</i> | 24 |
| | | <i>Ictiobus bubalus</i> | 421 |
| 182 | Plerocercoid-proteocephalid | <i>Dorosoma cepedianum</i> | 241 |
| 183 | <i>Prochristianella penaei</i> | <i>Penalus aztecus</i> | 1 |
| | | <i>Penalus setiferus</i> | 1 |
| 184 | <i>Proteocephalidae</i> | <i>Ambloplites rupestris</i> | 479 |
| | | <i>Cichlosoma cyanoguttatum</i> | 479 |
| | | <i>Ictalurus melas</i> | 479 |
| | | <i>Ictalurus natalis</i> | 479 |
| | | <i>Lepomis auritus</i> | 479 |
| | | <i>Lepomis gulosus</i> | 479 |
| | | <i>Lepomis macrochirus</i> | 479 |
| | | <i>Lepomis microlophus</i> | 479 |
| | | <i>Lepomis punctatus</i> | 479 |
| | | <i>Micropterus punctulatus</i> | 479 |
| | | <i>Micropterus salmoides</i> | 479 |
| 185 | <i>Proteocephalus</i> | <i>Anolis carolinensis</i> | 216 |
| | | <i>Fundulus grandis</i> | 24 |
| | | <i>Gambusia affinis</i> | 128 |
| | | <i>Ictalurus melas</i> | 40 |
| | | <i>Lepomis macrochirus</i> | 40 |
| | | <i>Lepomis marginatus</i> | 40 |
| | | <i>Micropterus salmoides</i> | 40 |
| 186 | <i>Proteocephalus ambloplitis</i> | <i>Ictalurus natalis</i> | 479 |
| | | <i>Ictalurus punctatus</i> | 211 |
| | | <i>Lepisosteus</i> | 109 |
| | | <i>Lepisosteus oculatus</i> | 479 |
| | | <i>Lepisosteus osseus</i> | 76 |
| | | <i>Lepisosteus productus</i> | 76 |
| | | <i>Lepisosteus spatula</i> | 76 |
| | | <i>Lepomis cyanellus</i> | 20, 308, 309 |
| | | <i>Lepomis gulosus</i> | 211 |
| | | <i>Lepomis humilis</i> | 308 |
| | | <i>Lepomis macrochirus</i> | 211, 308, 309, |

| | | |
|-----|---|---|
| | | 310 |
| | Lepomis megalotis | 20, 211, 308, 309, 310 |
| | Lepomis microlophus | 211 |
| | Micropterus punctulatus | 211, 479 |
| | Micropterus salmoides | 20, 23, 252, 309, 310, 479 |
| | Pomoxis annularis | 20, 211, 309, 310 |
| | Lepomis macrochirus | 280 |
| | Micropterus salmoides | 280 |
| 187 | <i>Proteocephalus ambloplitis</i> plerocercoid | Lepomis macrochirus 315 |
| | Lepomis megalotis | 315 |
| | Micropterus salmoides | 315 |
| | Pomoxis annularis | 315 |
| 188 | <i>Proteocephalus faranciae</i> | Farancia abacura 216 |
| 189 | <i>Proteocephalus magnus</i> | Rana catesbeiana 216 |
| | Rana clamitans | 216 |
| 190 | <i>Pseudolytoeceslus differtus</i> | Ictiobus bubalus 421 |
| 191 | <i>Raillietina</i> | Perognathus hispidus 316 |
| | Sigmodon hispidus | 316 |
| 192 | <i>Raillietina bakeri</i> | Sciurus niger 78 |
| | Sigmodon hispidus | 250, 298, 328 |
| 193 | <i>Raillietina cesticillus</i> | Colinus virginianus texanus 491 |
| 194 | <i>Raillietina colinia</i> | Colinus virginianus texanus 491 |
| 195 | <i>Raillietina klebergi</i> | Colinus virginianus texanus 492 |
| 196 | <i>Raillietina minuta</i> | Colinus virginianus texanus 492, 496 |
| 197 | <i>Raillietina retractilis</i> | Lepus californicus 404N |
| | Sylvilagus auduboni | 132N |
| 198 | <i>Raillietina stilesiella</i> | Sylvilagus floridanus 201 |
| 199 | <i>Raillientina williamsi</i> | Meleagris gallopavo intermedia 361 |
| 200 | <i>Rhabdometra odiosa</i> | Callipepla squamata 123 |
| | Canis latrans | 370 |
| | Colinus virginianus texanus | 491 |
| | Ortalis vetula maccalli | 105 |
| | Tympanuchus pallidicinctus | 363, 369 |
| 201 | <i>Rictularia lucifugus</i> | Myotis lucifugus 73N |
| 202 | <i>Schizotaenia americana</i> | Erethizon dorsatum 459A |
| 203 | <i>Schizotaenia sigmodontis</i> | Sigmodon hispidus 79 |
| 204 | <i>Scolex pleuronectis</i> | Fundulus grandis 24 |
| 205 | <i>Sobolevicanthus</i> | Dendrocygna autumnalis 205 |
| 206 | <i>Sobolevicanthus</i> <i>filumferens</i> | Anas clypeata 58 |
| | Anas cyanoptera septentrionalium | 499 |
| 207 | <i>Sobolevicanthus gracilis</i> | Anas creca 74 |
| 208 | <i>Sobolevicanthus</i> <i>krabbeella</i> | Anas creca 74 |
| 209 | <i>Sparganum mansoni</i> | Man 393 |
| 210 | <i>Spirometra</i> | Felis domesticus 199 |
| 211 | <i>Taenia</i> | Canis latrans 464 |
| | Dog | 430 |
| 212 | <i>Taenia hydatigena</i> | Dog 378 |
| 213 | <i>Taenia macrocystis</i> | Canis latrans 370 |
| | Felis rufus | 247 |
| | Wild Canids | 122 |
| 214 | <i>Taenia martis</i> | Bassariscus astutus 362 |
| 215 | <i>Taenia multiceps</i> | Canis latrans 370 |

| | | |
|------------------------------------|----------------------------|-------------|
| | Wild Canids | 122 |
| 216 <i>Taenia pisiformis</i> | Canis latrans | 364,365,370 |
| | Dog | 378,438 |
| | Felis rufus | 247 |
| | Sylvilagus auduboni | 132N |
| | Urocyon cinereoargenteus | 65 |
| | Wild Canids | 122 |
| 217 <i>Taenia rileyi</i> | Felis rufus | 247,365,440 |
| | Sigmodon hispidus | 238 |
| 218 <i>Taenia serialis</i> | Canis latrans | 364,365 |
| | Urocyon cinereoargenteus | 65 |
| 219 <i>Taenia taeniaeformis</i> | Cat | 378 |
| | Sigmodon hispidus | 328 |
| | Wild Canids | 122 |
| 220 <i>Thysanosoma actinioides</i> | Ammotragus lervia | 13N |
| | Antilocapra americana | 9N,18N,207N |
| | Cervus elaphus | 501N |
| | Domestic Sheep | 7N |
| | Goat | 378 |
| | Liposcellis bostrycophilus | 17N |
| | Odocoileus hemionus | 19A |
| | Ovis canadensis | 19A |
| | Phyopsocus | 17N,401A |
| | Ruminant | 12N |
| | Sheep | 4N,5N,378 |
| 221 <i>Variolepis farciminosa</i> | Sturnus vulgaris | 75 |
| 222 <i>Wyominia tetoni</i> | Ovis canadensis | 19A,401A |

Trematoda

| | | |
|--|--------------------------|-------------------|
| 223 <i>Acanthoparyphium spinulosum</i> | Oyster | 243 |
| 224 <i>Acolpenteron ureteroecetes</i> | Micropterus punctulatus | 40 |
| 225 <i>Actinocleidus</i> | Lepomis macrochirus | 40 |
| | Lepomis marginatus | 40 |
| | Lepomis megalotis | 40 |
| | Lepomis macrochirus | 308 |
| 226 <i>Actinocleidus articularis</i> | Lepomis macrochirus | 442 |
| | Lepomis megalotis | 20,310,314 |
| 227 <i>Actinocleidus fergusonii</i> | Lepomis cyanellus | 308 |
| | Lepomis humilis | 308 |
| | Lepomis macrochirus | 20,40,106,308,310 |
| | Lepomis megalotis | 308 |
| | Micropterus salmoides | 425 |
| 228 <i>Actinocleidus flagellatus</i> | Chaenobryttus gulosus | 40 |
| | Lepomis cyanellus | 20,310,315 |
| | Lepomis macrochirus | 20 |
| 229 <i>Actinocleidus furcatus</i> | Lepomis macrochirus | 40 |
| 230 <i>Actinocleidus fusiformes</i> | Chaenobryttus coronarius | 343 |
| | Lepomis cyanellus | 310,315 |
| | Lepomis megalotis | 309,314 |
| | Micropterus salmoides | 20,40,343 |

| | | | |
|-----|---|------------------------|--------------------|
| 231 | <i>Actinocleidus gracilis</i> | Chaenobryttus gulosus | 39 |
| | | Lepomis macrochirus | 40 |
| 232 | <i>Actinocleidus longus</i> | Lepomis cyanellus | 40,107,310, 343 |
| | | Lepomis macrochirus | 308 |
| 233 | <i>Actinocleidus maculatus</i> | Lepomis cyanellus | 308 |
| | | Lepomis humilis | 308 |
| | | Lepomis megalotis | 308 |
| 234 | <i>Actinocleidus unguis</i> | Lepomis marginatus | 40 |
| 235 | <i>Alaria</i> | Canis latrans | 364,365 |
| | | Dog | 378 |
| 236 | <i>Alaria americana</i> | Canis latrans | 464 |
| 237 | <i>Alaria marcianae</i> | Canis latrans | 364,365,370 |
| | | Felis rufus | 365,440 |
| 238 | <i>Allocreadium ictaluri</i> | Ictalurus melas | 40 |
| 239 | <i>Alloglossidium corti</i> | Ictalurus punctatus | 211 |
| 240 | <i>Alloglossidium geminus</i> | Ictalurus melas | 40 |
| 241 | <i>Alloglossidium kenti</i> | Ictalurus natalis | 40 |
| | | Ictalurus punctatus | 40 |
| 242 | <i>Amblyomma americanum</i> | Antilope cervicapra | 463 |
| 243 | <i>Amphimerus pseudofelineus</i> | Cat | 476 |
| | | Felis domesticus | 199 |
| 244 | <i>Analocreadium armatum</i> | Amnicola peracuta | 234 |
| | | No host given | 60 |
| 245 | <i>Ancyrocephalus felix</i> | Galeichtys felis | 418 |
| 246 | <i>Apatemon gracilis</i> | Dendrocygna autumnalis | 205 |
| 247 | <i>Armatae xiphidiocercaria</i> | Littorina irrorata | 185 |
| | | Cerithidea pliculosa | 185 |
| 248 | <i>Ascocotyle</i> | Adinia xenica | 24 |
| | | Fundulus grandis | 24 |
| | | Mollienesia formosa | 24 |
| | | Mollienesia latipinna | 24 |
| 249 | <i>Ascocotyle chandleri</i> | Adinia xenica | 24 |
| | | Cyprinodon variegatus | 419 |
| | | Fundulus grandis | 24 |
| | | Mollienesia latipinna | 24,419 |
| 250 | <i>Ascocotyle leighi</i> | Casmerodius albus | 433 |
| | | Cyprinodon variegatus | 433 |
| | | Mollienesia latipinna | 433 |
| 251 | <i>Ascocotyle leighi</i> <i>metacercaria</i> | Casmerodius albus | 433 |
| | | Cyprinodon variegatus | 433 |
| | | Mollienesia latipinna | 433 |
| 252 | <i>Ascocotyle mollienisicola</i> | Mollienesia formosa | 24 |
| | | Mollienesia latipinna | 24 |
| 253 | <i>Ascocotyle pachycystis</i> | Rallus elegans | 37 |
| | | Rallus longirostris | 37 |
| 254 | <i>Aspidogaster conchicola</i> | Amblema costata | 193 |
| | | Amblema perplicata | 200 |
| | | Amblema plicata | 193 |
| | | Anodonta corpulenta | 193,200 |
| | | Clam | 242 |
| | | Quadrula forsheyi | 200 |
| | | Quadrula houstonensis | 200 |
| 255 | <i>Athesmia heterolecithodes</i> | Canis latrans | 370 |
| | | Rallus elegans | 37 |

| | | | |
|-----|---|----------------------------------|-------------------|
| 256 | <i>Auridistomum chelydrae</i> | Chelydra serpentina | 216 |
| 257 | <i>Bilharziella polonica</i> | Anas cyanoptera septentrionalium | 499 |
| 258 | <i>Brachylaima thompsoni</i> | Sigmodon hispidus | 298 |
| 259 | <i>Brachylaime microti</i> | Microtus longicaudus | 277A |
| 260 | <i>Brachycoelium</i> | Opheodrys aestivus | 216 |
| 261 | <i>Brachycoelium daviesi</i> | Ambystoma texanum | 216 |
| | | Hyla cinerea | 216 |
| | | Pseudacris triseriata | 216 |
| | | Scincella lateralis | 216 |
| 262 | <i>Brachycoelium hospitale</i> | Rana sphenoccephala | 216 |
| 263 | <i>Brachycoelium meridionalis</i> | Notophthalmus meridionalis | 216 |
| 264 | <i>Brachycoelium storeriae</i> | Storeria dekayi | 216 |
| 265 | <i>Brachylaemus virginianus</i> | Armadillo | 96 |
| 266 | <i>Brachylecithum americanum</i> | Cassidix mexicanus | 137 |
| | | Corvus brachyrhynchos | 137 |
| | | Cyanocitta cristata | 137 |
| | | Quiscalus versicolor | 137 |
| | | Sturnella magna | 137 |
| 267 | <i>Brevifurcate apharyngeate cercaria</i> | Cerithidea pliculosa | 185 |
| 268 | <i>Bucephalid larva</i> | Donax | 237 |
| | | Ostrea equestris | 237 |
| 269 | <i>Bucephaliodes strongyluræ</i> | Menidia beryllina | 238 |
| | | Strongylura marina | 238 |
| 270 | <i>Bucephalus elegans</i> | Lampsilis anodontoides | 193 |
| | | Lampsilis radiata | 193 |
| | | Lepomis cyanellus | 308 |
| | | Lepomis humilis | 308 |
| | | Lepomis macrochirus | 308 |
| | | Lepomis megalotis | 308 |
| | | Proptera purpurata | 193 |
| 271 | <i>Bucephalus loeschi</i> | Donax variabilis | 239 |
| 272 | <i>Caecincola cooperi</i> | Lepomis macrochirus | 40 |
| 273 | <i>Caecincola latostoma</i> | Micropterus punctulatus | 479 |
| | | Micropterus salmoides | 479 |
| 274 | <i>Caecincola parvulus</i> | Chaenobryttus gulosus | 40 |
| | | Huro salmoides | 434 |
| | | Lepomis cyanellus | 309, 310, 314, |
| | | Micropterus dolomieu | 40 |
| | | Micropterus punctulatus | 40, 211 |
| | | Micropterus salmoides | 20, 40, 211, 252, |
| | | | 309 |
| | | Pomoxis annularis | 40 |
| 275 | <i>Carneophallus</i> | Rallus longirostris | 37 |
| 276 | <i>Cercaria flexicorpa</i> | Lepomis cyanellus | 308 |
| | | Lepomis humilis | 308 |
| | | Lepomis macrochirus | 308 |
| | | Lepomis megalotis | 308 |
| 277 | <i>Cercaria rabbi</i> | Musculium transversum | 153 |
| 278 | <i>Cercaria rangiae</i> | Rangia cuneata | 489 |
| 279 | <i>Cercaria ruddi</i> | Snail | 153 |
| 280 | <i>Cercaria tageli</i> | Tagelus plebeius | 489 |
| 281 | <i>Cercariaeum cercaria</i> | Melamysus bidentatus | 185 |
| 282 | <i>Cercorchis bairdi</i> | Sternotherus carinatus | 216 |
| | | Pseudemys scripta elegans | 216 |
| 283 | <i>Cercorchis texanus</i> | Pseudemys scripta elegans | 216 |

| | | | |
|-----|----------------------------------|------------------------------------|------------|
| 284 | <i>Choanura cercaria</i> | <i>Donax variabilis</i> | 239 |
| 285 | <i>Cleidodiscus</i> | <i>Lepomis cyanellus</i> | 20 |
| 286 | <i>Cleidodiscus bedardi</i> | <i>Lepomis cyanellus</i> | 20 |
| | | <i>Lepomis macrochirus</i> | 20 |
| | | <i>Lepomis marginatus</i> | 40 |
| | | <i>Lepomis megalotis</i> | 20,40,310 |
| 287 | <i>Cleidodiscus chelatus</i> | <i>Lepomis marginatus</i> | 40 |
| | | <i>Lepomis megalotis</i> | 40 |
| 288 | <i>Cleidodiscus diversis</i> | <i>Lepomis cyanellus</i> | 20,40,308 |
| | | <i>Lepomis humilis</i> | 308 |
| | | <i>Lepomis macrochirus</i> | 308 |
| | | <i>Lepomis megalotis</i> | 308 |
| 289 | <i>Cleidodiscus floridanus</i> | <i>Ictalurus punctatus</i> | 280 |
| 290 | <i>Cleidodiscus globus</i> | <i>Lepomis cyanellus</i> | 21 |
| 291 | <i>Cleidodiscus longus</i> | <i>Lepomis macrochirus</i> | 280 |
| | | <i>Micropterus salmoides</i> | 280 |
| | | <i>Pomoxis annularis</i> | 20,40,280 |
| | | | 310 |
| 292 | <i>Cleidodiscus nematocirrus</i> | <i>Lepomis macrochirus</i> | 20 |
| | | <i>Lepomis megalotis</i> | 30,310 |
| 293 | <i>Cleidodiscus pricei</i> | <i>Ameiurus natalis</i> | 343 |
| | | <i>Ictalurus punctatus</i> | 40,107,280 |
| 294 | <i>Cleidodiscus robustus</i> | <i>Lepomis cyanellus</i> | 308,343 |
| | | <i>Lepomis humilis</i> | 308 |
| | | <i>Lepomis macrochirus</i> | 40,308,310 |
| 295 | <i>Cleidodiscus similis</i> | <i>Lepomis cyanellus</i> | 21 |
| 296 | <i>Cleidodiscus uniformis</i> | <i>Pomoxis annularis</i> | 20,310 |
| 297 | <i>Cleidodiscus vanclideanus</i> | <i>Pomoxis annularis</i> | 20,40,310 |
| | | | 343 |
| | | <i>Pomoxis nigromaculatus</i> | 40 |
| 298 | <i>Clinostomum</i> | <i>Lepisosteus osseus</i> | 76 |
| | | <i>Lepisosteus spatula</i> | 76 |
| 299 | <i>Clinostomum marginatum</i> | <i>Chaenobryttus gulosus</i> | 40 |
| | | <i>Huro salmoides</i> | 434 |
| | | <i>Ictalurus punctatus</i> | 211 |
| | | <i>Lepomis auritus</i> | 479 |
| | | <i>Lepomis macrochirus</i> | 479 |
| | | <i>Lepomis microlophus</i> | 479 |
| | | <i>Lepomis punctatus</i> | 479 |
| | | <i>Micropterus punctulatus</i> | 40 |
| | | <i>Micropterus salmoides</i> | 40,252,479 |
| | | <i>Pimephales vigilax</i> | 40 |
| | | <i>Pomoxis annularis</i> | 280 |
| 300 | <i>Cornucopula adunca</i> | <i>Balistes carolinensis</i> | 235 |
| 301 | <i>Cotylapsis insignis</i> | <i>Anodonta corpulenta</i> | 193,200 |
| | | Clam | 242 |
| | | <i>Lampsilis anodontoides</i> | 193,200 |
| | | <i>Lampsilis radiata</i> | 193 |
| | | <i>Proptera purpurata</i> | 193 |
| | | <i>Quadrula forsheyi</i> | 200 |
| | | <i>Quadrula houstonensis</i> | 200 |
| 302 | <i>Crassicutis archosargii</i> | <i>Archosargus probatocephalus</i> | 358 |
| 303 | <i>Crassiphiala bulboglossa</i> | <i>Gambusia affinis</i> | 127,128 |
| 304 | <i>Crepidostomum</i> | <i>Gambusia affinis</i> | 128 |
| 305 | <i>Crepidostomum cooperi</i> | <i>Ambloplites rupstris</i> | 479 |
| | | <i>Lepomis auritus</i> | 479 |

| | | |
|-----|--|--|
| | Lepomis cyanellus | 20,309,310,315 |
| | Lepomis gulosus | 479 |
| | Lepomis macrochirus | 20,309,479 |
| | Lepomis marginatus | 40 |
| | Lepomis megalotis | 20,40,309,310, 314 |
| | Lepomis microlophus | 40,479 |
| | Lepomis punctatus | 479 |
| | Micropterus salmoides | 20,309,479 |
| | Pomoxis annularis | 211 |
| | Salmo gilae | 340A |
| 306 | <i>Crepidostomum cornutum</i> <i>cercaria</i> | Crayfish 339 |
| 307 | <i>Crepidostomum farionis</i> | Salmo apache Samo gairdneri Salmo trutta 340A 497 340A,497 |
| 308 | <i>Crepidostomum ictaluri</i> | Ictalurus punctatus 211 |
| 309 | <i>Crepidostomum illinoiense</i> | Hiodon alosoides 419 |
| 310 | <i>Creptotrematina aguirre</i> <i>pequenoii</i> | Astyanax mexicanus 479 |
| 311 | <i>Cyclocoelum mutabile</i> | Capella gallinago 287 |
| 312 | <i>Cyclocoelum obscurum</i> | Numenius americanus 146 Rallus elegans 37 |
| 313 | <i>Cylicostomum</i> | Horse Mule 378 378 |
| 314 | <i>Cysticercus</i> | Eumeces fasciatus 216 |
| 315 | <i>Dactylogyrus anchoratus</i> | Crassius auratus 343 |
| 316 | <i>Dactylogyrus auratus</i> | Notemigonus chrysoleucas 343 |
| 317 | <i>Dactylogyrus aureus</i> | Notemigonus chrysoleucas 40 |
| 318 | <i>Dactylogyrus extensus</i> | Cyprinus carpio 343 |
| 319 | <i>Dactylogyrus parvicirrus</i> | Notemigonus chrysoleucas 343 |
| 320 | <i>Dactylogyrus perlus</i> | Notropis lutrensis 107 |
| 321 | <i>Dactylogyrus vastator</i> | Crassius auratus 343 |
| 322 | <i>Dactylogyrus wegeneri</i> | Crassius auratus 343 |
| 323 | <i>Dasymetra conferta</i> | Nerodia fasciata 216 |
| 324 | <i>Dasymetra conferta</i> <i>metacercaria</i> | Acris crepitans 49 |
| 325 | <i>Dasymetra villicaeca</i> | Nerodia erythrogaster transversa 109 |
| 326 | <i>Dendritobilharzia</i> <i>pulverulenta</i> | Anas clypeata 55 Anas creca 74 |
| 327 | <i>Dichotoma cercaria</i> | Donax variabilis 239 |
| 328 | <i>Dicrocoelium rileyi</i> | Tadarida brasiliensis 71N,254 |
| 329 | <i>Dictyocaulus arnfieldi</i> | Horse Mule 378 378 |
| 330 | <i>Diplomonorchis leiostomi</i> | Leiostomus xanthurus 236 Micropogon undulatus 262 |
| 331 | <i>Diplostomatidae</i> | Lepomis auritus 479 Lepomis gulosus 479 Lepomis macrochirus 479 Lepomis punctatus 479 Micropterus salmoides 479 |
| 332 | <i>Diplostomulum scheuringi</i> | Gambusia affinis 127,128 |
| 333 | <i>Diplostomum</i> | Micropterus salmoides 252 |
| 334 | <i>Diplostomum crassum</i> | Quiscalus versicolor 99 |
| 335 | <i>Echinochasmus schwartzi</i> | Rallus elegans 37 |

| | | |
|---------------------------------------|----------------------------------|-------------------------------|
| | Rallus longirostris | 37 |
| 336 <i>Echinoparyphium flexum</i> | Duck | 68 |
| | Gadwall | 69 |
| | Pintail | 69 |
| | Shoveler | 69 |
| 337 <i>Echinoparyphium recurvatum</i> | Anas clypeata | 58 |
| | Phasianus colchicus | 366 |
| 338 <i>Echinostoma attenuatum</i> | Rallus elegans | 37 |
| | Rallus longirostris | 37 |
| 339 <i>Echinostoma revolutum</i> | Anas clypeata | 58 |
| | Anas crecca | 74 |
| | Anas cyanoptera septentrionalium | 499 |
| | Capella gallinago | 287 |
| | Dendrocygna autumnalis | 183 |
| | Duck | 68 |
| | Gadwall | 69 |
| | Pintail | 69 |
| | Shoveler | 69 |
| 340 <i>Echinostome cercaria</i> | Cerithidea pliculosa | 185 |
| | Nassarius vibex | 185 |
| 341 <i>Echinostome metacercaria</i> | Crayfish | 337 |
| 342 <i>Eurytrema allentoshi</i> | Didelphis virginiana | 135 |
| 343 <i>Eurytrema procyonis</i> | Procyon lotor | 93,134,136 |
| | Wild Canids | 122 |
| 344 <i>Fasciola hepatica</i> | Cattle | 116,138A,194, 197,378,502N |
| | Feral Hog | 197 |
| | Goat | 378 |
| | Lepus californicus | 345 |
| | Physa frontinalis acuta | 53 |
| | Sheep | 378,502N |
| | Sylvilagus floridanus | 345 |
| | White-tailed Deer | 197 |
| 345 <i>Fasciola magna</i> | Cattle | 197,378 |
| | Feral Hog | 197 |
| | White-tailed Deer | 197 |
| 346 <i>Fascioloides magna</i> | Cattle | 116,194 |
| | Cow | 424 |
| | Feral Hog | 194 |
| | Odocoileus virginianus | 194,208 |
| | Sus scrofa | 84 |
| | Tayassu tajacu | 409 |
| 347 <i>Fibricola texensis</i> | Procyon lotor | 89,93 |
| 348 <i>Galactosomum fregata</i> | Fregata magnificens | 103 |
| 349 <i>Genolopa</i> | Fundulus grandis | 24 |
| 350 <i>Glypthelmins quieta</i> | Rana catesbeiana | 233,425 |
| | Rana pipiens complex | 425 |
| 351 <i>Glypthelmins subtropica</i> | Rana catesbeiana | 216 |
| | Rana sphenoccephala | 216 |
| 352 <i>Gorgoderina minima</i> | Rana catesbeiana | 233 |
| 353 <i>Gorgoderina attenuata</i> | Rana catesbeiana | 233 |
| 354 <i>Gymnophalline cercaria</i> | Donax variabilis | 239 |
| 355 <i>Gyrodactylus</i> | Gambusia affinis | 40 |
| | Ictalurus melas | 40 |
| | Ictalurus natalis | 40 |
| 356 <i>Gyrodactylus elegans</i> | Crassius auratus | 343 |

| | | | |
|-----|------------------------------------|--|-------------------|
| 357 | <i>Haematoloechus</i> | <i>Rana catesbeiana</i> | 267,337 |
| | | <i>Rana pipiens complex</i> | 337 |
| 358 | <i>Haematoloechus breviplexus</i> | Amphibian | 339 |
| | | <i>Ferrissia</i> | 478,482 |
| | | <i>Rana catesbeiana</i> | 144N,233,425, 478 |
| | | <i>Rana pipiens complex</i> | 144N,425 |
| | | <i>Rana sphenoccephala</i> | 478 |
| 359 | <i>Haematoloechus coloradensis</i> | Anax | 143N, |
| | | <i>Enallagma</i> | 143N |
| | | <i>Libellula</i> | 143N |
| | | <i>Physa virgata</i> | 143N |
| | | <i>Rana pipiens complex</i> | 143N |
| | | <i>Tramea</i> | 143N |
| 360 | <i>Haematoloechus floedae</i> | <i>Rana catesbeiana</i> | 216,253 |
| | | <i>Rana clamitans</i> | 216 |
| 361 | <i>Haematoloechus longiplexus</i> | Amphibian | 339 |
| 362 | <i>Haematoloechus medioplexus</i> | Amphibian | 339 |
| 363 | <i>Haematoloechus uniplexus</i> | <i>Rana sphenoccephala</i> | 216 |
| 364 | <i>Halipegus</i> | <i>Lepomis gulosus</i> | 479 |
| | | <i>Lepomis microlophus</i> | 479 |
| | | <i>Lepomis punctatus</i> | 479 |
| 365 | <i>Haplocleidus</i> | <i>Gambusia affinis</i> | 40 |
| | | <i>Lepomis macrochirus</i> | 40 |
| 366 | <i>Haplocleidus biramosus</i> | <i>Lepomis megalotis</i> | 40 |
| 367 | <i>Haplocleidus dispar</i> | <i>Lepomis cyanellus</i> | 40 |
| | | <i>Lepomis macrochirus</i> | 40 |
| | | <i>Lepomis microlophus</i> | 40 |
| | | <i>Micropterus salmoides</i> | 40 |
| 368 | <i>Haplocleidus furcatus</i> | <i>Lepomis cyanellus</i> | 40 |
| | | <i>Lepomis marginatus</i> | 40 |
| | | <i>Micropterus salmoides</i> | 40 |
| 369 | <i>Harmostomum opisthotrias</i> | <i>Didelphis virginiana</i> | 83 |
| 370 | <i>Hasstilesia texensis</i> | <i>Lepus californicus</i> | 80 |
| | | <i>Sylvilagus floridanus</i> | 80 |
| 371 | <i>Hasstilesia tricolor</i> | Rabbit | 378 |
| 372 | <i>Hedruris</i> | <i>Lepomis microlophus</i> | 40 |
| | | <i>Pimephales vigilax</i> | 40 |
| 373 | <i>Hemiuridae juvenile</i> | <i>Dorosoma petenese</i> | 246 |
| 374 | <i>Henotosoma chelydrae</i> | <i>Chelydra serpentina</i> | 216 |
| | | <i>Kinosternon subrubrum hippocrepis</i> | 216 |
| | | <i>Pseudemys scripta elegans</i> | 216 |
| 375 | <i>Heterobilharzia americana</i> | <i>Lymnaea cubensis</i> | 209 |
| | | <i>Procyon lotor</i> | 379 |
| | | Raccoon | 209 |
| | | Wild Canids | 122 |
| 376 | <i>Heterophyid</i> | <i>Micropterus salmoides</i> | 309 |
| 377 | <i>Himasthla rhigedana</i> | <i>Numenius americanus</i> | 146 |
| 378 | <i>Homalometron armatum</i> | <i>Aplodinotus grunniens</i> | 211 |
| | | <i>Gambusia affinis</i> | 40 |
| | | <i>Lepomis macrochirus</i> | 211 |
| | | <i>Lepomis marginatus</i> | 40 |
| | | <i>Lepomis megalotis</i> | 20,309,310,315 |
| | | <i>Lepomis microlophus</i> | 211,479 |
| | | <i>Lepomis punctatus</i> | 479 |
| | | <i>Micropterus salmoides</i> | 211 |

| | | |
|--|----------------------------------|---------------|
| | Notropis lutrensis | 40 |
| 379 <i>Homalometron pallidum</i> | Fundulus grandis | 24 |
| | Fundulus similis | 24 |
| 380 <i>Hypoderaeum conoideum</i> | Anas creca | 74 |
| 381 <i>Immature digenetic fluke</i> | Carpiodes carpio | 420 |
| 382 <i>Laterocotyle padreinsulae</i> | Pogonis cromis | 423 |
| 383 <i>Lechriorchis validus</i> | Heterodon platyrhinos | 216 |
| | Lampropeltis getulus holbrooki | 216 |
| 384 <i>Leucochloridium certhiae</i> | Sturnus vulgaris | 43 |
| 385 <i>Levinseniella byrdi</i> | Rallus elegans | 37 |
| | Rallus longirostris | 37 |
| 386 <i>Lissorchis gullaris</i> | Ictiobus bubalus | 421 |
| | Ictiobus cyprinellus | 421 |
| | Ictiobus niger | 421 |
| 387 <i>Lissorchis hypentelii</i> | Moxostoma congestum | 479 |
| 388 <i>Lobatostoma</i> | Donax variabilis | 239 |
| 389 <i>Lyperosomum oswaldoi</i> | Numenius americanus | 146 |
| 390 <i>Lyperosomum sinuosum</i> | Numenius americanus | 146 |
| 391 <i>Macravestibulum</i> | Pseudemys concinna texana | 256 |
| | Pseudemys scripta troostii | 256 |
| 392 <i>Macroderoides spiniferis</i> | Gambusia affinis | 127,128 |
| 393 <i>Macrohaptor hopkinsi</i> | Lepomis macrochirus | 40 |
| | Lepomis marginatus | 40 |
| | Lepomis megalotis | 21 |
| 394 <i>Macrovalvitrematoides micropogoni</i> | Micropogon undulatus | 262 |
| 395 <i>Manodistomum occultum</i> | Notophthalmus meridionalis | 216 |
| 396 <i>Maritrema arenaria</i> | Numenius americanus | 146 |
| 397 <i>Mazocraeoides olentangiensis</i> | Dorosoma cepedianum | 241 |
| 398 <i>Megalodiscus americanus</i> | Rana sphenoccephala | 216 |
| 399 <i>Megalodiscus temperatus</i> | Hyla cinerea | 282 |
| | Pseudacris triseriata | 216 |
| | Rana areolata | 216 |
| | Rana catesbeiana | 216, 233, 425 |
| | Rana clamitans | 216 |
| | Rana pipiens complex | 425 |
| | Rana sphenoccephala | 216 |
| 400 <i>Megalurous cercaria</i> | Littorina irrorata | 185 |
| | Thais haemastoma | 185 |
| 401 <i>Megaperid cercaria</i> | Crepidula plana | 185 |
| 402 <i>Megnacercous cercaria</i> | Cerithidea pliculosa | 185 |
| 403 <i>Mesocoelium americanum</i> | Notophthalmus meridionalis | 216 |
| | Scincella lateralis | 216 |
| | Storeria dekayi | 216 |
| 404 <i>Mesostephorus longisaccus</i> | Dog | 102 |
| 405 <i>Metacercaria</i> | Acris crepitans | 49 |
| | Donax variabilis | 239 |
| 406 <i>Microcreadium parvum</i> | Amnicola peracuta | 234 |
| | No host given | 60 |
| 407 <i>Nematobothrium texomensis</i> | Ictiobus bubalus | 22A |
| | Ictiobus cyprinellus | 22A |
| | Ictiobus niger | 22A |
| 408 <i>Neogogatea pandionis</i> | Pandion haliaetus carolinensis | 99 |
| 409 <i>Neorenifer aniarum</i> | Nerodia erythrogaster transversa | 119 |
| 410 <i>Notocotylus attenuatus</i> | Anas clypeata | 58 |

| | | |
|-----|---|-----------------------------------|
| | Anas creca | 74 |
| | Anas cyanoptera septentrionalium | 499 |
| | Duck | 68 |
| | Gadwall | 69 |
| | Pintail | 69 |
| | Shoveler | 69 |
| 411 | <i>Notocotylus regis</i> | Rallus elegans 37 |
| 412 | <i>Ochotrenatrema labda</i> | Tadarida brasiliensis 71N,254 |
| 413 | <i>Oncocleidus accuminatus</i> | Lepomis megalotis 40 |
| 414 | <i>Oncocleidus acuminatus</i> | Lepomis marginatus 40 |
| 415 | <i>Oncocleidus cyanellus</i> | Lepomis cyanellus 40 |
| 416 | <i>Oncocleidus ferox</i> | Chaenobryttus gulosus 40 |
| | | Lepomis macrochirus 40 |
| | | Lepomis marginatus 40 |
| 417 | <i>Oncocleidus principalis</i> | Micropterus punctulatus 40 |
| | | Micropterus salmoides 40 |
| 418 | <i>Opecoeloides polyfimbriatus</i> | Synodus foetens 387 |
| 419 | <i>Opecoeloides polynemi</i> | Polynemus octonemus 387 |
| 420 | <i>Ophthalmophagus</i> | Rallus longirostris 37 |
| 421 | <i>Orchipedum jolliei</i> | Grus canadensis 67,198 |
| 422 | <i>Ornithodiplostomum ptychocheilus</i> | Gambusia affinis 127,128 |
| 423 | <i>Paralecithodendrium</i> | Tadarida brasiliensis 71N |
| 424 | <i>Paramacroderoides echinus</i> | Lepisosteus 109 |
| 425 | <i>Paramphistomum cervi</i> | Cattle 378 |
| 426 | <i>Paramphistomum stunkardi</i> | Huro salmoides 434 |
| 427 | <i>Parapleurolophocercous cercaria</i> | Cerithidea pliculosa 185 |
| 428 | <i>Paratrema numenii</i> | Numenius americanus 146 |
| 429 | <i>Parorchis acanthus</i> | Numenius americanus 146 |
| 430 | <i>Parvatrema donacis</i> | Donax variabilis 239 |
| 431 | <i>Pelmatostomum americanum</i> | Numenius americanus 146 |
| 432 | <i>Phagicola lageniformis</i> | Ondatra zibethicus 85 |
| 433 | <i>Pharyngostomoides procyonis</i> | Procyon lotor 93,214 |
| 434 | <i>Phyllodistomum</i> | Gambusia affinis 127,128 |
| | | Gambusia geiseri 127 |
| 435 | <i>Phyllodistomum caudatum</i> | Ictalurus melas 40 |
| | | Ictalurus natalis 40 |
| 436 | <i>Phyllodistomum lacustri</i> | Ictalurus punctatus 40 |
| 437 | <i>Phyllodistomum lohrenzi</i> | Huro salmoides 434 |
| | | Lepomis gulosus 479 |
| | | Micropterus salmoides 252,309,479 |
| | | Pomoxis annularis 40 |
| 438 | <i>Phyllodistomum pearsei</i> | Micropterus punctulatus 40 |
| | | Micropterus salmoides 40,252 |
| 439 | <i>Pisciamphistoma</i> | Micropterus salmoides 252 |
| 440 | <i>Pisciamphistoma minimum</i> | Lepomis megalotis 310 |
| 441 | <i>Pisciamphistoma reynoldsi</i> | Chaenobryttus gulosus 40 |
| | | Lepomis cyanellus 309,310,315 |
| | | Lepomis macrochirus 40 |
| | | Lepomis marginatus 40 |
| | | Lepomis megalotis 211,309,315 |
| | | Micropterus punctulatus 40 |
| | | Micropterus salmoides 20,40,309 |

| | | |
|--------------------------------------|-----------------------------------|---|
| | Pomoxis annularis | 20,309 |
| 442 <i>Pisciamphistoma stunkardi</i> | Lepomis cyanellus | 308 |
| | Lepomis humilis | 308 |
| | Lepomis macrochirus | 308 |
| | Lepomis megalotis | 308 |
| | Micropterus salmoides | 309 |
| 443 <i>Plagioporus sinitsini</i> | Dionda episcopa | 479 |
| | Gambusia affinis | 127,128 |
| | Notropis venustus | 479 |
| 444 <i>Plagiorchis</i> | Mormoops megalophylla | 254 |
| 445 <i>Plagiorchis goodmani</i> | No host given | 342 |
| 446 <i>Plagiorchis micracanthos</i> | Myotis lucifugus | 71N,73N |
| 447 <i>Pneumatophilus variabilis</i> | Nerodia erythrogaster transversa | 119 |
| 448 <i>Polylekithum ictaluri</i> | Amblema perplicata | 200 |
| | Anodonta corpulenta | 200 |
| | Lampsilis anodontoides | 200 |
| | Quadrula forsheyi | 200 |
| | Quadrula houstonensis | 200 |
| 449 <i>Polystoma hassalli</i> | Chelydra serpentina | 216 |
| | Kinosternon subrubrum hippocrepis | 216 |
| 450 <i>Polystoma megacotyle</i> | Pseudemys scripta elegans | 216 |
| 451 <i>Polystoma orbiculare</i> | Pseudemys scripta elegans | 216 |
| 452 <i>Polystoma terrapenis</i> | Terrapene carolina triunguis | 216 |
| 453 <i>Posthodiplostomum minimum</i> | Ambloplites rupestris | 479 |
| | Ardea herodias | 313 |
| | Astyanax mexicanus | 479 |
| | Chaenobryttus gulosus | 280,312 |
| | Cichlosoma cyanoguttatum | 479 |
| | Dionda episcopa | 479 |
| | Fundulus olivaceus | 312 |
| | Gambusia affinis | 40,127,128,312 |
| | Gambusia geiseri | 127 |
| | Hybognathus nuchalis | 40 |
| | Ictalurus melas | 312 |
| | Ictalurus natalis | 312 |
| | Ictalurus punctatus | 40,312 |
| | Lepomis auritus | 479 |
| | Lepomis cyanellus | 40,308,309,312 |
| | Lepomis gulosus | 211,479 |
| | Lepomis humilis | 308 |
| | Lepomis macrochirus | 40,211,280, 308,309,310, 212,479 |
| | Lepomis marginatus | 40 |
| | Lepomis megalotis | 40,211,308, 309,312 |
| | Lepomis microlophus | 40,211,312, 479 |
| | Lepomis punctatus | 479 |
| | Micropterus punctulatus | 211,311,479 |
| | Micropterus salmoides | 23,40,211, 252,280,309, 310,312,479 |
| | Minytrema melanops | 312 |
| | Moxostoma congestum | 312 |

| | | |
|---|----------------------------|---------|
| | Notemigonus chrysolecuas | 312 |
| | Notropis lutrensis | 40 |
| | Notropis umbratilis | 40 |
| | Notropis venustrus | 479 |
| | Percina | 40 |
| | Physa halei | 41,313 |
| | Pimephales vigilax | 40 |
| | Pomoxis annularis | 40,312 |
| | Pomoxis nigromaculatus | 312 |
| 454 <i>Posthodiplostomum minimum</i> <i>centrarchi</i> | Lepomis cyanellus | 20 |
| | Lepomis macrochirus | 20,315 |
| | Lepomis megalotis | 20,315 |
| | Micropterus salmoides | 20,315 |
| 455 <i>Postmonorchis orthopristsis</i> | Leiostomus xanthurus | 236 |
| | Orthopristsis chrysopterus | 236 |
| 456 <i>Proalaria variabilis</i> | Didelphis virginiana | 83 |
| 457 <i>Probolocorphyne glandulosa</i> | Numenius americanus | 146 |
| 458 <i>Prophyptiamus grusi</i> | Grus canadensis | 198 |
| 459 <i>Prosthogonimus</i> | Rallus longirostris | 37 |
| 460 <i>Prosthogonimus ovatus</i> | Rallus elegans | 37 |
| 461 <i>Protenes chapmani</i> | Pseudemys scripta elegans | 216 |
| 462 <i>Proteocephalus ambloplites</i> | Huro salmoides | 434 |
| 463 <i>Proterometra macrostoma</i> | Ambloplites rupestris | 479 |
| | Astyanax mexicanus | 479 |
| | Lepomis auritus | 479 |
| | Lepomis gulosus | 479 |
| | Lepomis macrochirus | 479 |
| | Lepomis microlophus | 479 |
| | Lepomis punctatus | 479 |
| | Micropterus salmoides | 479 |
| 464 <i>Pseudohaliotrema</i> <i>rhomboides</i> | Lagodon rhomboides | 418 |
| 465 <i>Pseudomagnivittellinum</i> <i>ictalurum</i> | Ictalurus melas | 147 |
| 466 <i>Pseudomurraytrema muelleri</i> | Catostomus commersonii | 343 |
| 467 <i>Raillietina bakeri</i> | Sciurus niger | 91 |
| 468 <i>Renifer aniarum</i> | Nerodia fasciata | 216 |
| 469 <i>Renifer kansensis</i> | Agkistrodon contortrix | 216 |
| | Sistrurus miliarius | 216 |
| 470 <i>Renifer texanus</i> | Heterodon platyrhinos | 216 |
| 471 <i>Rhabdosynochus</i> <i>rhabdosynochus</i> | Centropomus undecimalis | 418 |
| 472 <i>Rhamnocercus stichospinus</i> | Menticirrus littoralis | 418 |
| | Micropogon undulatus | 418 |
| 473 <i>Rhipidocotyle</i> | Etheostoma gracile | 40 |
| 474 <i>Rhipidocotyle angusticollis</i> | Sarder sardeu | 86 |
| 475 <i>Rhipidocotyle lepisostei</i> | Fundulus grandis | 24 |
| | Lepisosteus | 109 |
| | Lepisosteus osseus | 136,139 |
| | Lepisosteus spatula | 136,139 |
| 476 <i>Rhipidocotyle papillosa</i> | Gambusia affinis | 128 |
| 477 <i>Rhipidocotyle</i> <i>septapapillata</i> | Ambloplites rupestris | 479 |

| | | |
|-----|------------------------------------|--------------------------------------|
| | Chaenobryttus gulosus | 40 |
| | Elassoma zonatum | 40 |
| | Lepomis gulosus | 479 |
| | Lepomis marginatus | 40 |
| | Notropis umbratilis | 40 |
| 478 | <i>Rhipidocotyle transversale</i> | Menidia beryllina 238 |
| | | Strongylura marina 238 |
| 479 | <i>Rhopalias macracanthus</i> | Didelphis 388 |
| | | Didelphis virginiana 83 |
| 480 | <i>Saccocoelioides sogandaresi</i> | Adinia xenica 24 |
| | | Mollienisia latipinna 24,418 |
| 481 | <i>Schwartzitrema seamsteri</i> | Fregata magnificens 103 |
| 482 | <i>Sterrhurus texanus</i> | Sarder sardeu 86 |
| 483 | <i>Stomatrema pusilla</i> | Farancia abacura 216 |
| 484 | <i>Strigea eroliae</i> | Erolia melanotos 190 |
| 485 | <i>Tanaisia fedtschenoki</i> | Capella gallinago 287 |
| 486 | <i>Tergestia pectinata</i> | Bairdiella chrysur 235 |
| | | Cynoscion regalis 235 |
| | | Pomatomus saltatrix 235 |
| | | Trichiurus lepturus 235 |
| 487 | <i>Tetracotyle</i> | Gambusia affinis 127,128 |
| 488 | <i>Texanocotyle pogoniae</i> | Pogonias cromis 423 |
| 489 | <i>Textrema hopkinsi</i> | Lepomis punctatus 479 |
| | | Micropterus punctulatus 479 |
| | | Micropterus salmoides 145,479 |
| 490 | <i>Tremajoannes buckleyi</i> | Tadarida brasiliensis 71 |
| 491 | <i>Trematode</i> | Myotis velifer 254 |
| 492 | <i>Trichobilharzia</i> | Anas clypeata 58 |
| 493 | <i>Trichobilharzia physellae</i> | Anas acuta 368 |
| | | Anas creca 368 |
| | | Anas cyanoptera septentrionalium 499 |
| | | Anas discors 368 |
| | | Physa anatina 368 |
| 494 | <i>Trichocercous cercaria</i> | Anaschis avara 185 |
| 495 | <i>Tricholipeurus parallelus</i> | Antilope cervicapra 463 |
| 496 | <i>Typhlocoelum cucumerinum</i> | Anas clypeata 58 |
| 497 | <i>Typhlocoelum sisowi</i> | Anas creca 74 |
| 498 | <i>Ubiquita xiphidiocercaria</i> | Littorina irrorata 185 |
| 499 | <i>Urocleidus</i> | Huro salmoides 434 |
| 500 | <i>Urocleidus acer</i> | Lepomis macrochirus 20,40,310 |
| 501 | <i>Urocleidus acuminatus</i> | Lepomis megalotis 310 |
| 502 | <i>Urocleidus attenuatus</i> | Lepomis cyanellus 310 |
| | | Lepomis macrochirus 103,310 |
| | | Lepomis megalotis 40 |
| 503 | <i>Urocleidus chaenobryttus</i> | Chaenobryttus coronarius 343 |
| | | Chaenobryttus gulosus 40 |
| | | Lepomis cyanellus 308 |
| | | Lepomis humilis 308 |
| | | Lepomis macrochirus 308 |
| | | Lepomis megalotis 308 |
| 504 | <i>Urocleidus chrysops</i> | Lepibema chrysops 107 |
| 505 | <i>Urocleidus cyanellus</i> | Lepomis cyanellus 20,308,310 |
| | | Lepomis humilis 308 |
| | | Lepomis macrochirus 308 |
| | | Lepomis megalotis 308,310 |

| | | | |
|-----|-------------------------------|--------------------------|------------|
| 506 | <i>Urocleidus dispar</i> | Lepomis cyanellus | 308,310 |
| | | Lepomis humilis | 308 |
| | | Lepomis macrochirus | 20,308,310 |
| | | Lepomis megalotis | 20,308,310 |
| | | Micropterus salmoides | 20,310 |
| 507 | <i>Urocleidus ferox</i> | Lepomis cyanellus | 308 |
| | | Lepomis humilis | 308 |
| | | Lepomis macrochirus | 20,308,310 |
| | | Lepomis megalotis | 20,308 |
| 508 | <i>Urocleidus fundulus</i> | Fundulus notatus | 40 |
| 509 | <i>Urocleidus furcatus</i> | Micropterus salmoides | 20,23 |
| 510 | <i>Urocleidus grandis</i> | Chaenobryttus gulosus | 40 |
| | | Lepomis cyanellus | 20,40,310 |
| | | Lepomis megalotis | 310 |
| | | Pomoxis nigromaculatus | 107 |
| 511 | <i>Urocleidus helioides</i> | Micropterus salmoides | 343 |
| 512 | <i>Urocleidus principalis</i> | Lepomis cyanellus | 20 |
| | | Lepomis macrochirus | 310 |
| | | Micropterus punctulatus | 107 |
| | | Micropterus salmoides | 20 |
| 513 | <i>Urocleidus seculus</i> | Fundulus notatus | 40 |
| | | Gambusia affinis | 40 |
| | | Gambusia affinis affinis | 343 |
| 514 | <i>Urocleidus torquatus</i> | Lepomis macrochirus | 310 |
| | | Lepomis megalotis | 20,310 |
| 515 | <i>Urocleidus variabilis</i> | Lepomis marginatus | 40 |
| | | Lepomis megalotis | 40 |
| 516 | <i>Uvulifer ambloplitis</i> | Micropterus salmoides | 23,211 |
| 517 | <i>Uvulifer erraticus</i> | Toxostoma rufum | 99 |
| 518 | <i>Zonorchis allentoshi</i> | Canis latrans | 370 |
| 519 | <i>Zonorchis komareki</i> | Sigmodon hispidus | 298 |
| 520 | <i>Zonorchis petiolatus</i> | Corvus brachyrhynchos | 341 |
| 521 | <i>Zygocotyle lunata</i> | Anas clypeata | 58 |
| | | Anas creca | 74 |
| | | Dendrocygna autumnalis | 205 |
| | | Duck | 68 |
| | | Gadwall | 69 |
| | | Pintail | 69 |

Nemata

| | | | |
|-----|----------------------------------|--------------------------|---------------------|
| 522 | <i>Abbreviata</i> | Tadarida brasiliensis | 254N |
| 523 | <i>Acuaria anthuris</i> | Corvus brachyrhynchos | 341 |
| | | Corvus cryptoleucus | 371N |
| 524 | <i>Alifilaria pochardi</i> | Anas clypeata | 58 |
| 525 | <i>Amelostrogylus mentulatus</i> | Antilope cervicapra | 463 |
| 526 | <i>Amidostomum acutum</i> | Anas clypeata | 58 |
| | | Anas creca | 74 |
| 527 | <i>Amidostomum anseris</i> | Anas creca | 74 |
| 528 | <i>Ancylostoma braziliense</i> | Dog | 378 |
| | | Urocyon cinereoargenteus | 65 |
| 529 | <i>Ancylostoma caninum</i> | Canis familiaris | 217N |
| | | Canis latrans | 364,365,370, 464 |

| | | |
|-----|-----------------------------------|--|
| | Cat | 378 |
| | Dog | 223, 378, 430, 438 |
| | Felis rufus | 291 |
| | Urocyon cinereoargenteus | 65 |
| | Wild Canids | 122 |
| 530 | <i>Ancylostoma tubaeforme</i> | Felis rufus 365, 440 |
| 531 | <i>Anoplostrongylus delicatus</i> | Tadarida brasiliensis 254N |
| 532 | <i>Arduenna strongylina</i> | Pig 378 |
| 533 | <i>Arthrocephalus lotoris</i> | Bassariscus astutus 362 |
| 534 | <i>Ascarida columbae</i> | Columba livia 351 |
| 535 | <i>Ascaridia lineata</i> | Chicken 378 |
| 536 | <i>Ascaris lumbricoides</i> | Pig 378 |
| 537 | <i>Ascaris lumbricoides</i> | Pig 378 |
| 538 | <i>Ascaris suum</i> | Dicotyles tajacu 222 Sus scrofa 113 |
| 539 | <i>Ascarops larva</i> | Armadillo 96 |
| 540 | <i>Ascarops strongylina</i> | Myotis velifer 254 Onthophagus gazella 189 |
| 541 | <i>Aspiculuris</i> | Neotoma cinerea 274A Neotoma lepida 274A Neotoma stephensi 274A |
| 542 | <i>Aspiculuris ackerti</i> | Neotoma albigula 275A Neotoma cinerea 275A |
| 543 | <i>Aspidodera fasciata</i> | Armadillo 96 |
| 544 | <i>Asidodera harwood</i> | Didelphis virginiana 83 |
| 545 | <i>Atractis carolinae</i> | Terrapene carolina triunguis 216 |
| 546 | <i>Aulonocephalus lindquisti</i> | Callipepla squamata 123 Colinus virginianus texanus 491 |
| 547 | <i>Bunostomum</i> | Cattle 44 |
| 548 | <i>Bunostomum phlebotomum</i> | Cattle 378 Guinea Pig 50 Rabbit 50 Sheep 50 |
| 549 | <i>Bunostomum trigonocephalum</i> | Goat 378 Sheep 378 |
| 550 | <i>Camallanus</i> | Dorosoma petenese 241 Huro salmoides 434 Micropterus salmoides 309 |
| 551 | <i>Camallanus ancyloderus</i> | Ictiobus bubalus 421 Ictiobus cyprinellus 421 |
| 552 | <i>Camallanus oxycephalus</i> | Ambloplites rupestris 479 Ammocrypta vivax 479 Aplodinotus grunniens 211 Astyanax mexianus 479 Carpiodes carpio 420 Cichlosoma cyanoguttatum 479 Elassoma zonatum 40 Gambusia affinis 127, 128 Gambusia geiseri 127 Ictalurus punctatus 211 Lepomis auritus 479 Lepomis cyanellus 308, 309, 310, 479 Lepomis gulosus 211, 479 Lepomis humilis 308 |

| | | |
|--|-----------------------------------|---------------------------------------|
| | Lepomis macrochirus | 40,308,479 |
| | Lepomis megalotis | 308 |
| | Lepomis microlophus | 479 |
| | Lepomis punctatus | 479 |
| | Micropterus punctulatus | 211,479 |
| | Micropterus salmoides | 20,23,211, 252,280,309, 310,479 |
| | Pomoxis annularis | 20,211,280, 309,310 |
| 553 <i>Camallanus trispinosus</i> | Amyda ferox | 216 |
| | Chelydra serpentina | 216 |
| | Deirochelys reticularia | 216 |
| | Kinosternon subrubrum hippocrepis | 216 |
| | Lepomis cyanellus | 308 |
| | Lepomis humilis | 308 |
| | Lepomis macrochirus | 308 |
| | Lepomis megalotis | 308 |
| | Pseudemys scripta elegans | 216 |
| 554 <i>Capillaria</i> | Ambloplites rupestris | 479 |
| | Cattle | 138A |
| | Chicken | 378 |
| | Columba livia | 351 |
| | Lepomis auritus | 479 |
| | Lepomis cyanellus | 479 |
| | Lepomis gulosus | 479 |
| | Lepomis macrochirus | 479 |
| | Lepomis microlophus | 479 |
| | Lepomis punctatus | 479 |
| | Micropterus salmoides | 479 |
| | Rallus longirostris | 37 |
| 555 <i>Capillaria aerophila</i> | Canis latrans | 335N |
| | Felis rufus | 365,440 |
| 556 <i>Capillaria anatis</i> | Anas cyanoptera septentrionalium | 499 |
| 557 <i>Capillaria contorta</i> | Capella gallinago | 287 |
| 558 <i>Capillaria freitasi</i> | Passarella ilica ilica | 392 |
| 559 <i>Capillaria hepatica</i> | Sigmodon hispidus | 391 |
| | Wild Canids | 122 |
| 560 <i>Capillaria hathawayi</i> | Squalus acanthius | 390 |
| 561 <i>Capillaria heterodontis</i> | Heterodon platyrhinos | 216 |
| 562 <i>Capillaria ovopunctata</i> | Quiscalus quiscula | 32 |
| 563 <i>Capillaria palmata</i> | Eptisicus fuscus | 254 |
| 564 <i>Capillaria quisicali</i> | Quiscalus quiscula aeneus | 392 |
| 565 <i>Capillaria serpentina</i> | Chelydra serpentina | 216 |
| 566 <i>Chandlerella quisicali</i> | Quiscalus quiscula | 32 |
| 567 <i>Cheilospirura hamulosa</i> | Chicken | 378 |
| 568 <i>Choerostrogylus pudendotectus</i> | Pig | 378 |
| 569 <i>Codiostomum struthionis</i> | Ostrich | 378 |
| 570 <i>Conspicuum icteriodorum</i> | Quiscalus quiscula | 32 |
| 571 <i>Contracaecum</i> | Ambloplites rupestris | 479 |
| | Ammocrypta vivax | 479 |
| | Aplodinotus grunniens | 211 |
| | Astyanax mexicanus | 479 |
| | Cyprinodon variegatus | 24 |
| | Dorosoma petenese | 241 |

| | | |
|---------------------------------------|----------------------------|---------------------------|
| | Esox americanus | 40 |
| | Gambusia affinis | 30,127,128 |
| | Gambusia geiseri | 127 |
| | Huro salmoides | 434 |
| | Ictalurus nebulosus | 479 |
| | Ictalurus punctatus | 211 |
| | Lepisosteus osseus | 76 |
| | Lepisosteus productus | 76 |
| | Lepomis auritus | 479 |
| | Lepomis cyanellus | 479 |
| | Lepomis gulosus | 479 |
| | Lepomis macrochirus | 211,479 |
| | Lepomis megalotis | 211 |
| | Lepomis microlophus | 211,479 |
| | Micropterus punctulatus | 40,479 |
| | Micropterus salmoides | 23,40,211, 252,309,479 |
| | Notropis lutrensis | 40 |
| | Notropis umbratilis | 40 |
| | Pomoxis annularis | 211 |
| | Spinitectus carolini | 40 |
| 572 <i>Contracaecum bancrofti</i> | Pelecanus erythrorhynchoss | 305 |
| 573 <i>Contracaecum collieri</i> | Adinia xenica | 24 |
| | Cyprinodon variegatus | 24 |
| | Fundulus grandis | 24 |
| 574 <i>Contracaecum microcephalum</i> | Pelecanus erythrorhynchoss | 305 |
| 575 <i>Contracaecum spiculigerum</i> | Lepisosteus | 109 |
| | Lepisosteus osseus | 76 |
| | Lepisosteus productus | 76 |
| | Lepisosteus spatula | 76 |
| | Lepomis cyanellus | 20,308,309, 310 |
| | Lepomis humilis | 308 |
| | Lepomis macrochirus | 20,308, 309,310 |
| | Lepomis megalotis | 20,308,310 |
| | Micropterus salmoides | 20,309 |
| | Pomoxis annularis | 20,309 |
| 576 <i>Cooperia</i> | Cattle | 44,117 |
| 577 <i>Cooperia bisonia</i> | Sheep | 10N |
| 578 <i>Cooperia curticei</i> | Goat | 378 |
| | Sheep | 378 |
| 579 <i>Cooperia mcmasteri</i> | Bovine | 13A |
| | Cattle | 39A |
| 580 <i>Cooperia oncophora</i> | Ammotragus lervia | 13N |
| | Antilocapra americana | 207N |
| | Cattle | 39A |
| | Goat | 378 |
| | Odocoileus heminous | 19A |
| | Sheep | 378 |
| 581 <i>Cooperia pectinata</i> | Antilocapra americana | 207N |
| | Cattle | 39N |
| 582 <i>Cooperia punctata</i> | Antilocapra americana | 207N |
| | Cattle | 39N |
| | Goat | 378 |
| | Sheep | 378 |

| | | | |
|-----|------------------------------------|------------------------------|--------------------|
| 583 | <i>Cosmocephalus capellae</i> | Capella gallinago | 287 |
| 584 | <i>Cosmocercoides dukae</i> | Ambystoma talpoideum | 216 |
| | | Ambystoma texanum | 216 |
| | | Bufo terrestris | 216 |
| | | Bufo valliceps | 216 |
| | | Eumeces fasciatus | 216 |
| | | Gastrophyrne olivacea | 216 |
| | | Heterodon platyrhinos | 216 |
| | | Hyla squirella | 216 |
| | | Micrurus fulvius | 216 |
| | | Notophthalmus meridionalis | 216 |
| | | Ophisaurus ventralis | 216 |
| | | Pseudacris triseriata | 216 |
| | | Rana areolata | 216 |
| | | Rana catesbeiana | 216, 233, 425 |
| | | Rana clamitans | 216 |
| | | Rana palustris | 216 |
| | | Rana pipiens complex | 425 |
| | | Rana sphenoccephala | 216 |
| | | Scincella lateralis | 216 |
| | | Storeria dekayi | 216 |
| | | Terrapene carolina triunguis | 216 |
| | | Terrapene ornata | 216 |
| 585 | <i>Cruzia americana</i> | Dasyus novemcinctus | 63 |
| 586 | <i>Cruzia tentaculata</i> | Didelphis virginiana | 83 |
| 587 | <i>Cruzia tesfundinis</i> | Terrapene carolina triunguis | 216 |
| 588 | <i>Cryptosomum heyenemani</i> | Cophosaurus texanus | 486A |
| | | Holbrookia maculata | 486A |
| 589 | <i>Cryptosomum penneri</i> | Sceloporus magister | 486A |
| 590 | <i>Cyathospirura chavreuxi</i> | Felis rufus | 365, 440 |
| 591 | <i>Cylicospirura felineus</i> | Felis rufus | 365, 440 |
| 592 | <i>Cylicospirura subaequalis</i> | Felis rufus | 291 |
| 593 | <i>Cyrnea</i> | Canis latrans | 370 |
| | | Colinus virginianus texanus | 491 |
| 594 | <i>Dacnitoides cotylophora</i> | Ictalurus punctatus | 211 |
| 595 | <i>Dermatoxys veligera</i> | Canis latrans | 364, 365 |
| | | Lepus californicus | 404N |
| | | Sylvilagus auduboni | 132N |
| 596 | <i>Dichelyne lepisosteus</i> | Lepisosteus productus | 76 |
| | | Lepisosteus spatula | 76 |
| 597 | <i>Dichelyne robusta</i> | Ictalurus punctatus | 280 |
| 598 | <i>Dictyocaulus filaria</i> | Goat | 378 |
| | | Sheep | 225N, 378 |
| 599 | <i>Dictyocaulus viviparus</i> | Cervus elaphus | 501N |
| | | Cow | 424 |
| 600 | <i>Didelphonema longispiculata</i> | Felis rufus | 365, 440 |
| 601 | <i>Dipetalonema reconditum</i> | Dog | 266, 374 |
| 602 | <i>Diplotriaena</i> | Quiscalus quiscula | 32 |
| 603 | <i>Diplotriaena tricuspis</i> | Corvus cryptoleucus | 371N |
| 604 | <i>Dirofilaria acutiuscula</i> | Tayassu tajacu | 409 |
| 605 | <i>Dirofilaria immitis</i> | Canis latrans | 364, 365, 374, 464 |
| | | Dog | 266, 348, 378, 430 |
| | | Wild Canids | 121, 122 |

| | | | |
|-----|-----------------------------------|----------------------------------|--------------------------|
| 606 | <i>Dirofilaria scapiceps</i> | Sylvilagus floridanus | 475 |
| 607 | <i>Dirofilaria striata</i> | Dog | 348 |
| 608 | <i>Dirofilaria tenuis</i> | Procyon lotor | 93 |
| 609 | <i>Dispharynx nasuta</i> | Ortalis vetula maccalli | 105 |
| 610 | <i>Dracunculus insignis</i> | Dog | 477 |
| | | Procyon lotor | 90 |
| 611 | <i>Dracunculus ophdensis</i> | Nerodia erythrogaster transversa | 119 |
| 612 | <i>Echinuria uncinata</i> | Anas creca | 74 |
| | | Dendrocygna autumnalis | 205 |
| 613 | <i>Elaeophora schneideri</i> | Ammotragus lervia | 367 |
| | | Cervus elaphus | 131A, 501N |
| | | Elk | 131N |
| | | Horse Fly | 131N |
| | | Hybomitra | 221N |
| | | Hybomitra laticornis | 124N |
| | | Odocoileus hemionus | 367, 457A |
| | | Odocoileus virginianus | 196 |
| | | Sheep | 265N |
| 614 | <i>Epomidiostomum uncinatum</i> | Duck | 68 |
| | | Gadwall | 69 |
| | | Pintail | 69 |
| | | Shoveler | 69 |
| 615 | <i>Eucoleus aerophilus</i> | Urocyon cinereoargenteus | 65 |
| 616 | <i>Eustrongylides</i> | Ambloplites rupestris | 479 |
| | | Fundulus grandis | 24 |
| | | Lepisosteus osseus | 76 |
| | | Lepisosteus spatula | 76 |
| | | Lepomis gulosus | 479 |
| | | Micropterus salmoides | 479 |
| | | Nerodia erythrogaster transversa | 119 |
| 617 | <i>Falcaustra affine</i> | Terrapene carolina triunguis | 216 |
| 618 | <i>Falcaustra catesbeiana</i> | Rana catesbeiana | 216, 425 |
| | | Rana pipiens complex | 425 |
| 619 | <i>Falcaustra chelydrae</i> | Amyda ferox | 216 |
| | | Chelydra serpentina | 216 |
| 620 | <i>Falcaustra procera</i> | Pseudemys scripta elegans | 216 |
| 621 | <i>Fibricola cratera</i> | Mustela vison | 386 |
| 622 | <i>Filaria martis</i> | Conepatus mesoleucus | 466 |
| | | Mephitis mephitis | 466 |
| 623 | <i>Filaria texensis</i> | Mephitis mephitis | 98 |
| 624 | <i>Filaroides osleri</i> | Canis latrans | 294, 335A, 364, 365, 464 |
| 625 | <i>Foleyella americana</i> | Rana catesbeiana | 233 |
| 626 | <i>Globocephalus urosubulatus</i> | Sus scrofa | 113 |
| | | Pig | 505 |
| 627 | <i>Gnathostoma didelphis</i> | Didelphis virginiana | 83 |
| 628 | <i>Gnathostoma procyonis</i> | Procyon lotor | 93 |
| 629 | <i>Gongylonema</i> | Sigmodon hispidus | 298 |
| 630 | <i>Gongylonema baylisi</i> | Tayassu tajacu | 409 |
| 631 | <i>Gongylonema pulchrum</i> | Odocoileus virginianus | 183 |
| | | Onthophagus gazella | 189 |
| | | Sus scrofa | 113 |
| | | Ursus americanus | 100 |
| 632 | <i>Gongylonema ransomi</i> | Pig | 378 |
| 633 | <i>Gongylonema scutatum</i> | Goat | 378 |
| | | Sheep | 378 |

| | | | |
|-----|----------------------------------|----------------------------------|---------|
| 634 | <i>Gongylonema verrucosum</i> | Goat | 378 |
| | | Sheep | 378 |
| 635 | <i>Gyalocephalus equi</i> | Horse | 378 |
| | | Mule | 378 |
| 636 | <i>Habronema megastoma</i> | Horse | 378 |
| | | Mule | 378 |
| 637 | <i>Habronema microstoma</i> | Horse | 378 |
| | | Mule | 378 |
| 638 | <i>Habronema muscae</i> | Horse | 378 |
| | | Mule | 378 |
| 639 | <i>Haemonchus</i> | Cattle | 44N |
| | | Ovis aries | 403 |
| | | Ovis canadensis | 15N |
| | | Ovis trageiaphus | 15N |
| 640 | <i>Haemonchus contortus</i> | Antilope cervicapra | 463 |
| | | Cattle | 378 |
| | | Cow | 424 |
| | | Goat | 378 |
| | | Odocoileus virginianus | 183 |
| | | Sheep | 8N,378 |
| 641 | <i>Haemonchus placei</i> | Ammotragus lervia | 13N |
| | | Antilocapra americana | 207N |
| | | Cattle | 39N |
| | | Ovis canadensis | 11N |
| 642 | <i>Haemonchus similis</i> | Urocyon cinereoargenteus | 65 |
| 643 | <i>Hedrurus</i> | Elassoma zonatum | 40 |
| 644 | <i>Hassalstrongylus aduncus</i> | Sigmodon hispidus | 298 |
| 645 | <i>Heligmodendrium hassalli</i> | Sciurus carolinensis | 91 |
| | | Sciurus niger | 91 |
| 646 | <i>Heterakis</i> | Phasianus colchicus | 142 |
| 647 | <i>Heterakis gallinarum</i> | Meleagris gallopavo intermedia | 361 |
| | | Phasianus colchicus | 142,366 |
| 648 | <i>Heterakis isolonche</i> | Phasianus colchicus | 142 |
| | | Tympanuchus pallidicinctus | 363,369 |
| 649 | <i>Heterakis vesiculavis</i> | Chicken | 378 |
| 650 | <i>Hyostrongylus rubidus</i> | Pig | 378 |
| 651 | <i>Kalicephalis</i> | Agkistrodon contortrix | 216 |
| | <i>agkistrodontis</i> | Heterodon platyrhinos | 216 |
| | | Lampropeltis getulus holbrooki | 216 |
| | | Micrurus fulvius | 216 |
| | | Nerodia rhombifera | 216 |
| | | Nerodia fasciata | 216 |
| | | Pituophis melanoleucus sayi | 216 |
| | | Thamnophis proximus | 216 |
| 652 | <i>Kalicephalus</i> | Coluber constrictor flaviventris | 216 |
| | <i>agkistrodontis flagellus</i> | Masticophis flagellum | 216 |
| 653 | <i>Kalicephalus rectiphilus</i> | Coluber constrictor flaviventris | 216 |
| 654 | <i>Libyostrongylus douglassi</i> | Ostrich | 378 |
| 655 | <i>Litomosoides</i> | Neotoma micropus | 172 |
| 656 | <i>Litomosoides carinii</i> | Ondatra zibethicus | 85 |
| | | Sigmodon hispidus | 172,417 |
| 657 | <i>Litomosoides microfilaria</i> | Neotoma micropus | 172 |
| 658 | <i>Litomosoides sigmodontis</i> | Sigmodon hispidus | 82 |
| 659 | <i>Longistriata adunca</i> | Sigmodon | 416 |

| | | |
|---|---------------------------|-------------|
| | Sigmodon hispidus | 84,250 |
| 660 <i>Lungworm</i> | Bullfrog | 339 |
| | Leopard Frog | 339 |
| 661 <i>Macroderoides spiniferus</i> | Lepisosteus productus | 76 |
| 662 <i>Mastophorus muris</i> | Sigmodon hispidus | 328 |
| 663 <i>Mastophorus muris</i> <i>ascaroides</i> | Sigmodon hispidus | 250 |
| 664 <i>Mastophorus muris</i> <i>var. muris</i> | Rattus rattus | 250 |
| | Sigmodon hispidus | 250 |
| 665 <i>Metabronema salvelini</i> | Salmo apache | 340A |
| | Salmo gilae | 340A |
| | Salmo trutta | 340A |
| 666 <i>Metastrongylus</i> | Sus scrofa | 113 |
| 667 <i>Metastrongylus elongatus</i> | Pig | 378 |
| 668 <i>Metastrongylus salmi</i> | Pig | 378 |
| 669 <i>Metathelazia californica</i> | Felis rufus | 360,365,440 |
| 670 <i>Metathelazia felineus</i> | Felis rufus | 365 |
| 671 <i>Micipsella brevicauda</i> | Lepus californicus | 404N |
| 672 <i>Microfilaria</i> | Chonestes grammacus | 114 |
| | Cyanocitta cristata | 114 |
| | Hesperiphona vespertina | 506A |
| | Icterus spurius | 114 |
| | Peromyscus leucopus | 172 |
| | Spermophilus mexicanus | 172 |
| | Zenaida macroura | 114 |
| 673 <i>Microfilaria alpha</i> <i>and beta</i> | Sciurus niger | 91 |
| 674 <i>Microtetrameres helix</i> | Corvus brachyrhynchos | 341 |
| 675 <i>Molineus</i> | Conepatus mesoleucus | 466 |
| | Mephitis mephitis | 466 |
| | Spilogale putorius | 466 |
| 676 <i>Molineus barbatus</i> | Procyon lotor | 93 |
| 677 <i>Molinostrongylus delicatus</i> | Tadarida brasiliensis | 71N |
| 678 <i>Nematode</i> | Mormoops megalophylla | 254 |
| 679 <i>Nematode larva</i> | Cnemidophorus tessellatus | 206N |
| 680 <i>Nematodirella</i> <i>longispiculata</i> | Antilocapra americana | 207N |
| 681 <i>Nematodirus</i> | Cattle | 44 |
| | Lepus californicus | 404N |
| 682 <i>Nematodirus helvetianus</i> | Ammotragus lervia | 13N |
| | Cattle | 39N |
| 683 <i>Nematodirus lanceolatus</i> | Antilocapra americana | 207N |
| | Sheep | 16N |
| 684 <i>Nematodirus leporis</i> | Rabbit | 378 |
| 685 <i>Nematodirus neomexicanus</i> | Lepus californicus | 404N |
| 686 <i>Nematodirus spathiger</i> | Ammotragus lervia | 13N |
| | Antilope cervicapra | 463 |
| | Goat | 378 |
| | Ovis canadensis | 6N |
| | Sheep | 378 |
| 687 <i>Neodiplostomum lucidum</i> | Mustela vison | 386 |
| 688 <i>Obeliscoides cuniculi</i> | Rabbit | 378 |
| 689 <i>Oesophagostomum</i> | Antilope cervicapra | 463 |
| | Ovis canadensis | 11N |

| | | | |
|-----|--------------------------------------|----------------------------------|----------|
| 690 | <i>Oesophagostomum columbianum</i> | Odocoileus virginianus | 183 |
| 691 | <i>Oesophagostomum dentatum</i> | Pig | 378 |
| 692 | <i>Oesophagostomum radiatum</i> | Cattle | 39N,138A |
| 693 | <i>Onchocerca cervicalis</i> | Horse | 378 |
| | | Mule | 378 |
| 694 | <i>Ophidascaris</i> | Coluber constrictor flaviventris | 216 |
| 695 | <i>Oslerus osleri</i> | Canis latrans | 370,381 |
| | | Dog | 378 |
| | | Wild Canids | 122 |
| 696 | <i>Ostertagia</i> | Cattle | 44 |
| | | Odocoileus virginianus | 208 |
| 697 | <i>Ostertagia circumcincta</i> | Goat | 378 |
| | | Sheep | 378 |
| 698 | <i>Ostertagia orloffii</i> | Ammotragus lervia | 13N |
| 699 | <i>Ostertagia ostertagi</i> | Cattle | 39N,117 |
| | | Cervus elaphus | 501N |
| 700 | <i>Oswaldocruzia pipiens</i> | Bufo terrestris | 216 |
| | | Bufo valliceps | 216 |
| | | Eumeces fasciatus | 216 |
| | | Hyla cinerea | 216 |
| | | Rana palustris | 216 |
| | | Rana sphenoccephala | 216 |
| | | Sceloporus undulatus | 216 |
| | | Scincella lateralis | 216 |
| | | Terrapene carolina triunguis | 216 |
| | | Terrapene ornata | 216 |
| 701 | <i>Oxysomatium americanum</i> | Rana catesbeiana | 233 |
| 702 | <i>Oxysomatium variabilis</i> | Bufo valliceps | 215 |
| 703 | <i>Oxyspirura petrowi</i> | Callipepla squamata | 123,356 |
| | | Cyrtonyx montezumae | 356 |
| | | Phasianus colchicus | 366 |
| | | Tympanuchus pallidicinctus | 363,369 |
| 704 | <i>Oxyuris equi</i> | Horse | 378 |
| | | Mule | 378 |
| 705 | <i>Oxyuroid larva</i> | Ambystoma tigrinum | 505N |
| | | Carpiodes carpio | 420 |
| 706 | <i>Parabronema</i> | White-collared Peccary | 415 |
| 707 | <i>Parabronema pecariae</i> | Tayassu tajacu | 405N,409 |
| 708 | <i>Paramacroderoides echinus</i> | Lepisosteus osseus | 76 |
| 709 | <i>Parascaris equorum</i> | Horse | 378 |
| | | Mule | 378 |
| 710 | <i>Parathelandros texanus</i> | Cnemidophorus inornatus | 436 |
| | | Cnemidophorus scalaris | 436 |
| | | Cnemidophorus tigris | 436 |
| | | Cophosaurus texanus | 436 |
| | | Sceloporus merriami | 436 |
| | | Sceloporus undulatus | 436 |
| | | Urosaurus ornatus | 436 |
| 711 | <i>Parelaphostrongylus tenuis</i> | Goat | 212 |
| 712 | <i>Parhadjelia neglecta</i> | Dendrocygna autumnalis | 205 |
| 713 | <i>Parostertagia heterospiculum</i> | Tayassu tajacu | 409 |
| | | White-collared Peccary | 415 |
| 714 | <i>Parralintoshius parallintoshi</i> | Myotis velifer | 254 |

| | | | |
|-----|---|---|--------------------|
| 715 | <i>Passalurus nonanulatus</i> | <i>Lepus californicus</i> | 404 |
| 716 | <i>Pharyngodon kirbii</i> | <i>Cnemidophorus scalaris</i> | 435 |
| 717 | <i>Pharyngodon mudgi</i> | <i>Coleonyx brevis</i> | 435 |
| 718 | <i>Pharyngodon warneri</i> | <i>Cnemidophorus sexlineatus</i> | 216 |
| | | <i>Cnemidophorus tigris</i> | 31A |
| | | <i>Urosaurus ornatus</i> | 486A |
| 719 | <i>Pharyngodon warneri slim and stout</i> | <i>Cnemidophorus tessellatus</i> | 206N |
| 720 | <i>Philometra</i> | <i>Fundulus grandis</i> | 24 |
| | | <i>Lepomis macrochirus</i> | 40 |
| 721 | <i>Philometra nodulosa</i> | <i>Micropterus salmoides</i> | 40,211,252, 280 |
| 722 | <i>Physaloptera</i> | <i>Baiomys taylori</i> | 316 |
| | | <i>Bassariscus astutus</i> | 362 |
| | | <i>Canis latrans</i> | 464 |
| | | <i>Felis rufus</i> | 291 |
| | | <i>Sigmodon hispidus</i> | 328 |
| | | <i>Tympanuchus pallidicinctus</i> | 369 |
| 723 | <i>Physaloptera aquamatae</i> | <i>Agkistrodon contortrix</i> | 216 |
| 724 | <i>Physaloptera bispiculata</i> | <i>Sigmodon hispidus</i> | 298 |
| 725 | <i>Physaloptera larva</i> | Armadillo | 96 |
| 726 | <i>Physaloptera maxillaris</i> | <i>Conepatus mesoleucus</i> | 466 |
| | | <i>Mephitis mephitis</i> | 466 |
| | | <i>Spilogale putorius</i> | 466 |
| 727 | <i>Physaloptera phrynosoma</i> | <i>Phrynosoma cornutum</i> | 216,482 |
| 728 | <i>Physaloptera praeputialis</i> | <i>Felis rufus</i> | 365,440 |
| 729 | <i>Physaloptera rara</i> | <i>Canis latrans</i> | 364,365,370 |
| | | <i>Urocyon cinereoargenteus</i> | 65 |
| | | Wild Canids | 122 |
| 730 | <i>Physaloptera retusa</i> | <i>Sceloporus magister</i> | 486A |
| 731 | <i>Physaloptera squamata</i> | <i>Scincella lateralis</i> | 216 |
| 732 | <i>Physaloptera turgida</i> | <i>Didelphis virginiana</i> | 83 |
| 733 | <i>Physocephalus</i> | Armadillo | 96 |
| | | <i>Canis latrans</i> | 370 |
| | | <i>Tayassu tajacu</i> | 409 |
| 734 | <i>Physocephalus sexualatus</i> | Chicken | 3N |
| | | <i>Onthophagus gazella</i> | 189 |
| | | Pig | 378 |
| | | <i>Sus scrofa</i> | 113 |
| | | White-collared Peccary | 415 |
| 735 | <i>Pneumospirura bassarisci</i> | <i>Bassariscus astutus</i> | 360,362 |
| 736 | <i>Polydelphis</i> | <i>Coluber constrictor flaviventris</i> | 216 |
| 737 | <i>Porrocaecum</i> | <i>Dendrocygna autumnalis</i> | 205 |
| 738 | <i>Probstmayria vivipara</i> | Horse | 378 |
| | | Mule | 378 |
| 739 | <i>Procamallanus</i> | <i>Fundulus grandis</i> | 24 |
| 740 | <i>Proleptus</i> | <i>Cnemidophorus tessellatus</i> | 206N |
| 741 | <i>Proteracrum columbianum</i> | Goat | 378 |
| | | Sheep | 378 |
| 742 | <i>Proteracrum radiatum</i> | Cattle | 378 |
| 743 | <i>Protospirura</i> | <i>Perognathus penicillatus</i> | 25A |
| 744 | <i>Protospirura numidica</i> | <i>Canis latrans</i> | 370 |
| 745 | <i>Pseudostertagia bullosa</i> | <i>Ammotragus lervia</i> | 13N |
| | | <i>Antilocapra americana</i> | 207 |
| | | <i>Ovis canadensis</i> | 11N |
| | | Sheep | 346N |

| | | | |
|-----|---|---|---------------------------------------|
| 746 | <i>Pterygodermatites cahirensis</i> | <i>Canis latrans</i> | 364,365 |
| | | <i>Felis rufus</i> | 365,440 |
| 747 | <i>Quadriplotriaena hypsoykysta</i> | <i>Sturnella magna magna</i> | 284 |
| 748 | <i>Raphidascaris</i> | <i>Fundulus grandis</i> | 24 |
| 749 | <i>Rhabdias</i> | <i>Nerodia erythrogaster transversa</i> | 119 |
| 750 | <i>Rhabdias ranae</i> | <i>Acris crepitans</i> | 49 |
| | | <i>Bufo valliceps</i> | 231 |
| | | <i>Bufo woodhousei fowleri</i> | 337 |
| | | <i>Rana catesbeiana</i> | 216,233,337 |
| | | <i>Rana clamitans</i> | 337 |
| | | <i>Rana pipiens complex</i> | 337,425 |
| | | <i>Rana sphenoccephala</i> | 216 |
| 751 | <i>Rhabdias vellardi</i> | <i>Heterodon platyrhinos</i> | 216 |
| | | <i>Storeria dekayi</i> | 216 |
| | | <i>Thamnophis proximus</i> | 216 |
| | | <i>Virginia striatulus</i> | 216 |
| 752 | <i>Rhabditid nematode</i> | Japanese Beetle | 248 |
| 753 | <i>Rhabdochona</i> | <i>Cichlosoma cyanoguttatum</i> | 479 |
| | | <i>Gambusia affinis</i> | 127,128 |
| | | <i>Salmo gairdneri</i> | 497N |
| | | <i>Salmo trutta</i> | 497N |
| | | <i>Tilapia mossambica</i> | 479 |
| 754 | <i>Rhabdochona decaturensis</i> | <i>Aplodinotus grunniens</i> | 211 |
| | | <i>Ictalurus punctatus</i> | 211 |
| 755 | <i>Rictularia dipodomis</i> | <i>Dipodomys</i> | 467A |
| 756 | <i>Rictularia ondatrae</i> | <i>Ondatra zibethicus</i> | 85 |
| | | <i>Sigmodon hispidus</i> | 318 |
| 757 | Roundworm | <i>Dipodomys merriami</i> | 329 |
| 758 | <i>Sciadiocara</i> | <i>Grus canadensis</i> | 67,138A |
| 759 | <i>Setaria</i> | <i>Odocoileus virginianus</i> | 181 |
| 760 | <i>Setaria cervi</i> | Cattle | 138A |
| 761 | <i>Setaria equina</i> | Horse | 378 |
| | | Mule | 378 |
| 762 | <i>Skrjabinema</i> | <i>Ammotragus lervia</i> | 13N |
| | | <i>Ovis canadensis</i> | 6N,19A |
| 763 | <i>Skrjabinema caprae</i> | <i>Capra hircus</i> | 412N |
| 764 | <i>Skrjabinema ovis</i> | <i>Ovis canadensis</i> | 11N |
| 765 | <i>Skrjabinigylus chitwoodorum</i> | <i>Conepatus mesoleucus</i> | 466 |
| | | <i>Mephitis mephitis</i> | 466 |
| | | <i>Spilogale putorius</i> | 466 |
| 766 | <i>Skrjabinoptera phrynosoma</i> | <i>Phrynosoma cornutum</i> | 283 |
| 767 | <i>Spinitectus carolini</i> | <i>Ambloplites rupestris</i> | 479 |
| | | <i>Astyanax mexicanus</i> | 479 |
| | | <i>Gambusia affinis</i> | 128 |
| | | <i>Gambusia geiseri</i> | 127 |
| | | <i>Ictalurus natalis</i> | 479 |
| | | <i>Lepomis auritus</i> | 479 |
| | | <i>Lepomis cyanellus</i> | 20,308,309, 310,479 |
| | | <i>Lepomis gulosus</i> | 211,479 |
| | | <i>Lepomis humilis</i> | 306 |
| | | <i>Lepomis macrochirus</i> | 20,40,211, 280,308,309, 310,479 |

| | | |
|-------------------------------------|----------------------------------|-------------------------|
| | Lepomis megalotis | 20,211,308, 309 |
| | Lepomis microlophus | 479 |
| | Lepomis punctatus | 479 |
| | Micropterus punctulatus | 211,479 |
| | Micropterus salmoides | 23,211,252, 479 |
| | Pomoxis annularis | 20,211,309 |
| 768 <i>Spinitectus gracilis</i> | Aplodinotus grunniens | 211 |
| | Ictalurus punctatus | 211 |
| | Lepomis auritus | 479 |
| | Lepomis cyanellus | 479 |
| | Lepomis gulosus | 479 |
| | Lepomis macrochirus | 479 |
| | Lepomis microlophus | 479 |
| | Lepomis punctatus | 479 |
| 769 <i>Spinitectus micracanthus</i> | Ambloplites rupestris | 479 |
| | Astyanax mexicanus | 479 |
| | Ictalurus natalis | 479 |
| | Lepomis auritus | 479 |
| | Lepomis cyanellus | 479 |
| | Lepomis gulosus | 479 |
| | Lepomis macrochirus | 479 |
| | Lepomis microlophus | 479 |
| | Lepomis punctatus | 479 |
| | Micropterus punctulatus | 479 |
| | Micropterus salmoides | 479 |
| 770 <i>Spirocamallanus pereirai</i> | Micropogon undulatus | 259 |
| 771 <i>Spirocerca lupi</i> | Canis latrans | 335,364,365, 370,464 |
| | Dog | 430 |
| | Felis rufus | 365,440 |
| | Urocyon cinereoargenteus | 65 |
| | Wild Canids | 122 |
| 772 <i>Spirocerca sanguinolenta</i> | Dog | 378 |
| 773 <i>Spirometra</i> | Domestic Cat | 389 |
| 774 <i>Spiroonoura gracilis</i> | Ambystoma tigrinum | 505N |
| 775 <i>Spiroxys</i> | Gambusia affinis | 127,128 |
| | Gambusia geiseri | 127 |
| | Lepomis megalotis | 211 |
| | Lepomis microlophus | 211 |
| | Nerodia erythrogaster transversa | 120 |
| 776 <i>Spiroxys amydae</i> | Amyda ferox | 216 |
| 777 <i>Spiroxys contorta</i> | Ambloplites rupestris | 479 |
| | Astyanax mexicanus | 479 |
| | Cicholosoma cyanoguttatum | 479 |
| | Dendrochelys reticularia | 216 |
| | Ictalurus natalis | 479 |
| | Lepomis auritus | 479 |
| | Lepomis cyanellus | 479 |
| | Lepomis gulosus | 479 |
| | Lepomis macrochirus | 479 |
| | Lepomis microlophus | 479 |
| | Lepomis punctatus | 479 |
| | Micropterus punctulatus | 479 |
| | Micropterus salmoides | 479 |

| | | |
|---------------------------------------|----------------------------------|-------------|
| | Moxostoma congestum | 479 |
| | Tilapia mossambica | 479 |
| | Pseudemys scripta elegans | 216 |
| | Sternotherus carinatus | 216 |
| 778 <i>Spiruroidea</i> | Perognathus hispidus | 316 |
| 779 <i>Spiruroid larva</i> | Acris crepitans | 49 |
| 780 <i>Splendofilaria</i> | Corvus brachyrhynchos | 341 |
| | Quiscalus quiscula | 32 |
| 781 <i>Stephanofilaria stilesi</i> | Haematobia irritans | 227N |
| 782 <i>Stephanurus dentatus</i> | Pig | 378 |
| | Sus scrofa | 113 |
| 783 <i>Streptocara crassicanda</i> | Anas cyanoptera septentrioanlium | 499 |
| 784 <i>Strongylidae sigmodontis</i> | Sigmodon | 416 |
| 785 <i>Strongyloid larva</i> | Acris crepitans | 49 |
| 786 <i>Strongyloides</i> | Acris crepitans | 49 |
| | Dog | 94 |
| | Odocoileus virginianus | 208 |
| 787 <i>Strongyloides robustus</i> | Sciurus carolinensis | 91 |
| | Sciurus niger | 91 |
| 788 <i>Strongyloides sigmodontis</i> | Sigmodon hispidus | 318 |
| 789 <i>Strongyloides tumefaciens</i> | Felis domesticus | 199 |
| 790 <i>Strongylus edentatus</i> | Horse | 378 |
| | Mule | 378 |
| 791 <i>Strongylus equinus</i> | Horse | 378 |
| | Mule | 378 |
| 792 <i>Strongylus vulgaris</i> | Horse | 378 |
| | Mule | 378 |
| 793 <i>Sublurua</i> | Canis latrans | 364,365 |
| | Cynomys ludovicianus | 372N |
| 794 <i>Syngamus traceha</i> | Colinus virginianus texanus | 480 |
| 795 <i>Synhimantus</i> | Canis latrans | 370 |
| 796 <i>Synhimantus longigutturata</i> | Procyon lotor | 93 |
| 797 <i>Syphacia</i> | Canis latrans | 364,365 |
| 798 <i>Syphacia obvelata</i> | Rattus rattus | 250 |
| 799 <i>Syphacia sigmodontis</i> | Sigmodon hispidus | 298,328 |
| 800 <i>Tetrameres americana</i> | Columba livia | 351 |
| 801 <i>Tetrameres coloradensis</i> | Capella gallinago | 287 |
| 802 <i>Tetrameres crami</i> | Anas clypeata | 58 |
| | Anas creca | 74 |
| 803 <i>Tetrameres grusi</i> | Grus canadensis | 198 |
| 804 <i>Texicospirura turki</i> | Tayassu tajacu | 405N,409 |
| 805 <i>Thelandros salamandrae</i> | Plethedon neomexicanus | 352N |
| 806 <i>Thelazia californiensis</i> | Odocoileus hemionus | 412N |
| 807 <i>Thubunaea leiopismae</i> | Acris gryllus | 216 |
| | Scincella lateralis | 216 |
| 808 <i>Toxascaris leonina</i> | Canis familiaris | 217 |
| | Canis latrans | 364,365,370 |
| | Dog | 430 |
| | Felis rufus | 291,365,440 |
| | Wild Canids | 122 |
| 809 <i>Toxascaris limbata</i> | Dog | 378 |
| 810 <i>Toxocara canis</i> | Canis familiaris | 217N |
| | Dog | 223,378,430 |
| | | 438 |
| 811 <i>Toxocara mystax</i> | Cat | 378 |

| | | |
|---|-------------------------|-------------|
| | Felis rufus | 363,440 |
| 812 <i>Trichinella</i> | Ursus americanus | 281A |
| 813 <i>Trichinella spiralis</i> | Man | 509 |
| | Pig | 509 |
| 814 <i>Trichuridea</i> | Gambusia affinis | 127 |
| 815 <i>Trichiuroid</i> | Garfish | 109 |
| 816 <i>Trichospirura leptostoma</i> | Callithrix jacchus | 432 |
| 817 <i>Trichostrongylus</i> | Cattle | 44,138 |
| 818 <i>Trichostrongylus axei</i> | Ammotragus lervia | 13N |
| | Antilocapra americana | 207N |
| | Antilope cervicapra | 463 |
| | Cattle | 39N,117 |
| | Cervus elaphus | 501N |
| | Goat | 378 |
| | Horse | 378 |
| | Mule | 378 |
| | Sheep | 378 |
| 819 <i>Trichostrongylus calcaratus</i> | Ondatra zibethicus | 101 |
| | Rabbit | 378 |
| 820 <i>Trichostrongylus colubriformis</i> | Antilocapra americana | 207N |
| | Antilope cervicapra | 463 |
| | Cervus elaphus | 501 |
| | Goat | 378 |
| | Lepus californicus | 404N,405N |
| | Sheep | 378 |
| | Tayassu tajacu | 405N |
| 821 <i>Trichostrongylus longispicularis</i> | Ammotragus lervia | 13N |
| 822 <i>Trichostrongylus proboiurus</i> | Ammotragus lervia | 13N |
| | Antilope cervicapra | 463 |
| 823 <i>Trichostrongylus ransomi</i> | Sylvilagus floridanus | 139 |
| 824 <i>Trichostrongylus sigmodontis</i> | Meriones unguiculatus | 462 |
| | Mesocricetus auratus | 462 |
| | Oryzomys palustris | 462 |
| | Sigmodon hispidus | 462 |
| 825 <i>Trichostrongylus texianus</i> | Cynomys ludovicianus | 139 |
| 826 <i>Trichuridea</i> | Gambusia affinis | 128 |
| 827 <i>Trichuris</i> | Canis latrans | 364,365,370 |
| | Ovis canadensis | 6N |
| 828 <i>Trichuris dipodomis</i> | Dipodomys ordi | 394N,395 |
| 829 <i>Trichuris discolor</i> | Ovis canadensis | 11N |
| 830 <i>Trichuris globulosa</i> | Antilope cervicapra | 463 |
| 831 <i>Trichuris ovis</i> | Cattle | 378 |
| | Cervus nippon | 268 |
| | Goat | 378 |
| | Sheep | 378 |
| 832 <i>Trichuris perognathi</i> | Peromyscus californicus | 95A |
| 833 <i>Trichuris peromysci</i> | Peromyscus californicus | 95A |
| 834 <i>Trichuris skrjabini</i> | Cervus nippon | 268 |
| 835 <i>Trichuris tenuis</i> | Camelus dromedarius | 81 |
| 836 <i>Trichuris trichiura</i> | Pig | 378 |
| 837 <i>Trichuris vulpis</i> | Canis familiaris | 217N |

| | | |
|--|---------------------------------|-------------|
| | Dog | 378,430,438 |
| | Wild Canids | 122 |
| 838 <i>Triodontophorus tenuicollis</i> | Horse | 378 |
| | Mule | 378 |
| 839 <i>Troglostrongylus wilsoni</i> | <i>Felis rufus</i> | 291 |
| 840 <i>Tropisurus</i> | <i>Grus canadensis</i> | 67 |
| 841 <i>Tropisurus crami</i> | <i>Dendrocygna autumnalis</i> | 205 |
| 842 <i>Uncinaria lotoris</i> | <i>Bassariscus astutus</i> | 381 |
| 843 <i>Vampirolepis gertschi</i> | <i>Myotis thysanodes</i> | 73N |
| 844 <i>Vogeloides felis</i> | <i>Felis rufus</i> | 360,440 |
| 845 <i>Wehrdikmansia cervipedes</i> | <i>Odocoileus hemionus</i> | 226A |
| | <i>Odocoileus virginianus</i> | 226A |
| 846 <i>Wellcomia</i> | <i>Perognathus penicillatus</i> | 25A |
| 847 <i>Wellcomia perognathi</i> | <i>Perognathus intermedius</i> | 276A |

Acanthocephala

| | | |
|--|---|---------|
| 848 <i>Acanthocephala</i> | <i>Conepatus mesoleucus</i> | 466 |
| | <i>Mephitis mephitis</i> | 466 |
| | <i>Phrynosoma cornutum</i> | 482 |
| | <i>Spilogale putorius</i> | 466 |
| 849 <i>Arhythmorhynchus capellae</i> | <i>Capella gallinago</i> | 287 |
| 850 <i>Arhythmorhynchus duocinctum</i> | <i>Adinia xenica</i> | 24 |
| | <i>Fundulus grandis</i> | 24 |
| | <i>Fundulus similis</i> | 24 |
| 851 <i>Atactorhynchus verecendus</i> | <i>Fundulus grandis</i> | 24 |
| | <i>Cyprinodon variegatus</i> | 24 |
| | <i>Mollienesia formosa</i> | 24 |
| | <i>Mollienesia latipinna</i> | 24 |
| 852 <i>Centrorhynchus</i> | <i>Corvus brachyrhynchos</i> | 341 |
| 853 <i>Centrorhynchus larva</i> | <i>Rana catesbeiana</i> | 233 |
| 854 <i>Corynosoma</i> | <i>Dendrocygna autumnalis</i> | 205 |
| 855 <i>Corynosoma constrictum</i> | <i>Anas creca</i> | 74 |
| | <i>Anas cyanoptera septentrionalium</i> | 499 |
| | Duck | 68 |
| | Gadwall | 69 |
| | Pintail | 69 |
| | Shoveler | 69 |
| 856 <i>Corynosoma peposacae</i> | <i>Dendrocygna autumnalis</i> | 205 |
| 857 <i>Eocollis arcanus</i> | <i>Chaenobryttus coronarius</i> | 314 |
| | <i>Chaenobryttus gulosus</i> | 40 |
| | <i>Lepomis cyanelus</i> | 40 |
| | <i>Lepomis macrochirus</i> | 313,314 |
| | <i>Pomoxis annularis</i> | 40 |
| | <i>Pomoxis nigromaculatus</i> | 314 |
| 858 <i>Fillicollis anatis</i> | <i>Anas cyanoptera septentrionalium</i> | 499 |
| 859 <i>Gracilisentis gracilisentis</i> | <i>Dorosoma cepedianum</i> | 241 |
| | <i>Dorosoma petenese</i> | 241 |
| 860 <i>Hamanniella</i> | Armadillo | 96 |

| | | | |
|-----|---|-------------------------------|---------------|
| 861 | <i>Leptorhynchoides thecatus</i> | Ambloplites rupestris | 479 |
| | | Astyanax mexicanus | 479 |
| | | Cichlosoma cyanoguttatum | 479 |
| | | Gambusia geiseri | 127 |
| | | Ictalurus nebulosus | 479 |
| | | Lepomis auritus | 479 |
| | | Lepomis cyanellus | 479 |
| | | Lepomis gulosus | 479 |
| | | Lepomis macrochirus | 479 |
| | | Lepomis microlophus | 479 |
| | | Lepomis punctatus | 479 |
| | | Micropterus punctulatus | 479 |
| | | Micropterus salmoides | 479 |
| 862 | <i>Macracanthorhynchus hirudinaceus</i> | Pig | 35 |
| 863 | <i>Macracanthorhynchus ingens</i> | Bassariscus astutus | 362 |
| | | Procyon lotor | 93,331 |
| 864 | <i>Mediorhynchus</i> | Quiscalus quiscula | 32 |
| 865 | <i>Mediorhynchus colini</i> | Colinus virginianus texanus | 493 |
| 866 | <i>Mediorhynchus grandis</i> | Agelaius phoeniceus | 332 |
| | | Corvus brachyrhynchos | 341 |
| | | Corvus cryptoleucus | 371N |
| | | Molothrus ater ater | 332 |
| | | Passerherbulus caudacutus | 332 |
| | | Sturnella magna magna | 284 |
| 867 | <i>Mediorhynchus papillosus</i> | Numenius americanus | 70N |
| 868 | <i>Mediorhynchus robustum</i> | Sturnus vulgaris | 43 |
| 869 | <i>Moniliformis</i> | Rattus rattus | 77 |
| 870 | <i>Moniliformis clarki</i> | Cynomys ludovicianus | 372N |
| | | Erinaceus albiventris | 433 |
| | | Homo sapiens | 433 |
| | | Loris gracilis | 433 |
| | | Rattus norvegicus | 433 |
| | | Rattus rattus | 77,433,480 |
| | | Scalopus aquaticus | 433 |
| | | Sciurus niger | 433 |
| | | Spermophilus tridecemlineatus | 433 |
| 871 | <i>Moniliformis dubius</i> | Periplaneta americana | 330 |
| | | Rattus rattus | 87 |
| | | Sigmodon hispidus | 330 |
| 872 | <i>Neoechinorhynchus constrictus</i> | Pseudemys scripta elegans | 245,415 |
| | | Gambusia affinis | 128 |
| | | Huro salmoides | 434 |
| | | Lepomis cyanellus | 308,309,310 |
| | | Lepomis humilis | 308 |
| | | Lepomis macrochirus | 280,308 |
| | | Lepomis megalotis | 308 |
| | | Micropterus salmoides | 20,23,280,309 |
| | | Pomoxis annularis | 20,280,309 |
| 873 | <i>Neoechinorhynchus clyindratum</i> | Ambloplites rupestris | 479 |
| | | Astyanax mexicanus | 479 |
| | | Cichlosoma cyanoguttatum | 479 |
| | | Gambusia affinis | 127 |

| | | | |
|-----|--------------------------------------|---------------------------|---------------------|
| | Gambusia geiseri | 127 | |
| | Lepisosteus oculatus | 479 | |
| | Lepomis auritus | 479 | |
| | Lepomis cyanellus | 479 | |
| | Lepomis gulosus | 211,479 | |
| | Lepomis macrochirus | 211,479 | |
| | Lepomis marginatus | 40 | |
| | Lepomis megalotis | 40 | |
| | Lepomis microlophus | 479 | |
| | Lepomis punctatus | 479 | |
| | Micropterus punctulatus | 40,211,479 | |
| | Micropterus salmoides | 40,211,252, 479 | |
| | Spinitectus carolini | 40 | |
| 874 | <i>Neoechinorhynchus emydis</i> | Pseudemys scripta elegans | 216 |
| 875 | <i>Neoechinorhynchus emyditoides</i> | Pseudemys scripta elegans | 191,245 |
| 876 | <i>Neoechinorhynchus prolixus</i> | Carpiodes carpio | 420,481 |
| 877 | <i>Octospiniferoides chandleri</i> | Fundulus grandis | 66 |
| 878 | <i>Oncicola canis</i> | Canis latrans | 364,365,370, 381 |
| | | Dog | 378 |
| | | Felis rufus | 365,440 |
| 879 | <i>Oncicola canis larva</i> | Armadillo | 96 |
| 880 | <i>Pachysentis canicola</i> | Canis latrans | 364,365 |
| | | Urocyon cinereoargenteus | 65 |
| 881 | <i>Plagiorhynchus formosus</i> | Sturnus vulgaris | 43 |
| 882 | <i>Polymorphus</i> | Anas clypeata | 58 |
| 883 | <i>Polymorphus minutus</i> | Anas creca | 74 |
| 884 | <i>Polymorphus texensis</i> | Crocethia alba | 494 |
| 885 | <i>Pomphorhynchus bulbocolli</i> | Amblopites rupestris | 479 |
| | | Astyanax mexicanus | 479 |
| | | Lepomis auritus | 479 |
| | | Lepomis macrochirus | 479 |
| | | Lepomis microlophus | 479 |
| | | Lepomis punctatus | 479 |
| | | Moxostoma congestum | 479 |
| 886 | <i>Tanaorhamphus longirostris</i> | Dorosoma cepedianum | 241 |
| | | Dorosoma petenese | 241 |

Arthropoda-Acarina-Mite

| | | | |
|-----|-----------------------------------|-----------------------|---------|
| 887 | <i>Allodermanyssus sanguineus</i> | Mus musculus | 376A |
| 888 | <i>Amblyomma americanum</i> | Sigmodon hispidus | 250 |
| 889 | <i>Androlaelaps</i> | Perognathus flavus | 165 |
| | | Perognathus hispidus | 220 |
| 890 | <i>Androlaelaps farenholzi</i> | Brassariscus astutus | 120 |
| 891 | <i>Androlaelaps grandiculatus</i> | Perognathus flavus | 167,174 |
| | | Perognathus hispidus | 174 |
| | | Reithrodontomys | 167,174 |
| 892 | <i>Androlaelaps johnstoni</i> | Neotoma micropus | 170 |
| 893 | <i>Androlaelaps leviculus</i> | Onychomys leucogaster | 167,174 |

| | | |
|---|---------------------------|----------|
| | Perognathus hispidus | 167 |
| | Sigmodon hispidus | 167 |
| 894 <i>Aponomma elaphensis</i> | Elaphe subocularis | 133 |
| 895 <i>Astrida coccyzae</i> | Coccyzus americanus | 359 |
| 896 <i>Atricholaelaps glasgowi</i> | Sigmodon hispidus | 250 |
| | Urocyon cinereoargenteus | 166 |
| 897 <i>Agricholaelaps megaventralis</i> | Gray Squirrel | 442 |
| | Mus musculus | 442 |
| | Rattus norvegicus | 442 |
| | Rattus rattus | 250,323 |
| | Sciurus carolinensis | 442 |
| | Sigmodon hispidus | 442 |
| | Sturnus vulgaris | 43 |
| | Urocyon cinereoargenteus | 166 |
| 898 <i>Atricholaelaps sigmodoni</i> | Baiomys taylori | 385 |
| | Geomys bursarius | 385 |
| | Neotoma floridana | 385 |
| | Perognathus | 385 |
| | Peromyscus leucopus | 385 |
| | Sciurus carolinensis | 385 |
| | Sciurus niger | 385 |
| | Sigmodon hispidus | 385,441 |
| | Sylvilagus floridanus | 385 |
| 899 <i>Atyeous texanus</i> | Buteo swainsoni | 181 |
| 900 <i>Bdellonyssus bacoti</i> | Baiomys taylori | 220 |
| | Mus musculus | 220 |
| | Neotoma micropus | 174 |
| | Peromyscus boylii | 220 |
| | Peromyscus leucopus | 220 |
| | Peromyscus maniculatus | 220 |
| | Peromyscus pectoralis | 220 |
| | Rattus norvegicus | 174,448 |
| 901 <i>Boydaia pheucticola</i> | Pheucticus melanocephalus | 359 |
| 902 <i>Boydaia tyrannis</i> | Tyrannus verticalis | 59 |
| | Myiarchus cinerascens | 359 |
| 903 <i>Brevisterna morlani</i> | Cynomys gunnisoni | 451,451N |
| | Dipodomys ordi | 451,451N |
| | Neotoma | 451 |
| | Neotoma albigula | 451 |
| | Neotoma micropus | 451 |
| | Peromyscus leucopus | 451 |
| | Peromyscus maniculatus | 451 |
| | Peromyscus truei | 451 |
| | Sylvilagus auduboni | 451 |
| 904 <i>Brevisterna utahensis</i> | Neotoma | 451 |
| | Neotoma micropus | 174 |
| 905 <i>Cheyletiella</i> | Sturnus vulgaris | 43 |
| 906 <i>Cheyletus eruditus</i> | Bassariscus astutus | 120 |
| 907 <i>Chiroptonyssus robustipes</i> | Myotis velifer | 220 |
| 908 <i>Chiroptonyssus texensis</i> | Tadarida brasiliensis | 28,385 |
| 909 <i>Damalinia parallelus</i> | White-tailed Deer | 42A |
| 910 <i>Degeerriella nebulosa</i> | Sturnus vulgaris | 75 |
| 911 <i>Demodex</i> | Goat | 34,34N |
| | Horse | 34N,A |
| 912 <i>Demdex bovis</i> | Cattle | 192N |

| | | | |
|-----|---|-----------------------------------|---------|
| 913 | <i>Dermacarus hypudaei</i> | <i>Bassariscus astutus</i> | 120 |
| 914 | <i>Dermannyssidae</i> | <i>Perognathus hispidus</i> | 316 |
| | | <i>Sigmodon hispidus</i> | 316 |
| 915 | <i>Dermanyssus gallinae</i> | <i>Myotis velifer</i> | 204 |
| | | <i>Pipistrellus subflavus</i> | 204 |
| | | <i>Rattus rattus</i> | 385 |
| | | <i>Sturnus vulgaris</i> | 43 |
| 916 | <i>Dermanyssus prognepphilus</i> | <i>Sturnus vulgaris</i> | 43 |
| 917 | <i>Dermonoton serratus</i> | <i>Strix varia</i> | 181 |
| 918 | <i>Echidnolaelaps echidninus</i> | <i>Peromyscus boylii</i> | 220 |
| | | <i>Rattus norvegicus</i> | 182 |
| | | <i>Rattus rattus</i> | 385 |
| | | <i>Sylvilagus floridanus</i> | 385 |
| 919 | <i>Entonyssus</i> | <i>Crotalus atrox</i> | 203 |
| 920 | <i>Eubrachylaelaps crowei</i> | <i>Dipodomys spectabilis</i> | 174 |
| | | <i>Onychomys leucogaster</i> | 174 |
| 921 | <i>Eucheyletia hardyi</i> | <i>Bassariscus astutus</i> | 120 |
| 922 | <i>Eulaelaps stabularis</i> | <i>Baimoys taylori</i> | 385 |
| | | <i>Didelphis virginiana</i> | 385 |
| | | <i>Perognathus</i> | 385 |
| | | <i>Peromyscus leucopus</i> | 385 |
| | | <i>Urocyon cinereoargenteus</i> | 272 |
| 923 | <i>Euschoengastia bigenuala</i> | <i>Liomys irroratus</i> | 180,293 |
| | | <i>Perognathus penicillatus</i> | 293 |
| | | <i>Sigmodon</i> | 187 |
| 924 | <i>Euschoengastia eadsi</i> | <i>Bassariscus astutus</i> | 120 |
| 925 | <i>Euschoengastia hoplai</i> | <i>Perognathus flavus</i> | 293 |
| 926 | <i>Euschoengastia lacerta</i> | <i>Liomys irroratus</i> | 180,293 |
| 927 | <i>Euschoengastia loomisi</i> | <i>Liomys irroratus</i> | 180,282 |
| | | <i>Neotoma micropus</i> | 293 |
| | | <i>Peromyscus leucopus</i> | 293 |
| | | <i>Sigmodon hispidus</i> | 293 |
| | | <i>Spermophilus mexicanus</i> | 293 |
| 928 | <i>Eutrombicula alfreddugesi</i> | <i>Cnemidophorus sacki</i> | 293 |
| | | <i>Cnemidophorus sexlineatus</i> | 293 |
| | | <i>Cophosaurus texanus</i> | 293 |
| | | <i>Liomys irroratus</i> | 180 |
| | | <i>Neotoma micropus</i> | 293 |
| | | <i>Peromyscus leucopus</i> | 293 |
| | | <i>Phrynosoma cornutum</i> | 293 |
| | | <i>Scleoporus olivaceus</i> | 293 |
| | | <i>Sigmodon hispidus</i> | 293 |
| | | <i>Sylvilagus auduboni</i> | 293 |
| 929 | <i>Eutrombicula masoni</i> | <i>Rattus norvegicus</i> | 448 |
| 930 | <i>Gabucinia tistata</i> | <i>Buteo jamaicensis borealis</i> | 181A |
| 931 | <i>Gigantolaelaps crescetidarum</i> | <i>Oryzomys palustris</i> | 333 |
| 932 | <i>Haemolaelaps glasgowi</i> | Bird | 445 |
| | | <i>Cratogeomys castanops</i> | 165 |
| | | <i>Cyomys ludovicianus</i> | 174 |
| | | <i>Dipodomys spectabilis</i> | 174 |
| | | <i>Liomys irroratus</i> | 180 |
| | | Mammal | 445 |
| | | <i>Neotoma micropus</i> | 174 |
| | | <i>Onychomys leucogaster</i> | 174 |
| | | <i>Peromyscus boylii</i> | 176 |

| | | |
|--|-----------------------------|--------------|
| | Rattus norvegicus | |
| 174,376,376N,A | | |
| | Rattus rattus | 376,376NA |
| | Spermophilus pilosoma | 174 |
| | Spermophilus tereticaudus | 279A |
| | Spermophilus variegatus | 165 |
| 933 <i>Haemolaelaps megaventralis</i> | Bird | 445 |
| | Mammal | 445 |
| | Petrochelidon fulva pallida | 177 |
| | Rattus norvegicus | 174 |
| 934 <i>Hannemania amphili</i> | Bufo cognatus | 155N |
| | Bufo debilis | 155N |
| | Bufo microscaphus | 155N |
| | Bufo punctatus | 155N |
| | Bufo speciosus | 155 |
| | Bufo woodhousei | 155N |
| | Hyla arenicolor | 155N |
| | Scaphiopus bombifrons | 155N |
| | Scaphiopus hammondi | 155N |
| 935 <i>Hannemania eltoni</i> | Acris crepitans | 49 |
| 936 <i>Hirstionyssus</i> | Cratogeomys castanops | 165 |
| | Liomys irroratus | 180 |
| 937 <i>Hirstionyssus breviseta</i> | Bassariscus astutus | 120 |
| | Neotoma albigula | 449N |
| 938 <i>Hirstionyssus liomys</i> | Liomys irroratus | 223 |
| 939 <i>Hirstionyssus neotomae</i> | Liomys irroratus | 180 |
| | Neotoma albigula | 449N |
| 940 <i>Hirstionyssus staffordi</i> | Bassariscus astutus | 120 |
| | Mephitis mephitis | 447 |
| 941 <i>Hypoaspis dasypus</i> | Dasypus novemcinctus | 323 |
| 942 <i>Ichoronyssus</i> | Leptonycteris sanborni | 54A |
| | Pipistrellus hesperus | 54A |
| | Tadarida femorosacca | 54A |
| 943 <i>Ichoronyssus crosbyi</i> | Myotis velifer | 54A |
| 944 <i>Ichoronyssus dentipes</i> | Sigmodon hispidus | 443 |
| 945 <i>Ichoronyssus longisetosus</i> | Myotis velifer | 254 |
| | Plecotus townsendi | 254 |
| 946 <i>Ichoronyssus robustipes</i> | Tadarida brasiliensis | 54A,254,254N |
| 947 <i>Ischyropoda armatus</i> | Dipodomys ordi | 174 |
| | Dipodomys spectabilis | 174 |
| | Neotoma micropus | 174 |
| | Onychomys leucogaster | 174 |
| | Sylvilagus auduboni | 174 |
| 948 <i>Ixobioides liponyssoides</i> | Crotalus atrox | 485 |
| | Thamnophis proximus | 485 |
| 949 <i>Kramerella lunulata</i> <i>flammei</i> | Asio flammeus | 181 |
| 950 <i>Laelaps echidninus</i> | Rattus norvegicus | 376,448 |
| | Rattus rattus | 376 |
| 951 <i>Laelaps hawaiiensis</i> | Rattus rattus | 384 |
| 952 <i>Laelaps nuttalia</i> | Mus musculus | 220 |
| | Rattus norvegicus | 376 |
| | Rattus rattus | 376 |
| 953 <i>Laelaps nuttalli</i> | Rattus norvegicus | 448 |
| 954 <i>Laelaps stabularis</i> | Rattus rattus | 385 |
| 955 <i>Larinyssus orbicularis</i> | Larus argentatus | 444 |

| | | |
|--|------------------------------|----------|
| | Larus atricilla | 444 |
| | Larus delawarensis | 444 |
| | Sterna maxima | 444 |
| | Thalasseus maximus | 326 |
| 956 <i>Leptotrombidium panamensis</i> | Dipodomys merriami | 293 |
| | Liomys irroratus | 180,293 |
| | Peromyscus leucopus | 293 |
| | Spermophilus mexicanus | 293 |
| 957 <i>Liponyssus</i> | Perognathus hispidus | 316 |
| | Sigmodon hispidus | 316 |
| 958 <i>Liponyssus bacoti</i> | Didelphis virginiana | 385 |
| | Man | 47 |
| | Mus musculus | 385 |
| | Neotoma floridana | 385 |
| | Rat | 141 |
| | Rattus norvegicus | 47 |
| | Rattus rattus | 250,385 |
| | Sigmodon hispidus | 385 |
| | Urocyon cinereoargenteus | 166 |
| 959 <i>Liponyssus haemotophagus</i> | Pipistrellus hesperus | 165 |
| 960 <i>Liponyssus sylvarum</i> | Sternus vulgaris | 43 |
| 961 <i>Listrophorus</i> | Liomys irroratus | 180 |
| | Sylvilagus floridanus | 385 |
| 962 <i>Listrophorus klebergi</i> | Sigmodon hispidus | 304 |
| 963 <i>Macrocheles</i> | Rattus norvegicus | 174 |
| | Speotyto cunicularia | 174 |
| | Spermophilus spilosoma | 174 |
| 964 <i>Microtrombicula crossleyi</i> | Peromyscus leucopus | 293 |
| 965 <i>Microtrombicula merrhewi</i> | Tadarida brasiliensis | 130,130A |
| 966 <i>Neoichoronyssus incomptus</i> | Dipodomys ordi | 173,174 |
| 967 <i>Neoichoronyssus neotomae</i> | Neotoma micropus | 171 |
| 968 <i>Neoichoronyssus triacanthus</i> | Dipodomys ordi | 174 |
| 969 <i>Neonyssus belopolski</i> | Butorides virescens | 326 |
| 970 <i>Neonyssus hirsti</i> | Passer domesticus | 375 |
| 971 <i>Neonyssus triangulus</i> | Zenaida asiatica | 454 |
| 972 <i>Neonyssus zenaidurae</i> | Columbia livia domestica | 118 |
| | Columbigallina passerina | 347 |
| | Zenaida macroura | 347 |
| 973 <i>Neoschoengastea americana</i> | Amphispiza bilineata | 293 |
| | Chordeiles acutipennis | 293 |
| | Cynomys ludovicianus | 293 |
| | Geococcyx californianus | 174,293 |
| | Heleodytes brunneicapillus | 293 |
| | Lepus californicus | 293 |
| | Parabuteo unicinctus harrisi | 293 |
| | Salpinctes obsoletus | 293 |
| | Sylvilagus | 293 |
| | Thryomanes bewikii | 293 |
| | Toxosoma curvirostre | 293 |
| | Tyrannus melancholicus | 293 |
| 974 <i>Neotrombicula whartoni</i> | Lepus californicus | 293 |
| | Sigmodon hispidus | 293 |
| 975 <i>Nycteriglyphus bifolium</i> | Tadarida brasiliensis | 455 |
| 976 <i>Odontacarus arizonensis</i> | Crotaphytus collaris | 293 |
| 977 <i>Odontacarus cayolargoensis</i> | Liomys irroratus | 180,293 |

| | | |
|---|------------------------------|---------|
| | Peromyscus leucopus | 293 |
| | Sigmodon hispidus | 293 |
| 978 <i>Odontacarus galli</i> | Cynomys ludovicianus | 293 |
| | Lepus californicus | 293 |
| | Sylvilagus auduboni | 293 |
| 979 <i>Odontacarus plumosus</i> | Liomys irroratus | 180,293 |
| 980 <i>Olabidocarpus americanus</i> | Lasiurus intermedius | 303 |
| 981 <i>Ophthalmophagus striatus</i> | Columbigallina sparverius | 359 |
| 982 <i>Oribatidae</i> | Baiomys taylori | 316 |
| 983 <i>Ornithonyssus bacoti</i> | Liomys irroratus | 180 |
| 984 <i>Ornithonyssus sylvarium</i> | Liomys irroratus | 180 |
| | Eptesicus fuscus | 204 |
| | Myotis velifer | 204 |
| 985 <i>Paraneonyssus icteridius</i> | Molothrus ater | 452 |
| 986 <i>Passerculus japuibensis</i> | Dendroica auduboni | 202 |
| | Mniotilta varia | 202 |
| | Wilsonia citrina | 202 |
| | Wilsonia pusilla | 202 |
| 987 <i>Passerculus sandwichensis</i> | Melospiza lincolni | 202 |
| 988 <i>Periglischrus strandtmanni</i> | Mormoops megalophylla | 254,465 |
| 989 <i>Pneumocoptes penrosei</i> | Cynomys ludovicianus | 35 |
| 990 <i>Proctolaelaps</i> | Cynomy ludovicianus | 372N |
| 991 <i>Protalgas attenuatus</i> | Bubo virginianus | 181 |
| 992 <i>Pseudaloptinus milvulinus</i> | Parabuteo unicinctus harrisi | 181 |
| 993 <i>Pseudoschoengastia andyi</i> | Liomys irroratus | 180 |
| | Peromyscus leucopus | 293 |
| 994 <i>Pseudoschoengastia apista</i> | Bassariscus astutus | 120 |
| 995 <i>Pseudoschoengastia farneri</i> | Liomys irroratus | 180,293 |
| | Peromyscus leucopus | 293 |
| 996 <i>Pseudoschoengastia hungerfordi</i> | Perognathus hispidus | 293 |
| | Sylvilagus auduboni | 293 |
| 997 <i>Psoroptes</i> | Rattus norvegicus | 448 |
| 998 <i>Psoroptes ovis</i> | Sheep | 48N |
| 999 <i>Psoroptic cattle scabies</i> | Cattle | 33 |
| 1000 <i>Pteracarus chalinolobus</i> | Plecotus rafinesqui | 204 |
| 1001 <i>Ptilonyssus echinatus</i> | Petrochelidon pyrrhonota | 202 |
| 1002 <i>Ptilonyssus lanii</i> | Lanius ludovicianus | 202 |
| 1003 <i>Ptilonyssus mimi</i> | Mimus polyglottos | 202 |
| 1004 <i>Ptilonyssus nudus</i> | Passer domesticus | 375 |
| 1005 <i>Ptilonyssus perisorel</i> | Cyanocitta stelleri | 202N |
| | Perisoreus canadensis | 202N |
| 1006 <i>Ptilonyssus phainopeplae</i> | Phainopepla nitens | 202 |
| 1007 <i>Ptilonyssus richmondeniensis</i> | Richmondia cardinalis | 202 |
| 1008 <i>Ptilonyssus sairae</i> | Amphispiza bilineata | 202 |
| | Calamospiza melanocorys | 202 |
| | Chondestes grammacus | 202 |
| | Passerina cyanea | 202 |
| | Pipilo fuscus | 202N |
| | Poocetes gramineus | 202 |
| | Spizella passerina | 202 |
| 1009 <i>Ptilonyssus salpinctis</i> | Salpinctes obsoletus | 202 |
| 1010 <i>Ptilonyssus sialiae</i> | Sialia currucoides | 202 |

| | | | |
|------|-----------------------------------|-----------------------------------|---------|
| 1011 | <i>Radfordia ensifera</i> | <i>Rattus norvegicus</i> | 448 |
| 1012 | <i>Radfortiella</i> | <i>Leptoncyteris nivalis</i> | 373 |
| 1013 | <i>Rallinyssus candistigmus</i> | <i>Fulica americana</i> | 444 |
| | | <i>Rallus elegans</i> | 444 |
| 1014 | <i>Rhinonyssus coniventris</i> | <i>Arenaria interpres</i> | 446,450 |
| | | <i>Catoprophorus semipalmatus</i> | 446,450 |
| | | <i>Charadrius hiaticula</i> | 446,450 |
| | | <i>Crocathia alba</i> | 446,450 |
| | | <i>Erolia alpina</i> | 450 |
| | | <i>Erolia maritima</i> | 450 |
| | | <i>Squatarola squatarola</i> | 326 |
| | | <i>Totanus calidris</i> | 450 |
| | | <i>Totanus flavipes</i> | 446,450 |
| 1015 | <i>Rhinonyssus himantopus</i> | <i>Charadrius vociferus</i> | 453 |
| | | <i>Recurvirostra americana</i> | 453 |
| 1016 | <i>Rhinonyssus rhinolethrum</i> | <i>Anas acuta</i> | 446,450 |
| | | <i>Anas carolinensis</i> | 446 |
| | | <i>Anas clypeata</i> | 446,450 |
| | | <i>Anas platyrhynchos</i> | 446,450 |
| | | <i>Anas strepera</i> | 446,450 |
| | | Anatidae | 327 |
| | | <i>Aythya affinis</i> | 446,450 |
| | | <i>Branta canadensis</i> | 446,450 |
| | | <i>Mareca americana</i> | 446,450 |
| | | <i>Mergus</i> | 450 |
| 1017 | <i>Sarcoptes</i> | <i>Rattus norvegicus</i> | 448 |
| 1018 | <i>Sarcoptes myotis</i> | <i>Myotis velifer</i> | 54A,219 |
| 1019 | <i>Speleognathopsis benoiti</i> | Black-bellied Plover | 306 |
| 1020 | <i>Speleognathus australis</i> | <i>Bison bison</i> | 149 |
| 1021 | <i>Speleognathus striatus</i> | <i>Columba livia domestica</i> | 118 |
| 1022 | <i>Speleognathus sturni</i> | <i>Passer domesticus</i> | 375 |
| 1023 | <i>Spinturnix</i> | <i>Antrozous pallidus</i> | 54A |
| | | <i>Eptesicus fuscus</i> | 54A |
| | | <i>Myotis velifer</i> | 254 |
| | | <i>Pipistrellus hesperus</i> | 165 |
| 1024 | <i>Spinturnix americanus</i> | <i>Myotis leibii</i> | 54A |
| 1025 | <i>Spinturnix carloshoffmanni</i> | <i>Myotis leibii</i> | 54A |
| | | <i>Myotis thysanodes</i> | 54A |
| | | <i>Myotis velifer</i> | 54A,402 |
| | | <i>Tadarida brasiliensis</i> | 402 |
| 1026 | <i>Steatonyssus joaquimi</i> | <i>Eptesicus fuscus</i> | 254 |
| 1027 | <i>Sternosoma augei</i> | <i>Speotyto cunicularia</i> | 357 |
| 1028 | <i>Sternosoma crotophagae</i> | <i>Crotophaga sulcirostris</i> | 357 |
| 1029 | <i>Sternosoma pirangae</i> | <i>Piranga rubra</i> | 357 |
| 1030 | <i>Sternosoma porteri</i> | <i>Centurus aurifrons</i> | 357 |
| 1031 | <i>Sternosoma sayornis</i> | <i>Sayornis nigricans</i> | 357 |
| 1032 | <i>Sternosoma tracheacolum</i> | <i>Pipilo fuscus</i> | 357 |
| | | <i>Pyrrhuloxia sinuata</i> | 357 |
| 1033 | <i>Sternostoma boydi</i> | <i>Calidris canutus</i> | 326 |
| | | <i>Crocathia alba</i> | 327 |
| 1034 | <i>Sternostoma strandtmanni</i> | <i>Cassidix mexicanus</i> | 326 |
| 1035 | <i>Sternostoma tyrannus</i> | <i>Nuttaliornis borealis</i> | 59 |
| 1036 | <i>Streptolaelaps liomydis</i> | <i>Liomys irroratus</i> | 180 |
| 1037 | <i>Tecomatlana sandovali</i> | <i>Macrotus californicus</i> | 54A |
| 1038 | <i>Teinocoptes lasionycteris</i> | <i>Plecotus rafinesqui</i> | 204 |
| | | <i>Tadarida brasiliensis</i> | 204 |

| | | | |
|------|---|---------------------------|------|
| 1039 | <i>Tinaminyssus juxtamelloi</i> | Columba fasciata | 358N |
| 1040 | <i>Trichodectes pallidus</i> | Coati | 42A |
| 1041 | <i>Trombicula</i> | Liomys irroratus | 180 |
| | | Mormoops megalophylla | 254 |
| | | Tadarida brasiliensis | 254N |
| 1042 | <i>Trombicula allredi</i> | Liomys irroratus | 180 |
| 1043 | <i>Trombicula (Eutrombicula) alfreddugesi</i> | Cnemidophorus sexlineatus | 203 |
| | | Cnemidophorus tessellatus | 203 |
| | | Cnemidophorus tigris | 203 |
| | | Crotaphytus collaris | 203 |
| | | Heterodon nasicus | 203 |
| | | Sceloporus olivaceous | 203 |
| | | Uta stansburiana | 203 |
| 1044 | <i>Trombicula arenicola</i> | Spermophilus tereticaudus | 279A |
| 1045 | <i>Trombicula gurneyi</i> | Peromyscus leucopus | 293 |
| | | Sigmodon hispidus | 293 |
| 1046 | <i>Trombicula mexicana</i> | Myotis velifer | 54A |
| 1047 | <i>Trombicula montanensis</i> | Cynomys ludovicianus | 174 |
| | | Spermophilus mexicanus | 293 |
| 1048 | <i>Trombicula myotis</i> | Myotis leibii | 54A |
| | | Myotis thysanodes | 54A |
| | | Myotis velifer | 54A |
| 1049 | <i>Trombicula tibbettsi</i> | Mormoops megalophylla | 293 |
| 1050 | <i>Trouessartia rosterii</i> | Sturnus vulgaris | 43 |
| 1051 | <i>Tyranninyssus callinectodies</i> | Myiarchus cinerascens | 59 |
| 1052 | <i>Tyranninyssus spinosus</i> | Muscivora forficata | 59 |
| | | Tyrannus dominicensis | 59 |
| | | Tyrannus tyrannus | 59 |
| | | Tyrannus verticalis | 59 |
| 1053 | <i>Tyranninyssus tyrannisoides</i> | Empidonax minimus | 59 |
| | | Sayornis saya | 59 |
| 1054 | <i>Tyranninyssus tyrannus</i> | Nuttallornis borealis | 59 |
| 1055 | <i>Uropoda</i> | Dipodomys ordi | 174 |
| | | Neotoma micropus | 174 |
| | | Onychomys leucogaster | 174 |
| 1056 | <i>Whartonia perplexa</i> | Antrozous pallidus | 54A |
| | | Eptesicus fuscus | 254 |
| | | Pipistrellus hesperus | 54A |
| 1057 | <i>Whartonia whartoni</i> | Neotoma micropus | 293 |
| | | Peromyscus | 293 |
| 1058 | <i>Womersia strandtmani</i> | Least Tern | 293 |
| | | Sterna antillarum | 293 |

Arthropoda-Acarina-Ticks

| | | | |
|------|-----------------------------|--------------------------|---------------|
| 1059 | <i>Amblyomma americanum</i> | Aimophila ruficeps | 55,56 |
| | | Antrostomus carolinensis | 55,56 |
| | | Baelophus atricristatus | 55,56 |
| | | Baiomys taylori | 385 |
| | | Bassariscus astutus | 55,56,120,470 |

| | |
|---------------------------------|---------------------------|
| Bobwhite Quail | 490 |
| <i>Bos taurus</i> | 220 |
| <i>Buteo borealis</i> | 55,56 |
| <i>Buteo swainsoni</i> | 55,56 |
| <i>Canis familiaris</i> | 220,385 |
| <i>Canis latrans</i> | 158 |
| <i>Canis rufus</i> | 385 |
| Carnivore | 178 |
| <i>Carpodacus mexicanus</i> | 55,56 |
| <i>Cathartes aura</i> | 55,56 |
| <i>Catherpes mexicanus</i> | 55,56 |
| Cattle | 152 |
| <i>Chondestes grammacus</i> | 55,56 |
| <i>Coccyzus americanus</i> | 56 |
| <i>Colinus virginianus</i> | 55,56,355 |
| <i>Columbigallina passerina</i> | 55,56 |
| <i>Conepatus mesoleucus</i> | 55,56 |
| <i>Dasypus novemcinctus</i> | 55,56,220 |
| Deer | 490 |
| <i>Dendrocia caerulescens</i> | 55,56 |
| <i>Didelphis virginiana</i> | 55,56 |
| Domestic Animal | 178 |
| <i>Empidonax minimus</i> | 55,56 |
| Fringillidae | 55,56 |
| <i>Geococcyx californianus</i> | 55,56 |
| Gray Fox | 490 |
| Ground Bird | 178 |
| <i>Guiraca caerulea</i> | 55,56 |
| <i>Icteria virens</i> | 55,56 |
| Jack Rabbit | 490 |
| <i>Lanius ludovicianus</i> | 55,56 |
| <i>Lepus californicus</i> | 55,56 |
| Man | 490 |
| <i>Meleagris gallopavo</i> | 55,56 |
| <i>Mephitis mephitis</i> | 55,56,220 |
| <i>Mimus polyglottos</i> | 55,56 |
| <i>Muscivora forficata</i> | 55,56 |
| <i>Myiarchus crinitus</i> | 55,56 |
| <i>Myiochanes richardsonii</i> | 55,56 |
| <i>Neotoma floridana</i> | 341 |
| <i>Odocoileus virginianus</i> | 55,56,183, 264,385,407 |
| <i>Passerina ciris</i> | 55,56 |
| Perognathus | 385 |
| <i>Peromyscus leucopus</i> | 385 |
| <i>Phalaenoptilus nuttallii</i> | 55,56 |
| <i>Piranga rubra</i> | 55,56 |
| <i>Procyon lotor</i> | 55,56,385 |
| Rabbit | 178 |
| <i>Rattus rattus</i> | 55,56 |
| <i>Richmondia cardinalis</i> | 55,56 |
| Rodent | 178 |
| <i>Sciurus carolinensis</i> | 385 |
| <i>Sciurus niger</i> | 55,56,385 |
| <i>Sigmodon hispidus</i> | 55,55,385 |
| <i>Spermophilus mexicanus</i> | 55,56 |

| | | |
|------|---------------------------------------|---------------------|
| | Spermophilus variegatus | 55,56 |
| | Spinus psaltria | 55,56 |
| | Spizella pusilla | 56 |
| | Sus scrofa | 220 |
| | Sylvilagus floridanus | 55,56 |
| | Tayassu tajacu | 168 |
| | Thryomanes bewickii | 55,56 |
| | Thryothorus ludovicianus | 55,56 |
| | Urocyon cinereoargenteus | 55,56,166, 385 |
| | Wolf | 158 |
| | Zenaida macroura | 55,56 |
| 1060 | <i>Amblyomma cajennense</i> | |
| | Cow | 178 |
| | Dog | 178 |
| | Odocoileus virginianus | 178,407 |
| | Quail | 178 |
| | Tayassu tajacu | 168,178,409 |
| | Sus scrofa | 113 |
| 1061 | <i>Amblyomma cajennense maculatum</i> | |
| 1062 | <i>Amblyomma dissimile</i> | Boa Constrictor 178 |
| 1063 | <i>Amblyomma humerale</i> | Box Turtle 178 |
| 1064 | <i>Amblyomma imitator</i> | Burro 273,273N |
| | Cattle | 273,273N |
| | Deer | 273,273N |
| | Geococcyx californianus | 273,273N |
| | Goat | 273,273N |
| | Horse | 273,273N |
| | Man | 273,273N |
| | Meleagris gallopavo | 273,273N |
| | Opossum | 273,273N |
| | Peccary | 273,273N |
| | Sciurus aureogaster | 273,273N |
| 1065 | <i>Amblyomma inornatum</i> | |
| | Canis latrans | 158,178 |
| | Cotton Rat | 178 |
| | Cottontail Rabbit | 178 |
| | Dasypus novemcinctus | 178 |
| | Dog | 178 |
| | Jack Rabbit | 55,178 |
| | Felis rufus | 178 |
| | Mephitis mephitis | 178 |
| | Neotoma micropus | 178 |
| | Odocoileus virginianus | 407 |
| | Procyon lotor | 178 |
| | Tayassu tajacu | 168,178,409 |
| 1066 | <i>Amblyomma longirostre</i> | Viero griseus 178 |
| 1067 | <i>Amblyomma maculatum</i> | |
| | Canis familiaris | 385 |
| | Canis latrans | 158 |
| | Cattle | 152,178 |
| | Coyote | 178 |
| | Dog | 178 |
| | Odocoileus virginianus | 264,407 |
| | Quail | 178 |
| | Tayassu tajacu | 168 |
| | White-tailed Deer | 178 |
| 1068 | <i>Amblyomma philipi</i> | |
| | Canis | 111 |
| | Lepus | 111 |

| | | |
|--|-------------------------------|------------------|
| | Rabbit | 111 |
| | Sylvilagus | 111 |
| 1069 <i>Anocentor nitens</i> | Horse | 178 |
| | White-tailed Deer | 178 |
| 1070 <i>Antricola coprophilus</i> | Bat | 162 |
| | Tadarida brasiliensis | 178 |
| 1071 <i>Aponomma elaphensis</i> | Elaphe subocularis | 380 |
| 1072 <i>Argas megnini</i> | Horse | 424 |
| 1073 <i>Argas periscus</i> | Chicken | 61,178 |
| | Geococcyx californianus | 178 |
| | Black Vulture | 55 |
| 1074 <i>Argasid larva</i> | Crotaphytus collaris | 203 |
| 1075 <i>Boophilus annulatus</i> | Cattle | 178 |
| | Odocoileus virginianus | 264 |
| 1076 <i>Boophilus bovis</i> | Cow | 431 |
| 1077 <i>Ceratophyllus celsus</i> | Petrochelidon albifrons | 177 |
| 1078 <i>Ceratophyllus coahuilensis</i> | Petrochelidon fulva pallida | 177 |
| 1079 <i>Dermacentor albipictus</i> | Cattle | 152 |
| | Cervus elaphus | 501N |
| | Deer | 55 |
| | Horse | 178 |
| | Mule Deer | 178 |
| | Odocoileus hemionus | 19A,457A |
| | Odocoileus virginianus | 37A |
| | Ovis canadensis | 6N,19A,401A |
| | White-tailed Deer | 178 |
| 1080 <i>Dermacentor halli</i> | Texas Peccary | 311 |
| 1081 <i>Dermacentor hunteri</i> | Odocoileus hemionus | 19A |
| | Ovis canadensis | 19A,401A |
| 1082 <i>Dermacentor nigrolineatus</i> | Horse | 353 |
| | Odocoileus virginianus | 264 |
| 1083 <i>Dermacentor parumapertus</i> | Cottontail Rabbit | 178 |
| | Cynomys ludovicianus | 174,178 |
| | Dipodomys ordi | 174,178 |
| | Dipodomys spectabilis | 174 |
| | Jack Rabbit | 178 |
| | Lepus californicus | 173,174, 400N |
| | Neotoma micropus | 174 |
| | Onychomys leucogaster | 174 |
| | Packrat | 178 |
| | Peromyscus | 178 |
| | Rattus norvegicus | 178 |
| | Spermophilus spilosoma | 174 |
| | Spermophilus tereticaudus | 279A |
| | Spermophilus tridecemlineatus | 174,178 |
| | Spermophilus variegatus | 165 |
| | Sylvilagus auduboni | 174,400N |
| 1084 <i>Dermacentor variabilis</i> | Bassariscus astutus | 120 |
| | Canis familiaris | 220,385 |
| | Canis latrans | 158 |
| | Canis rufus | 178 |
| | Cotton Rat | 178 |
| | Cow | 178 |
| | Coyote | 55 |

| | | | |
|------|---|----------------------|-----|
| | Dog | 178 | |
| | Felis rufus | 178,439 | |
| | Goat | 178 | |
| | Gray Fox | 178 | |
| | House Cat | 178 | |
| | Javalina | 178 | |
| | Man | 178 | |
| | Opossum | 178 | |
| | Packrat | 178 | |
| | Peromyscus gossypinus | 178 | |
| | Raccoon | 178 | |
| | Rattus rattus | 178 | |
| | Sus scrofa | 113 | |
| | Taxidea taxus | 178 | |
| | Tayassu tajacu | 168,409 | |
| | Urocyon cinereoargenteus | 385 | |
| | Wolf | 158 | |
| | Odocoileus virginianus | 264 | |
| 1085 | <i>Haemaphysalis chordeilis</i> | Meadow Lark | 46 |
| 1086 | <i>Haemaphysalis leporis-</i> <i>palustris</i> | Amphispiza bilineata | 178 |
| | Bassariscus astutus | 120 | |
| | Bobwhite Quail | 55 | |
| | Cardinal | 55 | |
| | Centurus aurifrons | 178 | |
| | Colinus virginianus | 355 | |
| | Cottontail Rabbit | 55,178 | |
| | Deer | 55 | |
| | Dendrocopus scalaris | 178 | |
| | Dipodomys ordi | 174,178 | |
| | Felis rufus | 439 | |
| | Goat | 178 | |
| | Hylocichla gutta | 178 | |
| | Jack Rabbit | 55,178 | |
| | Lanius ludovicianus | 178 | |
| | Lepus californicus | 165,220,400N | |
| | Meadow Lark | 55 | |
| | Mimus polyglottis | 178 | |
| | Molothrus ater | 178 | |
| | Neotoma floridana | 385 | |
| | Neotoma micropus | 174 | |
| | Odocoileus virginianus | 264 | |
| | Oreoscoptes montanus | 178 | |
| | Poecetes gramineus | 178 | |
| | Richmondia cociferus | 178 | |
| | Roadrunner | 178 | |
| | Sialia currucoides | 178 | |
| | Spizella arborea | 178 | |
| | Sturnella neglecta | 178 | |
| | Sturnus vulgaris | 43 | |
| | Sylvilagus auduboni | 174,400N | |
| | Sylvilagus floridanus | 220,385 | |
| | Tayassu tajacu | 409 | |
| | Thryomanes bewickii | 178 | |
| | Toxostoma curvirostre | 178 | |
| | Zonotrichia leucophrys | 178 | |

| | | | |
|------|--------------------------|----------------------------|---------|
| 1087 | <i>Ixodes</i> | Baiomys taylori | 316,385 |
| | | Fox | 178 |
| | | Mephitis mephitis | 220 |
| | | Odocoileus virginianus | 407 |
| | | Perognathus | 385 |
| | | Peromyscus leucopus | 385 |
| | | Rattus rattus | 385 |
| | | Sigmodon hispidus | 316 |
| | | Urocyon cinereoargenteus | 166 |
| | | Wren | 55 |
| 1088 | <i>Ixodes angustus</i> | Neotoma micropus | 46 |
| 1089 | <i>Ixodes banksi</i> | Procyon lotor | 385 |
| 1090 | <i>Ixodes brunneus</i> | Sturnus vulgaris | 43 |
| 1091 | <i>Ixodes cadsi</i> | Liomys irroratus | 180 |
| 1092 | <i>Ixodes conepati</i> | Bassariscus astutus | 120 |
| | | Bat | 271 |
| | | Conepatus | 178 |
| | | Mephitis mephitis | 178,470 |
| 1093 | <i>Ixodes cookei</i> | Bassariscus astutus | 178 |
| | | Canis latrans | 158 |
| | | Coyote | 178 |
| | | Fox | 178 |
| | | Man | 178 |
| | | Procyon lotor | 165 |
| | | Raccoon | 178 |
| | | Skunk | 178 |
| | | Spermophilus variegatus | 165,178 |
| | | Urocyon cinereoargenteus | 166 |
| | | Wolf | 178 |
| 1094 | <i>Ixodes eadsi</i> | Liomys irroratus | 263 |
| | | Mus musculus | 263 |
| | | Peromyscus leucopus | 263 |
| | | Reithrodontomys fulvescens | 263 |
| | | Sigmodon hispidus | 263 |
| 1095 | <i>Ixodes hearlei</i> | Procyon lotor | 385 |
| 1096 | <i>Ixodes kingi</i> | Badger | 46,178 |
| | | Bassariscus astutus | 470 |
| | | Coyote | 178 |
| | | Cynomys ludovicianus | 174 |
| | | Prairie dog | 46,178 |
| | | Ringtail | 178 |
| | | Skunk | 46 |
| | | Wolf | 46 |
| 1097 | <i>Ixodes scapularis</i> | Bobcat | 178 |
| | | Canis rufus | 385 |
| | | Cattle | 152 |
| | | Coyote | 55 |
| | | Deer | 55,178 |
| | | Didelphis virginiana | 385 |
| | | Dog | 178 |
| | | Fox | 178 |
| | | Man | 55,178 |
| | | Odocoileus virginianus | 385,407 |
| | | Ophisaurus ventralis | 178 |
| | | Opossum | 178 |
| | | Procyon lotor | 385 |

| | | |
|--------------------------------------|-----------------------------------|-------------|
| | Raccoon | 178 |
| | Sceloporus olivaceus | 55 |
| | Sceloporus undulatus hyacinthinus | 55 |
| | Scincella lateralis | 55 |
| | Sciurus carolinensis | 385 |
| | Sciurus niger | 178,385 |
| | Sus scrofa | 113 |
| | Tayassu tajacu | 168 |
| | Urocyon cinereoargenteus | 385 |
| | Wolf | 178 |
| 1098 <i>Ixodes sculptus</i> | Spermophilus spilosoma | 174,178 |
| | Spermophilus variegatus | 46 |
| 1099 <i>Ixodes texanus</i> | Bassariscus astutus | 470 |
| | Fox | 178 |
| | Raccoon | 55,178 |
| | Urocyon cinereoargenteus | 166 |
| 1100 <i>Ixodes tovari</i> | Amphispiza bilineata | 175 |
| | Black-throated Sparrow | 78 |
| | Dipodomys ordi | 78 |
| | Goat | 78 |
| | Lepus californicus | 175,178 |
| | Meadow Lark | 178 |
| | Sturnella neglecta | 175 |
| | Sylvilagus auduboni | 175,178 |
| 1101 <i>Ixodes woodi</i> | Lepus californicus | 174 |
| | Neotoma | 272 |
| | Neotoma micropus | 174,178,229 |
| | Notiosorex crawfordi | 178 |
| | Onychomys leucogaster | 174 |
| | Opossum | 178 |
| | Packrat Den | 178 |
| | Peromyscus leucopus | 178 |
| | Peromyscus maniculatus | 178 |
| | Prairie Dog Den | 178 |
| | Sigmodon hispidus | 178 |
| | Skunk | 178 |
| | Speotyto cunicularia | 174 |
| | Spermophilus spilosoma | 174 |
| | Spermophilus tridecemlineatus | 174 |
| | Sylvilagus auduboni | 174 |
| | Tyrannus verticalis | 178 |
| 1102 <i>Ornithodoros</i> | Antrozous pallidus | 54A |
| | Cliff Swallow | 55 |
| | Eptesicus fuscus | 54A |
| | Leptonycteris sanborni | 54A |
| | Macrotus californicus | 54A |
| | Myotis leibii | 54A |
| | Myotis velifer | 254 |
| | Pipistrellus hesperus | 54A,165 |
| | Plecotus townsendi | 54A |
| | Sigmodon hispidus | 316 |
| | Tadarida brasiliensis | 385 |
| 1103 <i>Ornithodoros concanensis</i> | Bat | 178,271 |
| | Eptesicus fuscus | 254 |
| | Petrochelidon fulva pallida | 177 |

| | | |
|--------------------------------------|---------------------------|----------------------------|
| | Plecotus townsendi | 254 |
| 1104 <i>Ornithodoros kelleyi</i> | Antrozous pallidus | 54A |
| | Plecotus townsendi | 54A |
| 1105 <i>Ornithodoros megnini</i> | Antilocapra americana | 108N,411N |
| 1106 <i>Ornithodoros stageri</i> | Bat | 271 |
| | Free-tailed Bat | 178 |
| 1107 <i>Ornithodoros talaje</i> | Cliff Swallow | 55 |
| | Dipodomys | 269A |
| | Dipodomys ordi | 178 |
| | Liomys irroratus | 180 |
| | Neotoma micropus | 178 |
| | Perognathus hispidus | 178 |
| | Sigmodon hispidus | 172,178 |
| | Spermophilus mexicanus | 178 |
| | Spermophilus tereticaudus | 279 |
| 1108 <i>Ornithodoros turicata</i> | Armadillo | 178 |
| | Cynomys ludovicinaus | 174,372N |
| 1109 <i>Ornithodors yumatensis</i> | Bat | 178,271 |
| | Myotis thysanodes | 54A |
| | Myotis velifer | 54A,112,112A,170A,178,402, |
| | | 474 |
| | Myotis yumanensis | 54A |
| | Plecotus townsendi | 54A |
| | Tadarida brasiliensis | 402 |
| 1110 <i>Otobius megnini</i> | Antilocapra americana | 317N |
| | Bos taurus | 220 |
| | Canis latrans | 158 |
| | Cattle | 152,178,354 |
| | Deer | 55 |
| | Dog | 178 |
| | Man | 178 |
| | Odocoileus heminous | 178 |
| | Ovis canadensis | 6N |
| | Sheep | 178 |
| | Sylvilagus auduboni | 400N |
| 1111 <i>Pyrosoma bigeminum</i> | Cow | 431 |
| 1112 <i>Rhipicephalus sanguineus</i> | Canis familiaris | 385 |
| | Cow | 178 |
| | Dog | 178 |
| | Goat | 178 |
| | Man | 178 |
| | Mephitis mephitis | 456 |
| | Mule | 178 |
| | Odocoileus virginianus | 264 |
| | Raccoon | 178 |
| 1113 <i>Tick</i> | Bos taurus | 508 |
| 1114 <i>Winter Tick</i> | Deer | 278N |

Arthropoda-Crustacea-Copepoda

| | | |
|---------------------------------|-----------------------|-----|
| 1115 <i>Actheres micropteri</i> | Ictalurus punctatus | 211 |
| | Micropterus salmoides | 211 |
| 1116 <i>Argulus flavescens</i> | Aplodinotus grunniens | 211 |
| | Ictalurus punctatus | 211 |

| | | |
|---------------------------------|-------------------------|-----|
| | Lepomis macrochirus | 211 |
| | Micropterus salmoides | 211 |
| 1117 <i>Argulus longicaudus</i> | Pomoxis annularis | 211 |
| 1118 <i>Ergasilus arthrosis</i> | Aplodinotus grunniens | 211 |
| | Ictalurus punctatus | 211 |
| | Lepomis gulosus | 211 |
| | Lepomis macrochirus | 211 |
| | Lepomis megalotis | 211 |
| | Lepomis microlophus | 211 |
| | Micropterus punctulatus | 211 |
| | Micropterus salmoides | 211 |
| 1119 <i>Ergasilus tenax</i> | Pomoxis annularis | 211 |
| 1120 <i>Lernaea cyprinacea</i> | Ictalurus punctatus | 211 |
| | Lepomis macrochirus | 211 |
| | Pomoxis annularis | 211 |
| | Rana catesbeiana | 36 |
| 1121 <i>Lernaea tenuis</i> | Aplodinotus grunniens | 211 |

Arthropoda-Insecta-Dipteria

| | | |
|-------------------------------------|------------------------|------|
| 1122 <i>Basilisa antrozoi</i> | Antrozous pallidus | 54A |
| 1123 <i>Basilisa corynorhini</i> | Plecotus townsendi | 54A |
| 1124 <i>Basilisa forcipata</i> | Myotis californicus | 54A |
| | Myotis thysanodes | 54A |
| 1125 <i>Botfly larva</i> | Deer | 278N |
| 1126 <i>Callitroga hominivorax</i> | Odocoileus virginianus | 208 |
| 1127 <i>Cattle grub</i> | Cattle | 150 |
| 1128 <i>Cephenemyia</i> | Odocoileus hemionus | 457A |
| | Odocoileus virginianus | 208 |
| 1129 <i>Cochliomyia americana</i> | Castor canadensis | 110 |
| | Didelphis virginiana | 288 |
| | Lepus californicus | 288 |
| | Odocoileus virginianus | 288 |
| | Sylvilagus floridanus | 288 |
| 1130 <i>Cochliomyia macellaria</i> | Lepus californicus | 397 |
| | Sylvilagus floridanus | 288 |
| 1131 <i>Cuterebra</i> | Sylvilagus auduboni | 400N |
| 1132 <i>Gasterophilus</i> | Horse | 424 |
| 1133 <i>Hornfly</i> | Bos taurus | 508 |
| 1134 <i>Hypoderma lineatum</i> | Cattle | 148 |
| 1135 <i>Lipoptena</i> | Odocoileus virginianus | 183 |
| 1136 <i>Lipoptena mazamae</i> | Cattle | 151 |
| | Odocoileus virginianus | 129 |
| | White-tailed Deer | 410 |
| 1137 <i>Lucilia</i> | Lepus californicus | 288 |
| 1138 <i>Melophagus ovinus</i> | Odocoileus virginianus | 385 |
| | Ovis aries | 220 |
| 1139 <i>Microlychnia pusilla</i> | Colinus virginianus | 355 |
| | Zenaida macroura | 115 |
| 1140 <i>Nycteribiidae</i> | Myotis velifer | 254 |
| 1141 <i>Nycterophilia coxata</i> | Macrotus californicus | 54A |
| 1142 <i>Olfersia bisulcata</i> | Cathartes aura | 474 |
| 1143 <i>Ornithomyia fringillina</i> | Sturnus vulgaris | 43 |

| | | | |
|------|---------------------------------|-------------------------|-------------|
| 1144 | <i>Paratrichobius longicrus</i> | Choeronycteris mexicana | 54A |
| 1145 | <i>Protocalliphora</i> | Sturnus vulgaris | 43 |
| 1146 | <i>Sarcophaga plinthopyga</i> | Lepus californicus | 397 |
| 1147 | <i>Sarcophaga sulcata</i> | Lepus californicus | 397 |
| 1148 | <i>Screw worm</i> | Bos taurus | 508 |
| 1149 | <i>Synthesiomyia</i> | Lepus californicus | 288 |
| 1150 | <i>Trichobius adamsi</i> | Macrotus californicus | 54A |
| 1151 | <i>Trichobius corynorhini</i> | Plecotus townsendi | 54A |
| 1152 | <i>Trichobius major</i> | Bat | 271 |
| | | Myotis velifer | 220,253,402 |
| | | Plecotus townsendi | 254A |
| | | Tadarida brasiliensis | 402 |
| 1153 | <i>Trichobius mixtus</i> | Choeronycteris mexicana | 54A |
| 1154 | <i>Trichobius sphaeronotus</i> | Bat | 271 |
| | | Leptonycteris sanborni | 54A |

Arthropoda-Insecta-Hemiptera

| | | | |
|------|-----------------------------|-----------------------|---------|
| 1155 | <i>Cimex pilosellus</i> | Antrozous pallidus | 54A |
| | | Eptesicus fuscus | 54A,204 |
| | | Myotis thysanodes | 54A |
| | | Myotis velifer | 54A |
| | | Myotis yumanensis | 54A |
| | | Pipistrellus hesperus | 54A |
| 1156 | <i>Melanolestes picipes</i> | Man | 163 |
| 1157 | <i>Primicimex cavernis</i> | Bat | 271 |

Arthropoda-Insecta-Mallophaga and Anoplura

| | | | |
|------|---------------------------------|----------------------------------|-----|
| 1158 | <i>Acanthophthirius longa</i> | Tadarida brasiliensis | 254 |
| 1159 | <i>Acidoproctus</i> | Fulvous Tree Duck | 504 |
| 1160 | <i>Actornithophilus</i> | Totanus melanoleucus | 504 |
| 1161 | <i>Actornithophilus funebre</i> | Gelochelidon nilotica | 218 |
| | | Larus argentatus | 218 |
| | | Larus atricilla | 218 |
| | | Larus delawarensis | 218 |
| | | Sterna hirundo | 218 |
| | | Sterna maxima | 218 |
| 1162 | <i>Actornithophilus lari</i> | Larus argentatus smithsonianus | 504 |
| 1163 | <i>Anaticola anseris</i> | Domestic Goose | 504 |
| 1164 | <i>Anaticola crassicornis</i> | Anas clypeata | 58 |
| | | Anas creca | 74 |
| | | Anas cyanoptera septentrionalium | 499 |
| | | Anas discors | 504 |
| | | Aythya collaris | 504 |
| | | Melanitta doglandi | 504 |
| 1165 | <i>Anaticola dafilensis</i> | Dafila acuta | 504 |
| 1166 | <i>Anatoceous dentatus</i> | Anas clypeata | 58 |
| | | Anas platyrhynchos platyrhynchos | 504 |
| | | Aythya collaris | 504 |

| | | |
|--|----------------------------------|----------|
| | Branta canadensis leucopareia | 504 |
| 1167 <i>Anatoceous ictoerodes</i> | Anas clypeata | 58 |
| | Anas creca | 74 |
| | Anas discors | 504 |
| 1168 <i>Aquanirmus americanus</i> | Podiceps caspicus californicus | 504 |
| 1169 <i>Ardeicola</i> | Hydranassa tricolor ruficollis | 504 |
| 1170 <i>Ardeicola ajajae</i> | Ajaia ajaja | 473 |
| 1171 <i>Ardeicola botauri</i> | Botaurus lentiginosus | 473 |
| 1172 <i>Ardeicola celeris</i> | Ixobrychus exilis | 473 |
| 1173 <i>Ardeicola cruscula</i> | Ardea herodias | 473 |
| 1174 <i>Ardeicola elongata</i> | Eudocimus albus | 473 |
| 1175 <i>Ardeicola florida</i> | Florida caerulea | 473 |
| 1176 <i>Ardeicola florida nigra</i> | Butorides virescens | 473 |
| | Dichromanassa rufescens | 473 |
| | Hydranassa tricolor | 473 |
| 1177 <i>Ardeicola gaibagla</i> | Bubulcus ibis | 473 |
| | Casmerodius albus | 473 |
| | Florida thula | 473 |
| 1178 <i>Ardeicola goisagi</i> | Nycticorax nycticorax | 473 |
| 1179 <i>Ardeicola loculator</i> | Mycteria americana | 473 |
| 1180 <i>Ardeicola nyctanassa</i> | Nyctanassa violacea | 473 |
| 1181 <i>Ardeicola raphidius</i> | Plegadis chihi | 473 |
| | Plegadis falcinellus | 473 |
| | Plegadis falcinellus falcinellus | 504 |
| 1182 <i>Ardeicola robusta</i> | Eudocimus albus | 473 |
| 1183 <i>Ardeiphilus floridae</i> | Florida thula | 473, 474 |
| 1184 <i>Austromenopon</i> | Totanus melanoleucus | 504 |
| 1185 <i>Austromenopon crocatum</i> | Numenius americanus | 70N |
| 1186 <i>Austromenopon durisetosum</i> | Capella gallinago | 287 |
| 1187 <i>Austromenopon infrequens</i> | Larus argentatus | 218 |
| | Larus atricilla | 218 |
| | Larus delawarensis | 218 |
| 1188 <i>Austromenopon sachtlebeni</i> | Catoptrophorus semipalmatus | 504 |
| 1189 <i>Austromenopon sternophilum</i> | Sterna antillarum | 218 |
| 1190 <i>Austromenopon transversum</i> | Larus delawarensis | 504 |
| 1191 <i>Bruelia augustifrons</i> | Chondestes grammacus | 504 |
| 1192 <i>Bruelia domestica</i> | Hirundo rustica erythrogaster | 504 |
| 1193 <i>Bruelia illustris</i> | Colinus virginianus | 355 |
| 1194 <i>Bruelia longa</i> | Petrochelidon pyrrhonota | 504 |
| 1195 <i>Bruelia nebulosa</i> | Sturnus vulgaris vulgaris | 504 |
| 1196 <i>Bruelia ornatissima</i> | Agelaius phoeniceus | 504 |
| 1197 <i>Bruelia peninsularis</i> | Phainopepla nitens nitens | 504 |
| 1198 <i>Bruelia picturata</i> | Zonotrichia leucophrys gambelli | 504 |
| 1199 <i>Bruelia quiscalis</i> | Cassidix mexicanus prosopidicola | 504 |
| 1200 <i>Bruelia sialii</i> | Sialia sialis sialis | 504 |
| 1201 <i>Bruelia subtilis</i> | Passer domesticus domesticus | 504 |
| 1202 <i>Bruelia vulgata</i> | Mimus polyglottos leucopterus | 504 |
| | Richmondia cardinalis cardinalis | 504 |
| 1203 <i>Campanulotes bidentatus compar</i> | Domestic Pigeon | 504 |
| 1204 <i>Carduiceps complexivus</i> | Crocethia alba | 504 |
| 1205 <i>Carduiceps pusillus</i> | Tringa solitaria solitaria | 504 |
| 1206 <i>Carduiceps zonarius</i> | Calidris canutus rufa | 504 |
| 1207 <i>Chelopistes meleagridis</i> | Meleagris gallopavo intermedia | 504 |
| 1208 <i>Ciconiphilus</i> | Hydranassa tricolor ruficollis | 504 |
| 1209 <i>Ciconiphilus</i> | Ardea herodias | 473 |

| | | | |
|------|--|---|------|
| | <i>decimfascatus</i> | <i>Ardea occidentalis</i> | 473 |
| | | <i>Botaurus lentiginosus</i> | 473 |
| | | <i>Bubulcus ibis</i> | 473 |
| | | <i>Butorides virescens</i> | 473 |
| | | <i>Casmerodius albus</i> | 473 |
| | | <i>Dichromanassa rufescens</i> | 473 |
| | | <i>Florida caerulea</i> | 473 |
| | | <i>Florida thula</i> | 473 |
| | | <i>Hydranassa tricolor</i> | 473 |
| | | <i>Ixobrychus exilis</i> | 473 |
| | | <i>Mycteria americana</i> | 473 |
| | | <i>Nyctanassa violacea</i> | 473 |
| | | <i>Nycticorax nycticorax</i> | 473 |
| 1210 | <i>Ciconiphilus maculipes</i> | <i>Mycteria americana</i> | 473 |
| 1211 | <i>Colinicola numidiana</i> | <i>Colinus virginianus</i> | 355 |
| 1212 | <i>Colpocephalum ajajae</i> | <i>Ajaia ajaja</i> | 473 |
| 1213 | <i>Colpocephalum fusconigrum</i> | <i>Eudocimus albus</i> | 473 |
| 1214 | <i>Colpocephalum kelloggi</i> | <i>Cathartes aura</i> | 504 |
| 1215 | <i>Colpocephalum leptopygos</i> | <i>Plegadis chihi</i> | 473 |
| 1216 | <i>Colpocephalum polybori</i> | <i>Polyborus cheriway auduboni</i> | 504 |
| 1217 | <i>Colpocephalum scalariforme</i> | <i>Mycteria americana</i> | 473 |
| 1218 | <i>Columbicola</i> | <i>Zenaida asiatica asiatica</i> | 504 |
| 1219 | <i>Columbicola baculoides</i> | <i>Zenaida macroura marginella</i> | 504 |
| 1220 | <i>Columbicola columbae</i> | <i>Columba livia</i> | 504 |
| | | Pigeon | 299 |
| 1221 | <i>Columbicola macrourae</i> | <i>Zenaida macroura marginella</i> | 504 |
| 1222 | <i>Columbicola passerinae pallescens</i> | <i>Columbigallina passerina</i> | 504 |
| 1223 | <i>Comatomenopon dichromanassae</i> | <i>Dichromanassa rufescens</i> | 473 |
| 1224 | <i>Comatomenopon ibis</i> | <i>Bubulcus ibis</i> | 473 |
| 1225 | <i>Comatomenopon thulae</i> | <i>Florida thula</i> | 473 |
| 1226 | <i>Craspedorrhynchus dilatatus</i> | <i>Buteo swainsoni</i> | 504 |
| 1227 | <i>Craspedorrhynchus haliyeti</i> | <i>Haliaeetus leucocephalus leucocephalus</i> | 504 |
| 1228 | <i>Cuculicola splendidus</i> | <i>Geococcyx californianus</i> | 504 |
| 1229 | <i>Cuculiphilus coragypsis</i> | <i>Coragyps atratus</i> | 504 |
| 1230 | <i>Cummingsiella inexpectata</i> | <i>Recurvirostra americana</i> | 504 |
| 1231 | <i>Cummingsiella tongirostricola</i> | <i>Numenius americanus</i> | 504 |
| 1232 | <i>Damalinia limbata</i> | Angora Goat | 504 |
| 1233 | <i>Damalinia lipeuroides</i> | <i>Odocoileus hemionus</i> | 504 |
| | | White-tailed Deer | 457A |
| 1234 | <i>Degeeriella carruthi</i> | <i>Falco sparverius sparverius</i> | 504 |
| 1235 | <i>Degeeriella discocephalus</i> | <i>Aguila chrysaetos canadensis</i> | 504 |
| 1236 | <i>Degeeriella giebeli</i> | <i>Buteo swainsoni</i> | 504 |
| | | <i>Circus cyaneus hudsonius</i> | 504 |
| | | <i>Falco sparverius sparverius</i> | 504 |
| 1237 | <i>Degeeriella illustris</i> | <i>Sturnus vulgaris</i> | 43 |
| 1238 | <i>Degeeriella nebulosa</i> | <i>Sturnus vulgaris</i> | 43 |
| 1239 | <i>Dennysus bruneri</i> | <i>Aeronautes saxatalis</i> | 504 |
| | | <i>Chaetura pelagica</i> | 504 |
| 1240 | <i>Enderleinellus longiceps</i> | Merriam's Kangaroo Rat | 42A |
| 1241 | <i>Enderleinellus osborni</i> | <i>Spermophilus tereticaudus</i> | 279A |
| 1242 | <i>Enderleinellus suturalis</i> | Harris' Ground Squirrel | 42A |

| | | | |
|------|---|----------------------------|----------|
| | Spermophilus mexicanus | 322 | |
| | Spermophilus pilosoma | 322, 336N | |
| | Spermophilus tereticaudus | 336N | |
| 1243 | <i>Epidermoptes</i> | Anas creca | 74 |
| 1244 | <i>Eucolpocephalum femorale</i> | Ajaia ajaja | 473 |
| 1245 | <i>Fahrenholzia boleni</i> | Perognathus flavus | 307 |
| 1246 | <i>Fahrenholzia ehrlichi</i> | Liomys irroratus | 180, 255 |
| 1247 | <i>Fahrenholzia microcephala</i> | Liomys irroratus | 322 |
| 1248 | <i>Fahrenholzia pinnata</i> | Dipodomys merriami | 336N |
| | Dipodomys ordi | 322, 336N | |
| | Dipodomys spectabilis | 336N | |
| | Merriam's Kangaroo Rat | 42A | |
| | Onychomys leucogaster | 322 | |
| | Perognathus flavus | 336N | |
| 1249 | <i>Fahrenholzia texana</i> | Liomys irroratus | 180, 255 |
| 1250 | <i>Fahrenholzia zacatacae</i> | Perognathus hispidus | 322 |
| 1251 | <i>Falcolipeurus marginalis</i> | Cathartes aura | 504 |
| 1252 | <i>Felicola subrostratus</i> | Domestic Cat | 504 |
| 1253 | <i>Fulicoffula longipila</i> | Fulica americana americana | 504 |
| 1254 | <i>Geomydoecus geomydis</i> | Civet Cat | 504 |
| | Cratogeomys castanops | 165, 504 | |
| | Geomys bursarius | 385, 504 | |
| | Gopher | 504 | |
| | Peromyscus | 504 | |
| 1255 | <i>Geomydoecus minor</i> | Thomomys bottae | 504N |
| 1256 | <i>Geomydoecus truncatus</i> | Geomys personatus | 504 |
| 1257 | <i>Gliricola porcelli</i> | Guinea Pig | 504 |
| 1258 | <i>Goniocotes gallinae</i> | Chicken | 504 |
| 1259 | <i>Goniodes gigas</i> | Chicken | 504 |
| 1260 | <i>Gonoides ortygis</i> | Colinus virginianus | 355, 503 |
| 1261 | <i>Gyropus ovalis</i> | Guinea Pig | 504 |
| 1262 | <i>Haematopinus</i> | Mephitis mephitis | 220 |
| 1263 | <i>Haematopinus eurysternus</i> | Cow | 322 |
| 1264 | <i>Haematopinus suis</i> | Sus scrofa | 220 |
| 1265 | <i>Haematopinus suis</i> <i>adventicus</i> | Dog | 322 |
| 1266 | <i>Haemotosiphon inodorus</i> | Cathartes aura | 474 |
| 1267 | <i>Heterodoxus spiniger</i> | Canis latrans | 158, 504 |
| | Domestic Dog | 504 | |
| | Wolf | 158 | |
| 1268 | <i>Holomenopon clypeilargum</i> | Anas clypeata | 58 |
| 1269 | <i>Holomenopon setigerum</i> | Anas clypeata | 58 |
| | Anas creca | 74 | |
| 1270 | <i>Holomenopon toomisi</i> | Aix sponsa | 504 |
| 1271 | <i>Hoplopleura</i> | Perognathus hispidus | 316 |
| 1272 | <i>Hoplopleura acanthropus</i> | Microtus longicaudus | 336N |
| 1273 | <i>Hoplopleura arboricola</i> | Cliff Chipmunk | 42A |
| | Tamias minimus | 336N | |
| | Tamias quadrivittatus | 336N | |
| 1274 | <i>Hoplopleura ferrisi</i> | Brush Mouse | 42A |
| | Rock Mouse | 42A | |
| 1275 | <i>Hoplopleura hesperomydis</i> | Microtus longicaudus | 336N |
| | Onychomys leucogaster | 322, 336N | |
| | Peromyscus | 322 | |
| | Peromyscus boylii | 336N | |
| | Peromyscus difficilis | 336N | |

| | | |
|--|----------------------------------|---------------------|
| | Peromyscus leucopus | 336N |
| | Peromyscus maniculatus | 336N |
| | Peromyscus truei | 336N |
| 1276 <i>Hoplopleura hirsuta</i> | Dicotyles tajacu | 322 |
| | Neotoma floridana | 385 |
| | Perognathus hispidus | 316 |
| | Sigmodon hispidus | 250,316,322, 385 |
| | Sylvilagus auduboni | 322 |
| 1277 <i>Hoplopleura oenomydis</i> | Rattus norvegicus | 322,376,382, 448 |
| | Rattus | 322 |
| | Rattus rattus | 376 |
| 1278 <i>Hoplopleura onchomydis</i> | Grasshopper Mouse | 42A |
| 1279 <i>Hoplopleura sciuricola</i> | Sciurus carolinensis | 385 |
| | Sciurus niger | 322,385 |
| 1280 <i>Ibidoecus bimaculatus</i> | Eudocimus albus | 473 |
| 1281 <i>Ibidoecus bisignatus</i> | Plegadis chihi | 473 |
| | Plegadis falcinellus | 473 |
| 1282 <i>Ibidoecus iberoamericanus</i> | Ajaia ajaja | 473 |
| 1283 <i>Ineidifrons transpositus</i> | Fulica americana americana | 504 |
| 1284 <i>Kurodaia painei</i> | Otus asio | 504 |
| 1285 <i>Kurodaia pectinata</i> | Owl | 504 |
| 1286 <i>Kurodaia subpachygaster</i> | Tyto alba pratincola | 504 |
| 1287 <i>Laemobothrion atrum</i> | Fulica americana americana | 504 |
| 1288 <i>Laemobothrion buteonivorum</i> | Chicken Hawk | 504 |
| 1289 <i>Laemobothrion mendesi</i> | Aguila chrysaetos canadensis | 504 |
| 1290 <i>Lagopoecus numidianus</i> | Colinus virginianus | 504 |
| 1291 <i>Linognathus africanus</i> | Goats | 322 |
| 1292 <i>Linognathus setosus</i> | Mus musculus | 321,322 |
| 1293 <i>Louse</i> | Bos taurus | 508 |
| 1294 <i>Lunaceps acrophilus</i> | Crocethia alba | 504 |
| 1295 <i>Lunaceps drusti</i> | Calidris canutus rufa | 504 |
| 1296 <i>Lunaceps numenii numenii</i> | Numenius americanus | 70N |
| 1297 <i>Lunaceps pusillus</i> | Tringa solitaria solitarius | 504 |
| 1298 <i>Machaerilaemus melospizae</i> | Melospiza melodia | 504 |
| 1299 <i>Menacanthus</i> | Colinus virginianus | 376 |
| | Colinus virginianus texanus | 504 |
| 1300 <i>Menacanthus spinosum</i> | Sturnus vulgaris | 43 |
| 1301 <i>Menopon gallinae</i> | Domestic Chicken | 504 |
| 1302 <i>Myrsidea cucullaris</i> | Sturnus vulgaris | 43 |
| 1303 <i>Myrsidea incerta</i> | Amphispiza bilineata | 504 |
| | Dendroica dominica | 504 |
| | Mimus polyglottos leucopterus | 504 |
| | Toxostoma curvirostre | 504 |
| | Tryannus melancholicus couchii | 504 |
| 1304 <i>Myrsidea interrupta</i> | Corvus brachyrhynchos | 504 |
| | Cassidix mexicanus prosopidicola | 504 |
| 1305 <i>Neohaematopinus</i> | Cynomys ludovicianus | 322 |
| | Spermophilus spilosoma | 322 |
| 1306 <i>Neohaematopinus citellinus</i> | Cynomys gunnisoni | 336N |
| | Harris' Ground Squirrel | 42A |
| | Spermophilus spilosoma | 336N |
| | Spermophilus tereticaudus | 188A,279A |
| | Spermophilus variegatus | 336N |

| | | | |
|------|-------------------------------------|--|---|
| 1307 | <i>Neohaematopinus laeviusculus</i> | Spermophilus variegatus | 165,322,336N |
| 1308 | <i>Neohaematopinus marmotae</i> | Cynomys ludovicianus Dipodomys ordi Marmota flaviventris Neotoma albigula Neotoma mexicana Neotoma micropus Spermophilus spilosoma | 229,322 336N 336N 336N 336N 229,336N 322 |
| 1309 | <i>Neohaematopinus neotomae</i> | Mexican Woodrat Neotoma micropus White-throated Woodrat | 42A 322 42A |
| 1310 | <i>Neohaematopinus pacificus</i> | Cliff Chipmunk Tamias quadrivittatus | 42A 336N |
| 1311 | <i>Neohaematopinus sciurinus</i> | Sciurus carolinensis Sciurus niger | 385 165,322,385 |
| 1312 | <i>Neophilopterus heteropygus</i> | Mycteria americana | 473 |
| 1313 | <i>Neotrichodectes thoracicus</i> | Bassariscus astutus | 120 |
| 1314 | <i>Ornithobius goniopleurus</i> | Branta canadensis leucopareia | 504 |
| 1315 | <i>Oxylipeurus callipeplus</i> | Callipepla squamata pallida | 504 |
| 1316 | <i>Oxylipeurus clavatus</i> | Colinus virginianus | 355,504 |
| 1317 | <i>Oxylipeurus polytrapezius</i> | Meleagris gallopavo intermedia | 504 |
| 1318 | <i>Pecaroecus javalii</i> | Tayassu tajacu | 30,165,168, 405N,409 |
| 1319 | <i>Pediculus humanus</i> | Humans | 322 |
| 1320 | <i>Penenirmus</i> | Amphispiza belli nevadensis | 504 |
| 1321 | <i>Penenirmus aurifrons</i> | Centurus aurifrons | 504 |
| 1322 | <i>Penenirmus jungens</i> | Colaptes auratus auratus | 504 |
| 1323 | <i>Philopterus</i> | Toxostoma curvirostre | 504 |
| 1324 | <i>Philopterus citrinellae</i> | Calamospiza melanocorys Chlorora chlorura Eremophila alpestris Lanius ludovicianus Myadestes townsendi townsendi Pyrrhuloxia sinuata Tangavius aeneus aeneus Tyrannus melancholicus couchii | 504 504 504 504 504 504 504 504 504 |
| 1325 | <i>Philopterus fringillae</i> | Passer domesticus domesticus | 504 |
| 1326 | <i>Philopterus mesosomaticus</i> | Hirundo rustica erythrogaster | 504 |
| 1327 | <i>Philopterus ocellatus</i> | Corvus brachyrhynchos | 504 |
| 1328 | <i>Philopterus sialii</i> | Sialia mexicana bairdi Sialia sialis sialis | 504 504 |
| 1329 | <i>Phthrius pubis</i> | Man | 322 |
| 1330 | <i>Physconelloides</i> | Columbigallina passerina pallescens | 504 |
| 1331 | <i>Physconelloides passerinae</i> | Zenaida macroura marginella | 504 |
| 1332 | <i>Physconelloides zenaidurne</i> | Columba livia Zenaida macroura marginella | 504 504 |
| 1333 | <i>Picicola foedus</i> | Muscivora forficata Sayornis saya saya Sparrow | 504 504 504 |
| 1334 | <i>Picicola marginatulus</i> | Dryocopus pileatus pileatus | 504 |
| 1335 | <i>Plegadiphilus eudocimus</i> | Eudocimus albus | 473 |
| 1336 | <i>Plegadiphilus plegadis</i> | Plegadis chihi | 473 |

| | | |
|---|----------------------------------|-------------------------|
| | Plegadis falcinellus | 473 |
| 1337 <i>Polyplax auricularis</i> | Deer Mouse | 42A |
| | Grasshopper Mouse | 42A |
| | Onychomys leucogaster | 322, 336N |
| | Peromyscus difficilis | 336N |
| | Peromyscus maniculatus | 336N |
| | Peromyscus truei | 336N |
| 1338 <i>Polyplax serrata</i> | Mus musculus | 321, 322 |
| 1339 <i>Polyplax spinulosa</i> | Baimoys taylori | 316 |
| | Mus musculus | 220, 385 |
| | Perognathus hispidus | 316 |
| | Rattus norvegicus | 322, 376, 448 |
| | Rattus rattus | 250, 322, 376, 384, 448 |
| | Sigmodon hispidus | 316 |
| 1340 <i>Polyplax spinulosus</i> | Lepus californicus | 220 |
| 1341 <i>Precaroeocus javalli</i> | Tayassu tajacu | 322 |
| 1342 <i>Pseudomenopon nilsum</i> | Porzana carolina | 504 |
| 1343 <i>Pseudomenopon pacificum</i> | Fulica americana americana | 504 |
| 1344 <i>Quadriceps boeophilus</i> | Charadrius vociferus vociferus | 504 |
| 1345 <i>Quadriceps elongatus</i> | Rynchops nigra | 218 |
| 1346 <i>Quadriceps euprepes</i> | Larus atricilla | 218 |
| 1347 <i>Quadriceps felis</i> | Larus argentatus | 218 |
| | Larus atricilla | 218 |
| | Larus delawarensis | 218 |
| 1348 <i>Quadriceps gloriosa</i> | Sterna fuscata | 218 |
| 1349 <i>Quadriceps hospes</i> | Sandpiper | 504 |
| 1350 <i>Quadriceps lineolatus</i> | Larus argentatus | 218 |
| | Sterna maxima | 218 |
| 1351 <i>Quadriceps nigrolimbatus</i> | Limnodromus griseus | 504 |
| 1352 <i>Quadriceps phaeonota</i> | Hydrachelidon nigra surinamensis | 218 |
| | Sterna hirundo | 218 |
| 1353 <i>Quadriceps praestans</i> | Gelochelidon nilotica | 218 |
| | Sterna hirundo | 218 |
| | Sterna maxima | 218 |
| 1354 <i>Quadriceps similis</i> | Totanus melanoleucus | 504 |
| 1355 <i>Rhinonyssus rhinolethrum</i> | Anas discors | 325 |
| | Aythya valisineria | 325 |
| | Chen caerulescens | 325 |
| | Chen hyperborea | 325 |
| | Chen rossi | 325 |
| | Mergus serrator | 325 |
| 1356 <i>Rhynonirmus scolopacis</i> | Capella gallinago | 287 |
| | Capella gallinago delicata | 504 |
| 1357 <i>Ricinus arcuatus</i> | Muscivora forficata | 504 |
| 1358 <i>Ricinus diffusus</i> | Geothlypis trichas | 504 |
| 1359 <i>Ricinus melospizae</i> | Melospiza melodia | 504 |
| 1360 <i>Ricinus microcephalus</i> | Spizella arborea | 504 |
| 1361 <i>Ricinus subhastatus</i> | Chlorura chlorura | 504 |
| 1362 <i>Saemundssonina</i> | Rynchops nigra | 218 |
| 1363 <i>Saemundssonina anisorhamphos</i> | Bird | 504 |
| | Black Skimmer | 504 |
| 1364 <i>Saemundssonina breviappendiculata</i> | Larus argentatus | 504 |
| 1365 <i>Saemundssonina brevicornis</i> | Sandwich Tern | 504 |

| | | | |
|------|--------------------------------------|----------------------------------|------|
| 1366 | <i>Saemundssonina gonothroax</i> | Larus argentatus | 218 |
| | | Larus atricilla | 218 |
| | | Larus delawarensis | 218 |
| 1367 | <i>Saemundssonina lari</i> | Laughing Gull | 504 |
| | | Ring-billed Gull | 504 |
| 1368 | <i>Saemundssonina laticuadata</i> | Royal Tern | 504 |
| 1369 | <i>Saemundssonina melanocephalus</i> | Gelochelidon nilotica | 218 |
| | | Rynchops nigra | 218 |
| | | Sterna antillarum | 218 |
| | | Sterna hirundo | 218 |
| | | Sterna maxima | 218 |
| 1370 | <i>Saemundssonina sterna</i> | Common Tern | 504 |
| | | Gull-billed Tern | 504 |
| 1371 | <i>Saemundssonina trinage</i> | Sanderling | 504 |
| | | Sandpiper | 504 |
| | | Semipalmated Plover | 504 |
| 1372 | <i>Solenopates binipilorus</i> | Odocoileus hemionus | 457A |
| 1373 | <i>Strigiphilus ceblebrachys</i> | Nyctea scandiaca | 504 |
| 1374 | <i>Strigiphilus cursor</i> | Bubo virginianus | 504 |
| 1375 | <i>Strigiphilus rostratus</i> | Tyto alba | 504 |
| 1376 | <i>Strigiphilus speotyti</i> | Owl | 504 |
| 1377 | <i>Strigiphilus syrnii</i> | Strix varia | 504 |
| 1378 | <i>Sturnidoecus simplex</i> | Turdus migratorius | 504 |
| 1379 | <i>Sturnidoecus sturni</i> | Sturnus vulgaris vulgaris | 504 |
| 1380 | <i>Suricatoecus octomaculatus</i> | Mephitis mephitis | 385 |
| | | Procyon lotor | 385 |
| 1381 | <i>Trichodectes canis</i> | Canis familiaris | 385 |
| | | Canis latrans | 158 |
| | | Domestic Dog | 504 |
| 1382 | <i>Trichodectes mephitis</i> | Mephitis mephitis | 504 |
| 1383 | <i>Trichodectes octomaculatus</i> | Procyon lotor | 504 |
| 1384 | <i>Trichodectes osborni</i> | Spilogale putorius | 504 |
| 1385 | <i>Trichodectes quadraticeps</i> | Urocyon cinereoargenteus | 166 |
| 1386 | <i>Trichodectes thoracicus</i> | Bassariscus astutus | 504 |
| 1387 | <i>Trinoton</i> | Gadwall | 504 |
| | | Lophodytes cucullatus | 504 |
| 1388 | <i>Trinoton lituratum</i> | Anas carolinensis | 504 |
| | | Anas discors | 504 |
| 1389 | <i>Trinoton querquedulae</i> | Anas acuta | 504 |
| | | Anas carolinensis | 504 |
| | | Anas clypeata | 58 |
| | | Anas creca | 74 |
| | | Anas cyanoptera septentrionalium | 499 |
| | | Anas platyrhynchos platyrhynchos | 504 |
| | | Aythya collaris | 504 |
| | | Aythya valisineria | 504 |
| | | Mareca americana | 504 |

Arthropoda-Insecta-Siphonaptera

| | | | |
|------|-------------------------------|------------------------------|----------|
| 1390 | <i>Actenophthalmus</i> | Grasshopper Mouse | 164 |
| | | <i>Onychomys leucogaster</i> | 160 |
| 1391 | <i>Amphalius recopinus</i> | <i>Ochotona princeps</i> | 232N |
| 1392 | <i>Anomiopsyllis</i> | <i>Neotoma micropus</i> | 159,230N |
| 1393 | <i>Anomiopsyllus hiemalis</i> | Cottontail Rabbit | 164 |
| | | <i>Cynomys ludovicianus</i> | 160 |

HOST-PARASITE BIBLIOGRAPHY

| Host Scientific Name | Parasite Number |
|---|---|
| <i>Acris crepitans</i> | 33, 35, 36, 88, 324, 405, 750, 779, 785, 786, 935 |
| <i>Acris gryllus</i> | 88, 807 |
| <i>Adinia xenica</i> | 89, 248, 249, 480, 573, 850 |
| <i>Aeronautes saxatalis</i> | 1239 |
| <i>Agelaius phoeniceus</i> | 30, 866, 1196 |
| <i>Agkistrodon contortrix</i> | 469, 651, 723 |
| <i>Aguila chrysaetos canadensis</i> | 1235, 1289 |
| <i>Aimophila ruficeps</i> | 1059 |
| <i>Aix sponsa</i> | 1270 |
| <i>Ajaia ajaja</i> | 1170, 1212, 1244, 1282 |
| <i>Amblema costata</i> | 254 |
| <i>Amblema perplicata</i> | 254 |
| <i>Amblema plicata</i> | 254, 448 |
| <i>Ambloplites rupestris</i> | 184, 305, 453, 463, 477, 522, 554, 571, 616, 767, 769, 777, 861, 873, 885 |
| <i>Ambystoma talpoideum</i> | 584 |
| <i>Ambystoma texanum</i> | 261, 584 |
| <i>Ambystoma tigrinum</i> | 14, 16, 109, 180, 705, 774 |
| <i>Ameiurus natalis</i> | 293 |
| <i>Ammocrypta vivax</i> | 552, 571 |
| <i>Ammospermophilus harrisi</i> | 1515 |
| <i>Ammotragus lervia</i> | 220, 580, 613, 641, 682, 686, 698, 745, 762, 818 |
| <i>Amnicola peracuta</i> | 244, 406 |
| <i>Amphispiza bilineata</i> | 973, 1008, 1086, 1100, 1303 |
| <i>Amphispiza belli nevadensis</i> | 1320 |
| <i>Amyda ferox</i> | 553, 619, 776 |
| <i>Anas acuta</i> | 7, 56, 493, 1016, 1389 |
| <i>Anas americana</i> | 56 |
| <i>Anas carolinensis</i> | 1016, 1388, 1389 |
| <i>Anas clypeata</i> | 43, 81, 99, 101, 102, 108, 110, 150, 206, 326, 337, 339, 410, 492, 496, 521, 524, 526, 802, 882, 1016, 1164, 1166, 1167, 1268, 1269, 1389 |
| <i>Anas creca</i> | 7, 56, 81, 108, 110, 207, 208, 326, 339, 380, 410, 493, 497, 521, 526, 527, 612, 802, 855, 883, 1164, 1167, 1243, 1269, 1389 |
| <i>Anas cyanoptera septentrionalium</i> | 81, 101, 107, 150, 206, 257, 339, 410, 493, 556, 783, 855, 858, 1164, 1389 |
| <i>Anas discors</i> | 493, 1164, 1167, 1355, 1388 |
| <i>Anas platyrhynchos</i> | 1016 |
| <i>Anas platyrhynchos platyrhynchos</i> | 7, 56, 1167, 1389 |
| <i>Anas strepera</i> | 56, 1016 |
| <i>Anaschis avara</i> | 494 |
| <i>Anatidae</i> | 1016 |
| <i>Anax</i> | 359 |
| <i>Anodonta corpulenta</i> | 254, 301, 448 |
| <i>Anolis carolinensis</i> | 160, 185 |
| <i>Antilocapra americana</i> | 153, 220, 580, 581, 582, 641, 680, 683, 745, 818, 820, 1105, 1110 |
| <i>Antilope cervicapra</i> | 95, 242, 495, 525, 640, 686, 689, 818, 820, 822, 830 |
| <i>Antrostomus carolinensis</i> | 1059 |
| <i>Antrozous pallidus</i> | 37, 1023, 1056, 1102, 1104, 1122, 1155 |
| <i>Aplodinotus grunniens</i> | 378, 552, 571, 754, 768, 1116, 1118, 1121 |
| <i>Archosargus probatocephalus</i> | 302 |
| <i>Ardea herodias</i> | 453, 1173, 1209 |
| <i>Ardea herodias herodias</i> | 115 |
| <i>Ardea occidentalis</i> | 1209 |

| | | | |
|---|---|----------------------|-----|
| <i>Arenaria interpres</i> | 1014 | <i>Asio flammeus</i> | 949 |
| <i>Astyanax mexicanus</i> | 310, 453, 463, 552, 571, 767, 769, 777, 861, 873, 885 | | |
| <i>Aythya affinis</i> | 1016 | | |
| <i>Aythya collaris</i> | 1164, 1166, 1389 | | |
| <i>Aythya valisineria</i> | 1355, 1389 | | |
| <i>Baelophus atricristatus</i> | 1059 | | |
| <i>Baiomys taylori</i> | 722, 898, 900, 922, 982, 1059, 1087, 1339, 1416, 1428, 1473, 1474, 1521 | | |
| <i>Bairdiella chrysura</i> | 486 | | |
| <i>Balistes carolinensis</i> | 300 | | |
| <i>Bassariscus astutus</i> | 142, 214, 533, 722, 735, 842, 863, 890, 906, 913, 921, 924, 937, 940, 994, 1059, 1084, 1086, 1092, 1093, 1096, 1099, 1313, 1386, 1394, 1416, 1423, 1433, 1476, 1488, 1491, 1493 | | |
| <i>Bison bison</i> | 1020 | | |
| <i>Bos taurus</i> | 1059, 1110, 1113, 1133, 1148, 1293, 1410, 1491 | | |
| <i>Botaurus lentiginosus</i> | 1171, 1209 | | |
| <i>Branta canadensis</i> | 1016 | | |
| <i>Branta canadensis leucopareia</i> | 1166, 1314 | | |
| <i>Bubo virginianus</i> | 991, 1374 | | |
| <i>Bubulcus ibis</i> | 1177, 1209, 1224 | | |
| <i>Bufo cognatus</i> | 934 | | |
| <i>Bufo debilis</i> | 934 | | |
| <i>Bufo microscaphus</i> | 934 | | |
| <i>Bufo punctatus</i> | 934 | | |
| <i>Bufo speciosus</i> | 934 | | |
| <i>Bufo terrestris</i> | 584, 700 | | |
| <i>Bufo valliceps</i> | 584, 700, 702, 750 | | |
| <i>Bufo woodhousei</i> | 934 | | |
| <i>Bufo woodhousei fowleri</i> | 750 | | |
| <i>Buteo borealis</i> | 1059 | | |
| <i>Buteo jamaicensis borealis</i> | 930 | | |
| <i>Buteo swainsoni</i> | 899, 1059, 1226, 1236 | | |
| <i>Butorides virescens</i> | 969, 1176, 1209 | | |
| <i>Calamospiza melanocorys</i> | 1008, 1324 | | |
| <i>Calidris canutus</i> | 1033 | | |
| <i>Calidris canutus rufa</i> | 1206, 1295 | | |
| <i>Callipepla squamata</i> | 200, 546, 703 | | |
| <i>Callipepla squamata pallida</i> | 1315 | | |
| <i>Callithrix jacchus</i> | 816 | | |
| <i>Camelus dromedarius</i> | 835 | | |
| <i>Canis familiaris</i> | 47, 529, 808, 810, 837, 1059, 1067, 1084, 1112, 1381, 1409, 1410, 1416, 1473, 1488, 1491 | | |
| <i>Canis latrans</i> | 26, 47, 52, 143, 145, 200, 211, 213, 215, 216, 218, 235, 236, 237, 255, 518, 529, 555, 593, 595, 605, 624, 695, 722, 729, 733, 744, 746, 771, 793, 795, 797, 808, 827, 878, 880, 1059, 1065, 1067, 1084, 1093, 1110, 1267, 1381, 1410, 1416, 1423, 1476, 1491 | | |
| <i>Canis rufus</i> | 1059, 1084, 1097, 1410, 1416, 1491, 1495, 1496 | | |
| <i>Canis</i> | 1068 | | |
| <i>Capella gallinago</i> | 57, 116, 117, 118, 119, 122, 123, 311, 339, 485, 557, 583, 801, 849, 1186, 1356 | | |
| <i>Capella gallinago delicata</i> | 1356 | | |
| <i>Capra hircus</i> | 763 | | |
| <i>Carpododes carpio</i> | 67, 112, 381, 552, 705, 876 | | |
| <i>Carpodacus mexicanus</i> | 1059 | | |
| <i>Casmerodius albus</i> | 250, 251, 1177, 1209 | | |
| <i>Cassidix mexicanus</i> | 22, 30, 266, 1034 | | |
| <i>Cassidix mexicanus prosopidicola</i> | 1199, 1304 | | |
| <i>Castor canadensis</i> | 1129 | | |
| <i>Cathartes aura</i> | 1059, 1142, 1214, 1251, 1266 | | |
| <i>Catherpes mexicanus</i> | 1059 | | |
| <i>Catoprophorus semipalmatus</i> | 1014, 1188 | | |

| | |
|--|--|
| <i>Catostomus clarki</i> | 136 |
| <i>Catostomus commersonii</i> | 466 |
| <i>Catostomus insignis</i> | 136 |
| <i>Centropomus undecimalis</i> | 471 |
| <i>Centurus aurifrons</i> | 1030, 1086, 1321 |
| <i>Cerithidea pliculosa</i> | 247, 267, 340, 402, 427 |
| <i>Cervus elaphus</i> | 51, 220, 599, 613, 699, 818, 820, 1079 |
| <i>Cervus nippon</i> | 831, 834 |
| <i>Chaenobryttus coronarius</i> | 230, 503, 510, 857 |
| <i>Chaenobryttus gulosus</i> | 228, 231, 274, 299, 416, 441, 453, 477, 503, 857 |
| <i>Chaetura pelagica</i> | 1239 |
| <i>Charadrius hiaticula</i> | 1014 |
| <i>Charadrius vociferus</i> | 1015 |
| <i>Charadrius vociferus vociferus</i> | 1344 |
| <i>Chelydra serpentina</i> | 256, 374, 449, 553, 565, 619 |
| <i>Chelydra serpentina serpentina</i> | 19, 23 |
| <i>Chen caerulescens</i> | 1355 |
| <i>Chen hyperborea</i> | 1355 |
| <i>Chen rossi</i> | 1355 |
| <i>Chlorora chlorura</i> | 1324, 1361 |
| <i>Choeronycteris mexicana</i> | 1144, 1153 |
| <i>Chonestes grammacus</i> | 672, 1008, 1059, 1191 |
| <i>Chordeiles acutipennis</i> | 973 |
| <i>Chordeiles minor minor</i> | 66, 179 |
| <i>Cichlosoma cyanoguttatum</i> | 184, 453, 552, 753, 777, 861, 873 |
| <i>Circus cyaneus hudsonius</i> | 1236 |
| <i>Citellus variegatus</i> | 1423 |
| <i>Clethrionomys gapperi</i> | 1432, 1494 |
| <i>Cnemidophorus inornatus</i> | 710 |
| <i>Cnemidophorus sacki</i> | 928 |
| <i>Cnemidophorus scalaris</i> | 710, 716 |
| <i>Cnemidophorus sexlineatus</i> | 718, 928, 1043 |
| <i>Cnemidophorus tessellatus</i> | 90, 161, 164, 167, 679, 719, 740, 1043 |
| <i>Cnemidophorus tigris</i> | 710, 718, 1043 |
| <i>Coccyzus americanus</i> | 895, 1059 |
| <i>Colaptes auratus auratus</i> | 1322 |
| <i>Coleonyx brevis</i> | 717 |
| <i>Colinus virginianus</i> | 1059, 1086, 1139, 1193, 1211, 1260, 1290, 1299, 1316, 1473 |
| <i>Colinus virginianus texanus</i> | 177, 193, 194, 195, 196, 200, 546, 593, 794, 865, 1299 |
| <i>Coluber constrictor flaviventris</i> | 652, 653, 694, 736 |
| <i>Columba fasciata</i> | 1039 |
| <i>Columba livia</i> | 6, 44, 50, 534, 554, 800, 1220, 1332 |
| <i>Columba livia domestica</i> | 972, 1021 |
| <i>Columbigallina passerina</i> | 972, 981, 1059 |
| <i>Columbigallina passerina pallescens</i> | 1222, 1330 |
| <i>Columbigallina sparverius</i> | 981 |
| <i>Conepatus mesoleucus</i> | 47, 142, 158, 622, 675, 726, 765, 848, 1059 |
| <i>Conepatus</i> | 1092 |
| <i>Cophosaurus texanus</i> | 158, 165, 167, 588, 710, 928 |
| <i>Coragyps atratus</i> | 1229 |
| <i>Corvus brachyrhynchos</i> | 61, 125, 266, 520, 523, 674, 780, 852, 866, 1304, 1327 |
| <i>Corvus cryptoleucus</i> | 61, 125, 523, 603, 866 |
| <i>Crassius auratus</i> | 315, 321, 322, 356 |
| <i>Cratogeomys castanops</i> | 932, 936, 1254, 1414 |
| <i>Crepidula plana</i> | 401 |
| <i>Crocethia alba</i> | 884, 1014, 1033, 1204, 1294 |
| <i>Crotalus atrox</i> | 919, 948 |
| <i>Crotalus viridis viridis</i> | 20 |
| <i>Crotaphytus collaris</i> | 976, 1043, 1074 |
| <i>Crotophaga sulcirostris</i> | 1028 |

| | |
|-------------------------------------|--|
| <i>Cyanocitta cristata</i> | 266, 672 |
| <i>Cyanocitta stelleri</i> | 1005 |
| <i>Cynomys gunnisoni</i> | 903, 1306, 1424, 1467, 1468 |
| <i>Cynomys ludovicianus</i> | 793, 825, 870, 932, 973, 978, 989, 990, 1047, 1083, 1096, 1108, 1305, 1308, 1393, 1416, 1423, 1467, 1476, 1491, 1493 |
| <i>Cynomys</i> | 1514 |
| <i>Cynoscion regalis</i> | 486 |
| <i>Cyprinodon variegatus</i> | 34, 84, 89, 113, 114, 249, 250, 251, 571, 573, 851 |
| <i>Cyprinus carpio</i> | 318 |
| <i>Cyrtonyx montezumae</i> | 703 |
| <i>Dafila acuta</i> | 1665 |
| <i>Dasypus novemcinctus</i> | 52, 139, 585, 941, 1059, 1065, 1495, 1496 |
| <i>Deirochelys reticularis</i> | 553, 777 |
| <i>Dendrocopos scalaris</i> | 1086 |
| <i>Dendrocygna autumnalis</i> | 59, 81, 97, 205, 246, 339, 521, 612, 712, 737, 841, 854, 856 |
| <i>Dendroica auduboni</i> | 986 |
| <i>Dendroica caerulescens</i> | 1059 |
| <i>Dendroica dominica</i> | 1303 |
| <i>Dichromanassa rufescens</i> | 1176, 1209, 1223 |
| <i>Dicotyles tajacu</i> | 538, 1276, 1492 |
| <i>Didelphis virginiana</i> | 52, 144, 342, 369, 456, 479, 544, 586, 732, 922, 958, 1059, 1097, 1129, 1410, 1416, 1473, 1488, 1491, 1493, 1496, 1526 |
| <i>Didelphis</i> | 479, 1496 |
| <i>Dionda episcopa</i> | 443, 453 |
| <i>Dipodomys agilis</i> | 1433 |
| <i>Dipodomys deserti</i> | 1445 |
| <i>Dipodomys elator</i> | 1433 |
| <i>Dipodomys merriami</i> | 10, 11, 757, 956, 1248, 1445, 1449, 1517 |
| <i>Dipodomys ordi</i> | 8, 11, 12, 828, 903, 947, 966, 968, 1055, 1083, 1086, 1100, 1107, 1248, 1308, 1419, 1423, 1440, 1444, 1445, 1446, 1447, 1453, 1454, 1476, 1491, 1511, 1512 |
| <i>Dipodomys spectabilis</i> | 18, 920, 932, 947, 1083, 1248, 1439, 1441, 1443, 1446, 1447, 1448, 1489, 1511 |
| <i>Dipodomys</i> | 755, 1107, 1514 |
| <i>Donax variabilis</i> | 271, 284, 327, 354, 388, 405, 430 |
| <i>Donax</i> | 268 |
| <i>Dorosoma cepedianum</i> | 182, 397, 859, 886 |
| <i>Dorosoma petenese</i> | 373, 550, 571, 859, 886 |
| <i>Dryocopus pileatus pileatus</i> | 1334 |
| <i>Elaphe obsoleta lindheimerii</i> | 162 |
| <i>Elaphe subocularis</i> | 894, 1071 |
| <i>Elassoma zonatum</i> | 477, 552, 643 |
| <i>Empidonax minimus</i> | 1053, 1059 |
| <i>Enallagma</i> | 359 |
| <i>Eptesicus fuscus</i> | 563, 984, 1023, 1026, 1056, 1102, 1103, 1155, 1461 |
| <i>Eremophila alpestris</i> | 1324 |
| <i>Erethizon dorsatum</i> | 47, 202, 1423, 1419 |
| <i>Erinaceus albiventris</i> | 870 |
| <i>Erolia alpina</i> | 1014 |
| <i>Erolia maritima</i> | 1014 |
| <i>Erolia melanotos</i> | 484 |
| <i>Esox americanus</i> | 571 |
| <i>Etheostoma gracile</i> | 473 |
| <i>Eudocimus albus</i> | 1174, 1182, 1213, 1280, 1335 |
| <i>Eumeces fasciatus</i> | 148, 163, 314, 584, 700 |
| <i>Falco sparverius sparverius</i> | 1234, 1236 |
| <i>Farancia abacura</i> | 159, 188, 483 |
| <i>Felis domesticus</i> | 47, 103, 210, 243, 789, 1410, 1416, 1430, 1491, 1526 |
| <i>Felis rufus</i> | 47, 143, 213, 216, 217, 237, 529, 530, 555, 590, 591, 592, 600, 669, 670, 722, 728, 746, 771, 808, 811, 839, 844, 878, 1065, 1084, 1086, 1416, 1420, 1423, |

| | |
|---|---|
| | 1429, 1491, 1493 |
| <i>Ferrissia</i> | 358 |
| <i>Florida caerulea</i> | 1175, 1209 |
| <i>Florida thula</i> | 1177, 1183, 1209, 1225 |
| <i>Fregata magnificens</i> | 348, 481 |
| <i>Fringillidae</i> | 1059 |
| <i>Fulica americana</i> | 1013 |
| <i>Fulica americana americana</i> | 1253, 1283, 1287, 1343 |
| <i>Fundulus grandis</i> | 89, 113, 181, 185, 204, 248, 249, 349, 379, 475, 573, 616, 720, 739, 748, 850, 851, 877 |
| <i>Fundulus notatus</i> | 508, 513 |
| <i>Fundulus olivaceus</i> | 453 |
| <i>Fundulus similis</i> | 34, 113, 379, 850 |
| <i>Galeichtys felis</i> | 245 |
| <i>Gambusia affinis</i> | 69, 74, 174, 185, 303, 304, 332, 355, 365, 378, 392, 422, 434, 443, 453, 476, 487, 513, 552, 571, 753, 767, 775, 814, 826, 872, 873 |
| <i>Gambusia affinis affinis</i> | 513 |
| <i>Gambusia geiseri</i> | 69, 434, 453, 552, 571, 767, 775, 861, 873 |
| <i>Gastrophyrne olivacea</i> | 584 |
| <i>Gelochelidon nilotica</i> | 1161, 1353, 1369 |
| <i>Geococcyx californianus</i> | 973, 1059, 1064, 1073, 1228 |
| <i>Geomys bursarius</i> | 122, 898, 1254, 1410, 1512 |
| <i>Geomys personatus</i> | 122, 1256 |
| <i>Geothlypis trichas</i> | 1358 |
| <i>Gopherus berlandieri</i> | 19 |
| <i>Grus canadensis</i> | 59, 421, 458, 758, 803, 840 |
| <i>Guiraca caerulea</i> | 1059 |
| <i>Haematobia irritans</i> | 781 |
| <i>Haliaeetus leucocephalus leucocephalus</i> | 1127 |
| <i>Heleodytes brunneicapillus</i> | 973 |
| <i>Hesperiphona vespertina</i> | 32, 51, 672 |
| <i>Heterodon nasicus</i> | 1043 |
| <i>Heterodon platyrhinos</i> | 383, 470, 561, 584, 651, 751 |
| <i>Hiodon alosoides</i> | 71, 309 |
| <i>Hirundo rustica erythrogaster</i> | 1192, 1326 |
| <i>Holbrookia maculata</i> | 158, 588 |
| <i>Homo sapiens</i> | 870 |
| <i>Huro salmoides</i> | 274, 299, 426, 437, 462, 499, 550, 571, 872 |
| <i>Hybognathus nuchalis</i> | 453 |
| <i>Hybomitra laticornis</i> | 613 |
| <i>Hybomitra</i> | 613 |
| <i>Hydrachelidon nigra surinamensis</i> | 1352 |
| <i>Hydranassa tricolor</i> | 1176, 1209 |
| <i>Hydranassa tricolor ruficollis</i> | 1169, 1208 |
| <i>Hyla arenicolor</i> | 934 |
| <i>Hyla cinerea</i> | 261, 399, 700 |
| <i>Hyla squirella</i> | 88, 584 |
| <i>Hylocichla gutta</i> | 1086 |
| <i>Ictalurus melas</i> | 184, 185, 238, 240, 355, 435, 453, 465 |
| <i>Ictalurus natalis</i> | 184, 186, 241, 355, 435, 453, 571, 767, 769, 777 |
| <i>Ictalurus nebulosus</i> | 82, 861 |
| <i>Ictalurus punctatus</i> | 69, 82, 83, 186, 239, 241, 289, 293, 299, 308, 436, 453, 552, 571, 594, 597, 754, 768, 1115, 1116, 1118, 1120 |
| <i>Icteria virens</i> | 1059 |
| <i>Icterus spurius</i> | 672 |
| <i>Ictiobus bubalus</i> | 72, 111, 112, 181, 190, 386, 407, 551 |
| <i>Ictiobus cyprinellus</i> | 112, 386, 407, 551 |
| <i>Ictiobus niger</i> | 112, 386, 407 |
| <i>Ixobrychus exilis</i> | 1172, 1209 |
| <i>Kinosteron subrubrum hippocrepis</i> | 19, 374, 449, 553 |

| | |
|---------------------------------------|---|
| <i>Lagodon rhomboides</i> | 464 |
| <i>Lampropeltis getulus holbrooki</i> | 383, 651 |
| <i>Lampsilis anodontoides</i> | 270, 301, 448 |
| <i>Lampsilis radiata</i> | 270, 301 |
| <i>Lanius ludovicianus</i> | 1002, 1059, 1086, 1324 |
| <i>Larus argentatus</i> | 955, 1161, 1187, 1347, 1350, 1364, 1366 |
| <i>Larus argentatus smithsonianus</i> | 1162 |
| <i>Larus atricilla</i> | 955, 1161, 1187, 1346, 1347, 1366 |
| <i>Larus delawarensis</i> | 955, 1161, 1187, 1190, 1347, 1366 |
| <i>Lasiurus cinereus</i> | 85 |
| <i>Lasiurus intermedius</i> | 980 |
| <i>Leiopisma laterale</i> | 88 |
| <i>Leiostomus xanthurus</i> | 29, 330, 455 |
| <i>Lepibema chrysops</i> | 504 |
| <i>Lepisosteus oculatus</i> | 186, 873 |
| <i>Lepisosteus osseus</i> | 186, 298, 475, 571, 575, 616, 708 |
| <i>Lepisosteus productus</i> | 186, 571, 575, 596, 661 |
| <i>Lepisosteus spatula</i> | 186, 298, 475, 575, 596, 616 |
| <i>Lepisosteus</i> | 186, 424, 475, 575 |
| <i>Lepomis auritus</i> | 68, 69, 184, 299, 305, 331, 453, 463, 552, 554, 571, 767, 768, 769, 777, 861, 873, 885 |
| <i>Lepomis cyanellus</i> | 70, 186, 227, 228, 230, 232, 233, 270, 274, 276, 285, 286, 288, 290, 294, 295, 305, 367, 368, 415, 441, 442, 453, 454, 502, 503, 505, 506, 507, 510, 512, 552, 553, 554, 571, 575, 767, 768, 769, 777, 857, 861, 872, 873 |
| <i>Lepomis gulosus</i> | 69, 184, 186, 305, 331, 364, 437, 453, 463, 477, 552, 554, 571, 767, 768, 769, 777, 861, 873, 1118 |
| <i>Lepomis humilis</i> | 70, 186, 227, 233, 270, 276, 288, 294, 442, 453, 503, 505, 506, 507, 552, 553, 575, 767, 872 |
| <i>Lepomis macrochirus</i> | 68, 70, 184, 185, 186, 187, 225, 226, 227, 228, 229, 231, 232, 270, 272, 276, 286, 288, 291, 292, 294, 299, 305, 331, 365, 367, 378, 393, 416, 441, 442, 453, 454, 463, 500, 502, 503, 505, 506, 507, 512, 514, 552, 553, 554, 571, 575, 720, 767, 768, 769, 777, 857, 861, 872, 873, 885, 1116, 1118, 1120 |
| <i>Lepomis marginatus</i> | 185, 225, 234, 286, 287, 305, 368, 378, 393, 414, 416, 441, 453, 477, 515, 873 |
| <i>Lepomis megalotis</i> | 70, 186, 187, 225, 226, 227, 230, 233, 270, 276, 286, 287, 288, 292, 305, 366, 378, 393, 413, 440, 441, 442, 453, 454, 501, 502, 503, 505, 506, 507, 510, 514, 515, 552, 553, 571, 575, 767, 775, 872, 873, 1118 |
| <i>Lepomis microlophus</i> | 184, 186, 299, 305, 364, 367, 372, 378, 453, 463, 552, 554, 571, 767, 768, 769, 775, 777, 861, 873, 885, 1118 |
| <i>Lepomis punctatus</i> | 69, 184, 299, 305, 331, 364, 378, 453, 463, 489, 552, 554, 767, 768, 769, 777, 861, 873, 885 |
| <i>Leptonycteris nivalis</i> | 1012 |
| <i>Leptonycteris sanborni</i> | 942, 1102, 1154 |
| <i>Lepus alleni</i> | 1423 |
| <i>Lepus californicus</i> | 47, 156, 197, 344, 370, 595, 671, 681, 685, 715, 820, 973, 974, 978, 1059, 1083, 1086, 1100, 1101, 1129, 1130, 1137, 1146, 1147, 1149, 1340, 1400, 1401, 1416, 1423, 1425, 1491, 1512, 1526 |
| <i>Lepus</i> | 1068, 1400, 1423 |
| <i>Libellula</i> | 359 |
| <i>Limnodromus griseus</i> | 64, 124, 131, 1351 |
| <i>Limnys irroratus</i> | 923, 926, 927, 928, 932, 936, 938, 939, 956, 961, 977, 979, 983, 984, 993, 995, 1036, 1041, 1042, 1091, 1094, 1107, 1246, 1247, 1249, 1488 |
| <i>Liposcellis bostrycophilus</i> | 220 |
| <i>Littorina irrorata</i> | 247, 400, 498 |
| <i>Lophodytes cucullatus</i> | 1387 |
| <i>Lophortyx gambelii</i> | 22 |
| <i>Loris gracilis</i> | 870 |
| <i>Lymnaea cubensis</i> | 375 |
| <i>Lynx</i> | 1466 |
| <i>Macrotus californicus</i> | 1037, 1102, 1141, 1150 |
| <i>Mareca americana</i> | 1016, 1389 |

| | |
|---------------------------------------|---|
| <i>Marmota flaviventris</i> | 1308, 1516 |
| <i>Masticophis flagellum</i> | 652 |
| <i>Melamysus bidentatus</i> | 281 |
| <i>Melanitta doglandi</i> | 1164 |
| <i>Meleagris gallopavo</i> | 1059, 1064 |
| <i>Meleagris gallopavo intermedia</i> | 137, 149, 199, 647, 1207, 1317 |
| <i>Melopelia asiatica mearnsi</i> | 37, 51 |
| <i>Melospiza lincolni</i> | 987 |
| <i>Melospiza melodia</i> | 1298, 1359 |
| <i>Menidia beryllina</i> | 269, 478 |
| <i>Menticirrus littoralis</i> | 472 |
| <i>Mephitis mephitis</i> | 47, 142, 158, 622, 623, 675, 726, 765, 848, 940, 1059, 1065, 1087, 1092, 1112, 1262, 1380, 1382, 1410, 1476, 1491, 1495, 1512 |
| <i>Mephitis</i> | 1514, 1519 |
| <i>Mergus serrator</i> | 1355 |
| <i>Mergus</i> | 1016 |
| <i>Meriones unguiculatus</i> | 824 |
| <i>Mesocricetus auratus</i> | 824 |
| <i>Micropogon undulatus</i> | 330, 394, 472, 770 |
| <i>Micropterus dolomieu</i> | 274 |
| <i>Micropterus punctulatus</i> | 184, 186, 224, 273, 274, 299, 417, 438, 441, 453, 489, 512, 552, 571, 767, 769, 777, 861, 873, 1118 |
| <i>Micropterus salmoides</i> | 69, 184, 185, 186, 187, 227, 230, 273, 274, 291, 299, 305, 331, 333, 367, 368, 376, 378, 417, 437, 438, 439, 441, 442, 453, 454, 463, 489, 506, 509, 511, 512, 516, 550, 552, 554, 571, 575, 616, 721, 767, 769, 777, 861, 872, 873, 1115, 1116, 1118 |
| <i>Microtus longicaudus</i> | 259, 1272, 1275, 1398, 1426, 1435, 1482 |
| <i>Microtus</i> | 1459 |
| <i>Micrurus fulvius</i> | 584, 651 |
| <i>Mimus polyglottos</i> | 1003, 1059, 1086 |
| <i>Mimus polyglottos leucopterus</i> | 1202, 1303 |
| <i>Minytrema melanops</i> | 453 |
| <i>Mniotilta varia</i> | 986 |
| <i>Mollienesis formosa</i> | 248, 252, 851 |
| <i>Mollienesis latipinna</i> | 89, 248, 249, 250, 251, 252, 480, 851 |
| <i>Molothrus ater</i> | 985, 1086 |
| <i>Molothrus ater ater</i> | 866 |
| <i>Mormoops megalophylla</i> | 444, 678, 988, 1041, 1049 |
| <i>Moxostoma congestum</i> | 387, 453, 777, 885 |
| <i>Mugil cephalus</i> | 34 |
| <i>Mus musculus</i> | 52, 126, 887, 897, 900, 952, 958, 1094, 1292, 1338, 1339, 1416, 1430, 1526 |
| <i>Mus</i> | 1396, 1433, 1481, 1485, 1523 |
| <i>Muscivora forficata</i> | 1052, 1059, 1333, 1357 |
| <i>Musculium transversum</i> | 277 |
| <i>Mustela frenata</i> | 1459 |
| <i>Mustela vison</i> | 621, 687 |
| <i>Myadestes townsendi townsendi</i> | 1324 |
| <i>Mycteria americana</i> | 1179, 1209, 1210, 1217, 1312 |
| <i>Myiarchus cinerascens</i> | 902, 1051 |
| <i>Myiarchus crinitus</i> | 1059 |
| <i>Myiochanes richardsonii</i> | 1059 |
| <i>Myotis californicus</i> | 1124 |
| <i>Myotis leibii</i> | 1024, 1025, 1048, 1102 |
| <i>Myotis lucifugus</i> | 201, 446, 1462 |
| <i>Myotis thysanodes</i> | 843, 1025, 1048, 1109, 1124, 1155, 1461 |
| <i>Myotis velifer</i> | 55, 75, 491, 540, 714, 907, 915, 943, 945, 984, 1018, 1023, 1025, 1046, 1048, 1102, 1109, 1140, 1152, 1155, 1460, 1461 |
| <i>Myotis yumanensis</i> | 1109, 1155 |
| <i>Nassarius vibex</i> | 340 |

| | |
|---|---|
| <i>Neotoma albigula</i> | 52, 542, 903, 937, 939, 1308, 1394, 1436, 1439, 1476, 1478, 1479, 1499 |
| <i>Neotoma cinerea</i> | 541, 542, 1437, 1476 |
| <i>Neotoma floridana</i> | 898, 958, 1059, 1086, 1276 |
| <i>Neotoma lepida</i> | 15, 541, 1443, 1477, 1479, 1517 |
| <i>Neotoma mexicana</i> | 1308, 1475, 1499 |
| <i>Neotoma micropus</i> | 52, 53, 655, 657, 892, 900, 903, 904, 927, 928, 932, 947, 967, 1055, 1057, 1065, 1083, 1086, 1088, 1101, 1107, 1308, 1309, 1392, 1393, 1395, 1396, 1417, 1419, 1421, 1423, 1436, 1438, 1439, 1443, 1444, 1447, 1453, 1471, 1476, 1491, 1499, 1511, 1512 |
| <i>Neotoma stephensi</i> | 541, 1499, 1515 |
| <i>Neotoma</i> | 903, 904, 1101, 1436, 1438, 1443, 1459, 1474, 1479, 1514 |
| <i>Nerodia erythrogaster transversa</i> | 325, 409, 447, 611, 616, 749, 775, 1527 |
| <i>Nerodia fasciata</i> | 323, 468, 651 |
| <i>Nerodia rhombifera</i> | 166, 651 |
| <i>Notemigonus chrysoleucas</i> | 316, 317, 319, 453 |
| <i>Notiosorex crawfordi</i> | 1101 |
| <i>Notophthalmus meridionalis</i> | 263, 395, 403, 584 |
| <i>Notropis lutrensis</i> | 320, 378, 453, 571 |
| <i>Notropis venustris</i> | 443, 453 |
| <i>Numenius americanus</i> | 79, 312, 377, 389, 390, 396, 428, 429, 431, 457, 867, 1185, 1231, 1296 |
| <i>Nuttalliornis borealis</i> | 1035, 1054 |
| <i>Nyctea scandiaca</i> | 1373 |
| <i>Nyctanassa violacea</i> | 1180, 1209 |
| <i>Nycticorax nycticorax</i> | 1178, 1209 |
| <i>Ochotona princeps</i> | 1391, 1407, 1439, 1459, 1525 |
| <i>Odocoileus hemionus</i> | 1, 95, 220, 580, 613, 806, 845, 1079, 1081, 1110, 1128, 1233, 1372 |
| <i>Odocoileus virginianus</i> | 1, 2, 3, 45, 46, 51, 346, 613, 631, 640, 690, 696, 759, 786, 845, 1059, 1060, 1065, 1068, 1075, 1079, 1082, 1084, 1086, 1087, 1097, 1112, 1126, 1128, 1129, 1135, 1136, 1138 |
| <i>Ondatra zibethicus</i> | 432, 656, 756, 819 |
| <i>Onthophagus gazella</i> | 540, 631, 734 |
| <i>Onychomys leucogaster</i> | 893, 920, 932, 947, 1055, 1083, 1101, 1248, 1275, 1337, 1390, 1393, 1411, 1414, 1416, 1421, 1423, 1440, 1441, 1444, 1447, 1450, 153, 1454, 1457, 1467, 1476, 1483, 1511, 1512, 1514, 1517, 1520 |
| <i>Onychomys</i> | 1514 |
| <i>Opheodrys aestivus</i> | 260 |
| <i>Ophisaurus ventralis</i> | 584, 1097 |
| <i>Oreoscoptes montanus</i> | 1086 |
| <i>Ortalis vetula maccalli</i> | 200, 609 |
| <i>Orthopristis chrysopterus</i> | 455 |
| <i>Oryzomys palustris</i> | 824, 931 |
| <i>Ostrea equestris</i> | 268 |
| <i>Otus asio</i> | 1284 |
| <i>Otus trichopsis</i> | 22 |
| <i>Ovis aries</i> | 639, 1138 |
| <i>Ovis canadensis</i> | 95, 220, 222, 639, 641, 686, 689, 745, 827, 829, 1079, 1081, 1110 |
| <i>Ovis trageiaphus</i> | 639, 762, 764 |
| <i>Pandion haliaetus carolinensis</i> | 408 |
| <i>Parabuteo unicinctus harrisi</i> | 973, 992 |
| <i>Passarella ilica ilica</i> | 558 |
| <i>Passer domesticus</i> | 970, 1004, 1022 |
| <i>Passer domesticus domesticus</i> | 30, 31, 41, 1201, 1325 |
| <i>Passerina ciris</i> | 1059 |
| <i>Passerina cyanea</i> | 1008 |
| <i>Passerherbulus caudacutus</i> | 866 |
| <i>Pelecanus erythrorhynchoss</i> | 575 |
| <i>Penalus aztecus</i> | 183 |
| <i>Penalus setiferus</i> | 183 |
| <i>Percina</i> | 453 |
| <i>Periplaneta americana</i> | 871 |

| | |
|---|--|
| <i>Perisoreus canadensis</i> | 1005 |
| <i>Perognathus flavus</i> | 889, 891, 925, 1245, 1248, 1442 |
| <i>Perognathus hispidus</i> | 126, 138, 191, 778, 889, 891, 893, 914, 957, 996, 1107, 1250, 1271, 1276, 1339, 1474 |
| <i>Perognathus intermedius</i> | 847 |
| <i>Perognathus nelsoni</i> | 1442 |
| <i>Perognathus penicillatus</i> | 743, 846, 923, 1442 |
| <i>Perognathus</i> | 898, 922, 1059, 1087, 1521 |
| <i>Peromyscus boylii</i> | 52, 54, 900, 918, 932, 1275, 1396, 1433, 1457, 1474 |
| <i>Peromyscus californicus</i> | 25, 832, 833 |
| <i>Peromyscus crinitus</i> | 1433 |
| <i>Peromyscus difficilis</i> | 1275, 1337, 1436, 1457, 1500 |
| <i>Peromyscus eremicus</i> | 1431, 1433 |
| <i>Peromyscus eremicus nest</i> | 1518 |
| <i>Peromyscus gossypinus</i> | 1084 |
| <i>Peromyscus leucopus</i> | 672, 898, 900, 903, 922, 927, 932, 956, 964, 977, 993, 995, 1045, 1059, 1087, 1094, 1101, 1275, 1428, 1457, 1474, 1515, 1521 |
| <i>Peromyscus maniculatus</i> | 27, 73, 900, 903, 1101, 1275, 1337, 1396, 1419, 1422, 1426, 1433, 1434, 1436, 1439, 1447, 1457, 1458, 1459, 1474, 1483, 1484, 1487, 1499 |
| <i>Peromyscus pectoralis</i> | 900, 1427, 1474, 1486 |
| <i>Peromyscus truei</i> | 54, 903, 1275, 1337, 1436, 1457, 1459, 1474 |
| <i>Peromyscus</i> | 1057, 1083, 1254, 1275, 1418, 1419, 1427, 1436, 1450, 1474, 1514 |
| <i>Petrochelidon albifrons</i> | 1077 |
| <i>Petrochelidon fulva pallida</i> | 933, 1078, 1103 |
| <i>Petrochelidon pyrrhonota</i> | 1001, 1194, 1406 |
| <i>Peucedramus olivaceus</i> | 32 |
| <i>Phainopepla nitens</i> | 1006 |
| <i>Phainopepla nitens nitens</i> | 1197 |
| <i>Phalaenoptilus nuttallii</i> | 1059 |
| <i>Phasianus colchicus</i> | 77, 337, 646, 647, 648, 703 |
| <i>Pheucticus melanocephalus</i> | 901 |
| <i>Phrynosoma cornutum</i> | 49, 98, 727, 766, 848, 928 |
| <i>Phrynosoma douglassi</i> | 168 |
| <i>Phyopsocus</i> | 220 |
| <i>Physa anatina</i> | 493 |
| <i>Physa frontinalis acuta</i> | 344 |
| <i>Physa halei</i> | 453 |
| <i>Physa virgata</i> | 359 |
| <i>Pimephales vigilax</i> | 299, 372, 453 |
| <i>Pipilo fuscus</i> | 1008, 1032 |
| <i>Pipistrellus hesperus</i> | 37, 55, 942, 959, 1023, 1056, 1102, 1155, 1461 |
| <i>Pipistrellus subflavus</i> | 915 |
| <i>Piranga rubra</i> | 1029, 1059 |
| <i>Pituophis melanoleucus sayi</i> | 651 |
| <i>Plecotus rafinesqii</i> | 28, 1000, 1038 |
| <i>Plecotus townsendi</i> | 945, 1102, 1103, 1104, 1109, 1123, 1151, 1152 |
| <i>Plegadis chihi</i> | 1181, 1215, 1281, 1336 |
| <i>Plegadis falcinellus</i> | 1281, 1336 |
| <i>Plegadis falcinellus falcinellus</i> | 1181 |
| <i>Plethodon neomexicanus</i> | 805 |
| <i>Podiceps caspicus californicus</i> | 1168 |
| <i>Pogonias cromis</i> | 24, 382, 488 |
| <i>Polyborus cheriway auduboni</i> | 1216 |
| <i>Polynemus octonemus</i> | 419 |
| <i>Pomatomus saltatrix</i> | 486 |
| <i>Pomoxis annularis</i> | 186, 187, 274, 291, 296, 297, 299, 305, 437, 441, 453, 552, 571, 575, 767, 857, 72, 1117, 1119, 1120 |
| <i>Pomoxis nigromaculatus</i> | 297, 453, 510, 857 |
| <i>Pooecetes gramineus</i> | 1008, 1086 |
| <i>Procyon lotor</i> | 147, 169, 343, 347, 375, 433, 608, 610, 628, 676, 796, 863, 1059, 1065, 1089, |

| | |
|---|---|
| | 093, 1095, 1097, 1380, 1383, 1401, 1410, 1416, 1423, 1473, 1491, 1495, 1496 |
| <i>Proptera purpurata</i> | 270, 301 |
| <i>Pseudacris triseriata</i> | 88, 261, 399, 584 |
| <i>Pseudemys alabamensis</i> | 19 |
| <i>Pseudemys concinna texana</i> | 19, 391 |
| <i>Pseudemys scripta elegans</i> | 19, 23, 282, 283, 374, 450, 451, 461, 553, 620, 777, 872, 874, 875 |
| <i>Pseudemys scripta troostii</i> | 391 |
| <i>Ptychocheilus lucius</i> | 172 |
| <i>Pyrrhuloxia sinuata</i> | 1032, 1324 |
| <i>Quadrula foresheyi</i> | 254, 301, 448 |
| <i>Quadrula houstonensis</i> | 254, 301, 448 |
| <i>Quiscalus quiscula</i> | 562, 566, 570, 602, 780, 864 |
| <i>Quadrula quiscula aeneus</i> | 564 |
| <i>Quiscalus versicolor</i> | 266, 334 |
| <i>Rallus elegans</i> | 86, 253, 312, 335, 338, 385, 411, 460, 1013 |
| <i>Rallus longirostris</i> | 86, 175, 177, 253, 255, 275, 335, 338, 385, 420, 459, 554 |
| <i>Rana areolata</i> | 399, 584 |
| <i>Rana catesbeiana</i> | 173, 189, 350, 351, 352, 353, 357, 358, 360, 399, 584, 618, 625, 701, 750, 853, 1120 |
| <i>Rana clamitans</i> | 189, 360, 399, 584, 750 |
| <i>Rana palustris</i> | 584, 700 |
| <i>Rana pipiens complex</i> | 87, 173, 350, 357, 358, 359, 399, 584, 618, 750 |
| <i>Rana sphenocephala</i> | 262, 351, 358, 363, 398, 399, 584, 700, 750 |
| <i>Rangia cuneata</i> | 278 |
| <i>Rattus mephiles</i> | 1394 |
| <i>Rattus norvegicus</i> | 9, 133, 870, 897, 900, 918, 929, 932, 933, 950, 952, 953, 958, 963, 997, 1011, 1017, 1083, 1277, 1339, 1403, 1409, 1410, 1414, 1415, 1416, 1423, 1430, 1455, 1463, 1465, 1476, 1488, 1491, 1506, 1512, 1526 |
| <i>Rattus rattus</i> | 93, 126, 170, 664, 798, 869, 870, 871, 897, 915, 918, 932, 950, 951, 954, 958, 1059, 1084, 1087, 1277, 1339, 1409, 1410, 1416, 1430, 1455, 1465, 1491, 1526 |
| <i>Rattus</i> | 1277, 1408, 1416, 1430, 1465, 1485, 1491, 1526 |
| <i>Recurvirostra americana</i> | 1015, 1230 |
| <i>Reithrodontomys fulvescens</i> | 1094 |
| <i>Reithrodontomys megalotis</i> | 1474 |
| <i>Reithrodontomys</i> | 891 |
| <i>Richmondia cardinalis</i> | 1007, 1059 |
| <i>Richmondia cardinalis cardinalis</i> | 1202 |
| <i>Richmondia cociferus</i> | 1086 |
| <i>Rynchops nigra</i> | 1345, 1362, 1369 |
| <i>Salmo apache</i> | 307, 665 |
| <i>Salmo gairdneri</i> | 307, 753 |
| <i>Salmo gilae</i> | 305, 665 |
| <i>Salmo trutta</i> | 7, 307, 665, 753 |
| <i>Salpinctes obsoletus</i> | 973, 1009 |
| <i>Sardar sardeu</i> | 474, 482 |
| <i>Sayornis nigricans</i> | 1031 |
| <i>Sayornis saya</i> | 1053 |
| <i>Sayornis saya saya</i> | 1333 |
| <i>Scalopus aquaticus</i> | 870 |
| <i>Scaphiopus bombifrons</i> | 934 |
| <i>Scaphiopus hammondi</i> | 934 |
| <i>Scardafella inca</i> | 48 |
| <i>Sceloporus clarki</i> | 38 |
| <i>Sceloporus jarrovi</i> | 38 |
| <i>Sceloporus magister</i> | 171, 589, 730 |
| <i>Sceloporus merriami</i> | 710 |
| <i>Sceloporus olivaceus</i> | 928, 1043, 1097 |
| <i>Sceloporus poinsetti</i> | 37 |
| <i>Sceloporus undulatus</i> | 700, 710 |

| | |
|--|--|
| <i>Sceloporus undulatus hyacinthinus</i> | 1097 |
| <i>Scincella lateralis</i> | 90, 261, 403, 584, 700, 731, 807, 1097 |
| <i>Sciurus aberti</i> | 1469 |
| <i>Sciurus arizonensis</i> | 1470 |
| <i>Sciurus carolinensis</i> | 645, 787, 897, 898, 1059, 1097, 1297, 1311, 1473 |
| <i>Sciurus niger</i> | 95, 192, 467, 645, 673, 787, 870, 898, 1059, 1064, 1097, 1297, 1311, 1410, 1469, 1473 |
| <i>Sialia currucoides</i> | 1010, 1086 |
| <i>Sialia mexicana bairdi</i> | 1328 |
| <i>Sialia sialis sialis</i> | 1200, 1328 |
| <i>Sigmodon hispidus</i> | 93, 121, 126, 155, 191, 192, 217, 219, 258, 519, 559, 629, 644, 656, 658, 659, 662, 663, 664, 722, 724, 756, 788, 799, 824, 871, 888, 893, 896, 897, 898, 914, 927, 928, 944, 957, 958, 962, 974, 977, 1045, 1059, 1094, 1101, 1102, 1107, 1276, 1339, 1393, 1416, 1423, 1465, 1488, 1497, 1526, 659, 784, 923, 1087 |
| <i>Sigmodon</i> | 659, 784, 923, 1087 |
| <i>Sistrurus miliarius</i> | 469 |
| <i>Speotyto cunicularia</i> | 963, 1027, 1101 |
| <i>Spermophilus lateralis</i> | 1415, 1480 |
| <i>Spermophilus mexicanus</i> | 672, 927, 956, 1047, 1059, 1107, 1242, 1512, 1514 |
| <i>Spermophilus spilosoma</i> | 932, 963, 1083, 1098, 1101, 1242, 1305, 1306, 1308, 1416, 1423, 1453, 1512 |
| <i>Spermophilus tereticaudus</i> | 932, 1044, 1083, 1107, 1241, 1242, 1306, 1416, 1508 |
| <i>Spermophilus tridecemlineatus</i> | 870, 1083, 1101, 1453, 1510, 1512, 1513 |
| <i>Spermophilus variegatus</i> | 4, 932, 1059, 1083, 1093, 1098, 1306, 1307, 1415, 1416, 1424, 1491, 1515 |
| <i>Spermophilus</i> | 1514 |
| <i>Spilogale putorius</i> | 142, 158, 675, 726, 765, 848, 1384 |
| <i>Spinitectus carolini</i> | 571, 873 |
| <i>Spinus psaltria</i> | 1059 |
| <i>Spizella arborea</i> | 1086 |
| <i>Spizella passerina</i> | 1008 |
| <i>Sizella pusilla</i> | 1059 |
| <i>Squalus acanthius</i> | 560 |
| <i>Squatarola squatarola</i> | 1014 |
| <i>Sterna antillarum</i> | 1058, 1189, 1369 |
| <i>Sterna fuscata</i> | 1348 |
| <i>Sterna hirundo</i> | 1161, 1352, 1353, 1369 |
| <i>Sterna maxima</i> | 955, 1161, 1350, 1353, 1369 |
| <i>Sternotherus carinatus</i> | 19, 282, 777 |
| <i>Storeria dekayi</i> | 264, 403, 584, 751 |
| <i>Strix varia</i> | 917, 1377 |
| <i>Strongylura marina</i> | 269, 478 |
| <i>Sturnella magna</i> | 30, 39, 40, 41, 266 |
| <i>Sturnella magna magna</i> | 58, 78, 747, 866 |
| <i>Sturnella neglecta</i> | 1086, 1100 |
| <i>Sturnus vulgaris</i> | 76, 127, 140, 141, 176, 178, 221, 384, 868, 881, 897, 905, 910, 915, 916, 960, 1050, 1086, 1090, 1143, 1145, 1237, 1238, 1300, 1302, 1404, 1419, 1528 |
| <i>Sturnus vulgaris vulgaris</i> | 1195, 1379 |
| <i>Sus scrofa</i> | 42, 346, 538, 626, 631, 666, 734, 782, 1059, 1061, 1084, 1097, 1264, 1491 |
| <i>Sylvilagus aquaticus</i> | 1399, 1423 |
| <i>Sylvilagus auduboni</i> | 197, 216, 595, 903, 928, 947, 978, 996, 1083, 1086, 1100, 1101, 1110, 1131, 1276, 1393, 1400, 1416, 1423, 1424, 1425, 1444, 1467, 1476, 1491, 1511, 1512 |
| <i>Sylvilagus floridanus</i> | 80, 198, 344, 370, 606, 823, 898, 919, 961, 1059, 1086, 1129, 1130, 1401, 1416, 1423 |
| <i>Sylvilagus nuttalli</i> | 1400 |
| <i>Sylvilagus</i> | 47, 973, 1068, 1399, 1400, 1423, 1526 |
| <i>Synodus foetens</i> | 418 |
| <i>Tachycineta thalassina</i> | 1405 |
| <i>Tadarida brasiliensis</i> | 21, 128, 328, 412, 423, 490, 531, 677, 908, 946, 965, 975, 1025, 1038, 1041, 1070, 1102, 1109, 1152, 1158, 1460, 1498, 1502, 1505 |

| | |
|---------------------------------------|--|
| <i>Tadarida femorosacca</i> | 942 |
| <i>Tagelus plebeius</i> | 280 |
| <i>Tamias dorsalis</i> | 1452 |
| <i>Tamias minimus</i> | 1273, 1451 |
| <i>Tamias quadrivittatus</i> | 1273, 1310, 1451 |
| <i>Tamias</i> | 1436, 1437 |
| <i>Tamiasciurus hudsonicus</i> | 1456, 1472, 1480, 1507 |
| <i>Tangavius aeneus aeneus</i> | 1324 |
| <i>Taxidea taxus</i> | 47, 1084, 1491 |
| <i>Tayassu tajacu</i> | 5, 152, 346, 604, 707, 713, 733, 804, 820, 1059, 1060, 1065, 1067, 1084, 1086, 1097, 1318, 1429 |
| <i>Tayassu</i> | 1429 |
| <i>Terrapene carolina triunguis</i> | 452, 545, 584, 587, 617, 700 |
| <i>Terrapene ornata</i> | 19, 700 |
| <i>Thais haemastoma</i> | 400 |
| <i>Thalasseus maximus</i> | 955 |
| <i>Thamnophis proximus</i> | 651, 751, 948 |
| <i>Thomomys bottae</i> | 1255, 1413, 1421 |
| <i>Thomomys talpoides</i> | 1421, 1436 |
| <i>Thomomys</i> | 1414 |
| <i>Thryomanes bewickii</i> | 973, 1059, 1086 |
| <i>Thryothorus ludovicianus</i> | 1059 |
| <i>Tilapia mossambica</i> | 753, 777 |
| <i>Totanus calidris</i> | 1014 |
| <i>Totanus flavipes</i> | 1014 |
| <i>Totanus melanoleucus</i> | 1160, 1184, 1354 |
| <i>Toxosoma curvirostre</i> | 973, 1086, 1303, 1323 |
| <i>Toxostoma rufum</i> | 517 |
| <i>Tramea</i> | 359 |
| <i>Triatoma ambigua</i> | 52 |
| <i>Triatoma gerstreckeri</i> | 52 |
| <i>Triatoma longipes</i> | 52 |
| <i>Triatoma rubida</i> | 52 |
| <i>Triatoma sanguisuga</i> | 52 |
| <i>Trichiurus lepturus</i> | 486 |
| <i>Tringa solitaria solitaria</i> | 1205, 1297 |
| <i>Trionyx muticus</i> | 19 |
| <i>Trionyx spiniferus emoryi</i> | 19, 23 |
| <i>Troglodytes aedon</i> | 1405 |
| <i>Turdus migratorius</i> | 1378 |
| <i>Tympanuchus pallidicinctus</i> | 200, 648, 703, 722 |
| <i>Tyrannus dominicensis</i> | 1052 |
| <i>Tyrannus melancholicus</i> | 973 |
| <i>Tyrannus melancholicus couchii</i> | 1303, 1324 |
| <i>Tyrannus tyrannus</i> | 1052 |
| <i>Tyrannus verticalis</i> | 902, 1052, 1101 |
| <i>Tyto alba</i> | 1375 |
| <i>Tyto alba pratincola</i> | 1286 |
| <i>Urocyon cinereoargenteus</i> | 47, 146, 216, 218, 528, 529, 615, 642, 729, 771, 880, 896, 897, 922, 958, 1059, 1084, 1087, 1093, 1097, 1099, 1385, 1401, 1409, 1416, 1423, 1473, 1491, 1496 |
| <i>Urosaurus ornatus</i> | 37, 710, 718 |
| <i>Ursus americanus</i> | 631, 812 |
| <i>Uta stansburiana</i> | 1043 |
| <i>Viero griseus</i> | 1066 |
| <i>Virginia striatulus</i> | 751 |
| <i>Vulpes macrotis</i> | 13, 47 |
| <i>Vulpes vulpes</i> | 1401, 1495 |
| <i>VWilsonia citrina</i> | 986 |
| <i>VWilsonia pusilla</i> | 986 |

| | |
|--|------------------------|
| <i>Zapus</i> | 1515 |
| <i>Zenaida asiatica</i> | 971 |
| <i>Zenaida asiatica asiatica</i> | 1218 |
| <i>Zenaida macroura</i> | 672, 972, 1059, 1139 |
| <i>Zenaida macroura marginella</i> | 1219, 1221, 1331, 1332 |
| <i>Zonotrichia leucophrys</i> | 1086 |
| <i>Zonotrichia leucophrys gambelli</i> | 1198 |

| Host Common Name | Parasite Number |
|------------------------|--|
| Alexandrian Rat | 1416 |
| Amphibian | 358, 361, 362 |
| Angora Goat | 1232 |
| Armadillo | 265, 539, 543, 725, 733, 860, 879, 1108, 1495 |
| Badger | 1096, 1416, 1491 |
| Bat | 1070, 1092, 1103, 1106, 1109, 1152, 1154, 1157, 1461, 1503, 1504, 1505 |
| Bird | 932, 933, 1363, 1416 |
| Ground Ground Bird | 1059 |
| Black-bellied Plover | 1019 |
| Black Skimmer | 1363 |
| Black-throated Sparrow | 1100 |
| Black Vulture | 1073 |
| Boa Constrictor | 1062 |
| Bobcat | 1097, 1416, 1423, 1491 |
| Bobwhite Quail | 1086 |
| Bovine | 579 |
| Box Turtle | 1063 |
| Brush Mouse | 1274, 1397, 1415, 1426, 1487, 1501 |
| Bullfrog | 660 |
| Burro | 1064 |
| Burrowing Owl | 1416, 1491 |
| Calf | 1410, 1416 |
| Canids Wild | 143, 213, 215, 216, 219, 343, 375, 529, 605, 695, 729, 771, 808, 837 |
| Cardinal | 1086 |
| Carnivore | 1059 |
| Cat | 104, 105, 219, 243, 529, 811, 1409, 1410, 1416 |
| Domestic | 773, 1252 |
| House | 1084, 1430 |
| Cattle | 17, 91, 95, 151, 154, 344, 345, 346, 425, 547, 548, 554, 576, 580, 58, 582, 639, 40, 641, 681, 682, 692, 696, 699, 742, 760, 817, 818, 831, 912, 999, 1059, 1060, 1064, 1067, 1075, 1079, 1097, 1110, 1127, 1134, 1136 |
| Civet Cat | 1254, 1485 |
| Chicken | 535, 554, 567, 649, 734, 1258, 1259, 1416 |
| Domestic | 1301 |
| Chicken Hawk | 1288 |
| Clam | 254, 301 |
| Cliff Chipmunk | 1273, 1310, 1452 |
| Cliff Swallow | 1102, 1107 |
| Coati | 1040, 1490 |
| Common Tern | 1370 |
| Cotton Rat | 1065, 1084, 1444, 1488 |
| Cottontail Rabbit | 1065, 1083, 1086, 1393, 1400, 1416, 1423, 1444, 1467, 1495, 1511, 1512 |
| Cow | 17, 120, 346, 599, 640, 1076, 1084, 1111, 1112, 1263 |
| Coyote | 1067, 1084, 1093, 1096, 1097, 1416, 1423, 1476, 1491 |
| Crayfish | 306, 341 Deer 45, 1059, 1064, 1086, 1097, 1110, 1114, 1125 |
| Deer Mouse | 1337, 1416, 1419, 1426, 1457, 1481, 1487, 1501 |
| Dog | 104, 105, 157, 211, 212, 216, 235, 404, 528, 529, 601, 605, 607, 695, 771, 772, 786, 808, 809, 810, 837, 878, 1060, 1065, 1067, 1079, 1084, 1097, 1110, 1112, 1265, 1409, 1410, 1416, 1473, 1488, 1491 |
| Domestic | 1267, 1381 |
| Duck | 60, 65, 100, 101, 102, 129, 130, 132, 134, 135, 336, 339, 410, 521, 614, 855 |
| Elk | 613 |
| Fox | 1087, 1093, 1097, 1099, 1401, 1416, 1473, 1495 |
| Free-tailed Bat | 1106 |
| Fulvous Tree Duck | 1159 |
| Gadwall | 102, 130, 132, 135, 336, 339, 410, 521, 614, 855, 1387 |

| | |
|-------------------------|---|
| Garfish | 815 |
| Goat | 95, 220, 344, 549, 578, 580, 582, 598, 633, 634, 640, 686, 697, 711, 741, 818, 820, 831, 911, 1064, 1084, 1086, 1100, 1112, 1291 |
| Goose Domestic | 1163 |
| Gopher | 1254, 1416 |
| Grasshopper Mouse | 1278, 1337, 1390, 1393, 1414, 1416, 1423, 1444, 1447, 1453, 1457, 1467, 1476, 1483, 1511, 1512 |
| Gray Fox | 1059, 1084, 1423, 1491 |
| Gray Squirrel | 897, 1423 |
| Ground Squirrel | 1415, 1416, 1423, 1453, 1476, 1512, 1514 |
| Guinea Pig | 548, 1257, 1261 |
| Gull-billed Tern | 1370 Harris' |
| Ground Squirrel | 1242, 1306, 1416 |
| Hog, Feral | 344, 345, 346 |
| Horse | 62, 63, 313, 329, 635, 636, 637, 638, 693, 704, 709, 738, 761, 790, 791, 792, 818, 838, 911, 1064, 1069, 1072, 1079, 1082, 1132, 1410 |
| Horse Fly | 613 |
| Jack Rabbit | 1065, 1083, 1086, 1400, 1423, 1495, 1512 |
| Japanese Bettle | 752 |
| Javalina | 1084 |
| Kangaroo Rat | 1393, 1416, 1423, 1433, 1444, 1447, 1453, 1476, 1491, 1511, 1512 |
| Laughing Gull | 1367 |
| Least Tern | 1058 |
| Leopard Frog | 660 |
| Mammal | 932, 933, 1416 |
| Man | 104, 209, 813, 958, 1059, 1064, 1084, 1093, 1097, 1110, 1112, 1156, 1319, 1329, 1409, 1410, 1416, 1491, 1524, 1526 |
| Meadow Lark | 1085, 1086, 1100 |
| Merriam's Kangaroo Rat | 1240, 1248, 1394, 1416 |
| Mexican Free-tailed Bat | 1505 |
| Mexican Woodrat | 1309, 1394, 1471, 1478, 1481, 1487 |
| Mouse | 1430 |
| House Mouse | 1526 |
| Mule | 62, 63, 313, 329, 635, 636, 637, 638, 693, 704, 709, 738, 761, 790, 791, 792, 818, 838, 1112, 1410 |
| Mule Deer | 1079 |
| Norwegian Rat | 1416 |
| Opossum | 144, 1064, 1084, 1097, 1101, 1416, 1473, 1488, 1491, 1495, 1526 |
| Ostrich | 569, 654 |
| Owl | 1285, 1376 |
| Oyster | 223 |
| Packrat | 1083, 1084, 1393, 1444, 1447, 1453, 1476, 1511 |
| Den | 1101 |
| Peccary | 152, 1064 |
| Texas | 1080 |
| Pig | 92, 106, 523, 536, 537, 568, 626, 632, 650, 667, 668, 691, 734, 782, 813, 836, 862, 1410 |
| Domestic Pig | 92 |
| Pigeon | 22, 1220 |
| Domestic | 1203 |
| Pintail | 100, 101, 129, 130, 132, 134, 336, 339, 410, 521, 614, 855 |
| Pocket Gopher | 1414, 1512 |
| Pocket Mouse | 1416, 1453 |
| Prairie Dog | 1096, 1101, 1393, 1412, 1416, 1423, 1467, 1476, 1491 |
| Quail | 1060, 1067 |
| Rabbit | 80, 94, 96, 157, 371, 548, 684, 688, 819, 1059, 1068, 1400, 1410, 1416, 1423, 1526 |
| Domestic | 1416, 1423, 1526 |
| Raccoon | 147, 375, 1084, 1093, 1097, 1099, 1401, 1410, 1416, 1423, 1473, 1491, 1495 |
| Rat | 958 |
| Domestic | 1410, 1414, 1416, 1423, 1430, 1488, 1491, 1512, 1526 |
| Red Squirrel | 1469 |

| | |
|------------------------|---|
| Red Wolf | 1495 |
| Ring-billed Gull | 1367 |
| Ringtail | 1096, 1476, 1491 |
| Roadrunner | 1086 |
| Rock Squirrel | 1415, 1416 |
| Rock Mouse | 1274, 1397, 1415 |
| Rodent | 1059 |
| Royal Tern | 1368 |
| Ruminant | 220 |
| Sanderling | 1371 |
| Sandpiper | 1349, 1371 |
| Sandwich Tern | 1365 |
| Semipalmated Plover | 1371 |
| Sheep | 95, 153, 220, 344, 548, 549, 577, 578, 580, 582, 598, 613, 633, 634, 640, 683, 686, 697, 741, 745, 818, 820, 831, 998, 1110 |
| Domestic | 220 |
| Shoveler | 101, 132, 336, 339, 410, 614, 855 |
| Silvery Mouse | 1509 |
| Nest | 1509 |
| Skunk | 1093, 1096, 1101, 1410, 1416, 1476, 1487, 1495, 1512, 1514, 1522 |
| Snail | 279 |
| Sparrow | 1333 |
| Spiny Pocket Mouse | 1488 |
| Striped SKunk | 1491 |
| Swallow | 1402, 1406 |
| Tree Owl | 1416 |
| Tree Squirrel | 1473 |
| Turkey | 1416 |
| Western Harvest Mouse | 1487 |
| White-collared Peccary | 152,713, 734, 706 |
| White-tailed Deer | 344, 345, 909, 1067, 1069, 1079, 1136, 1233, 1491 |
| White-throated Woodrat | 1309 , 1394 |
| White-winged Dove | 50 |
| White-throated Woodrat | 1416, 1477 |
| Wolf | 1059,1084, 1093, 1096, 1097, 1267, 1410, 1416, 1491 |
| Wood Rat | 1416, 1423, 1443, 1491, 1512 |
| Wren | 1087 |

LITERATURE CITED

- 1 Aldrich, D.V. 1965. Observations on the ecology and life cycle of Prochristianella penaei Kruse (Cestoda: Trypanorhyncha). *Journal of Parasitology* 51:370-376.
- 2 Alicata, J.E. 1932. The occurrence of Moniezia benedeni in a peccary. *Journal of Parasitology* 19:83.
- 3 Allen, R.W. 1949. Note on the natural occurrence of farm-raised chickens of encysted third stage larvae of Physocephalus sexalatus, a spuriid stomach worm of swine. *Proceedings of the Helminthological Society of Washington* 16:1-3.
- 4 Allen, R.W., and P.K. Jackson. 1950. The pathologic changes associated with Thysanosoma actinioides. *Journal of Parasitology* 36(Suppl.):45.
- 5 Allen, R.W., and P.M. Kyles. 1952. Thysanosoma actinioides with five suckers. *Proceedings of the Helminthological Society of Washington* 19:37-38.
- 6 Allen, R.W. 1952. Parasites in bighorn sheep in New Mexico. *Proceedings of the Helminthological Society of Washington* 19:39.
- 7 Allen, R.W., and P.K. Jackson. 1952. Seasonal variation in acquisition of Thysanosoma actinioides. *Journal of Parasitology* 38(Suppl.):25.
- 8 Allen, R.W. 1953. Preliminary observations on the incidence and intensity of nematodes in sheep on irrigated pastures in New Mexico. *Report from the Conference on Parasites and Parasitic Diseases of Domestic Ruminants* p. 17.
- 9 Allen, R.W., and P.K. Jackson. 1953. The occurrence of the fringed tapeworm, Thysanosoma actinioides, in pronghorn antelope. *Proceedings of the Helminthological Society of Washington* 20:96-97.
- 10 Allen, R.W., and P.K. Jackson 1954. The domestic sheep, a new host for Cooperia bisonis. *Proceedings of the Helminthological Society of Washington* 21:61.
- 11 Allen, R.W. 1955. Parasites of mountain sheep in New Mexico, with new host records. *Journal of Parasitology* 41:583-587.
- 12 Allen, R.W. 1955. Studies on the life history of the fringed tapeworm, Thysanosoma actinioides. *Report from the Conference on Parasites and Parasitic Diseases of Domestic Ruminants* pp. 19-20.
- 13 Allen, R.W., W.W. Beckland, and R.E. Gilmore. 1956. Parasites of Barbary sheep. *Journal of Parasitology* 42(Suppl.):19.
- 14 Allen, R.W., and W.W. Beckland. 1958. Cooperia mcmasteri and Cooperia surnabada in the United States. *Journal of Parasitology* 44:503-506.
- 15 Allen, R.W., G.A. Schad, and K.S. Samson. 1958. Experimental cross-transmission of two strains of Haemonchus from wild ruminants to domestic sheep, with observations on their pathogenicity as compared with Haemonchus from domestic sheep. *Journal of Parasitology* 44(Suppl.):26.
- 16 Allen, R.W. 1959. The occurrence of Nematodirus lanceolatus in domestic sheep in the United States. *Journal of the Colorado Wyoming Academy of Science* 4:53.
- 17 Allen, R.W. 1959. Preliminary notes on the larval development of the fringed tapeworm in sheep, Thysanosoma actinioides Diesing, 1834, in psocids. (Psocoptera: Corrodentia). *Journal of Parasitology* 45:537-538.
- 18 Allen, R.W., and K.S. Samson. 1960. Further observations on the occurrence of Thysanosoma actinioides in the American pronghorn. *Journal of Parasitology* 46:671.
- 19 Allen, R.W., and H.G. Erling. 1964. Parasites of bighorn sheep and mule deer in Arizona with new host records. *Journal of Parasitology* 50(Suppl.):71.
- 20 Allison, T.C., and J.L. McGraw. 1967. The helminth parasites of Centrarchidae from the Navasota River system of Texas. *Texas Journal of Science* 19:326-328.
- 21 Allison, T.C. 1967. Three new species of monogenetic trematodes from the gills of Lepomis cyanellus Rafinesque and Lepomis megalotis Rafinesque of Texas and the proposal of a new genus, Macrohaptor. *Journal of Parasitology* 53:1005-1007.
- 22 Amin, O.M. 1969. Helminth fauna of suckers (Catostomidae) of the Gila River System, Arizona. 1. Nematobothrium texomensis McIntosh and Self, 1955 (Trematoda) and Glaridacris confusus Hunter, 1929 (Cestoda) from buffalofish. *American Midland Naturalist* 82:188-195.
- 23 Anderson, R.R. 1968. Helminth and copepod parasites of the largemouth bass, Micropeterus salmoides (Lacepede) from Lake Whitney, Bosque County, Texas. M.A. Thesis, Baylor University.
- 24 Armstrong, H.W. 1969. A survey of the helminth parasites of certain Cyprinodontidae and Poeciliidae of the Texas coast. M.S. Thesis, Texas A&M University.
- 25 Arnold, L.W. 1942. Notes on the life history of the sand pocket mouse. *Journal of Mammalogy* 23:339-341.
- 26 Augustson, G.F. 1944. The flea genus Thrassia and sylvatic plague, with a description of T.

- brennani n. sp. Journal of Parasitology 30:237-240.
- 27 Augustson, G.F. 1944. A new mouse flea, Pleochaetodies bullisji, n. gen., n. sp., from Texas. Journal of Parasitology 30:366-368.
 - 28 Augustson, G.F. 1945. A new genus, new species of dermanyssid mite (Acarina) from Texas. Bulletin of the Southern California Academy of Science 44:46-48.
 - 29 Augustson, G.F., and L.C. Ryan. 1948. The flea genus Phyncopsyllus in the United States. Bulletin of the Southern California Academy of Science 67:111-112.
 - 30 Babcock, O.G., and H.E. Ewing. 1938. A new genus and species of Anoplura from the peccary. Proceedings of the Entomological Society of Washington 40:197-201.
 - 31 Babero, B., and D. Matthias. 1967. Thubunaea cnemidophorus n. sp., and other helminths from lizards, Cnemidophorus tigris, in Nevada and Arizona. Transactions of the American Microscopical Society 86:173-177.
 - 32 Badley, J.E., and N.O. Dronen, 1979. Some helminth parasites of the common grackle of Southern Texas. Proceedings of the Helminthological Society of Washington 46:149-150.
 - 33 Baker, D.W., and W.F. Fisher. 1966. Demodectic parasites in livestock. 70th Annual Meeting of the United States Livestock Association pp. 409-416.
 - 34 Baker, D.W., and W.F. Fisher. 1969. The incidence of demodectic mites in the eyelids of various mammalian hosts. Journal of Economic Entomology 62:942.
 - 35 Baker, E.W. 1951. Pneumocoptes, a new genus of lung-inhibiting mite from rodents (Acarina: Epidermotidae). Journal of Parasitology 37:583-586.
 - 36 Baldauf, R.J. 1961. Another case of parasitic copepods on amphibians. Journal of Parasitology 47:195.
 - 37 Bates, B.H., and T.G. Meade. 1972. Helminth parasites of the king rail, Rallus elegans Aud., and clapper rail, Rallus longirostris Boddaert, of the Bolivar Peninsula, Texas. Proceedings of the Helminthological Society of Washington 39:146-147.
 - 38 Becker, E.R., and R.D. Burroughs. 1933. Rediscovery of Eimeria carinii Pinto. Journal of Parasitology 20:123.
 - 39 Beckland, W.W., and R.W. Allen. 1958. Worm parasites of cattle in New Mexico and Arizona. Veterinary Medicine 53:586-590.
 - 40 Bedinger, C.A. 1967. Helminth parasites of East Texas fishes. M.S. Thesis, Sam Houston State University.
 - 41 Bedinger, C.A., and T.G. Meade. 1967. Biology of a new cercaria for Posthodiplostomum minimum (Trematoda: Diplostomidae). Journal of Parasitology 53:985-988.
 - 42 Beer, J.R., E.F. Cook, and R.G. Schwab. 1959. The ectoparasites of some mammals from the Chiricahua Mountains, Arizona. Journal of Parasitology 45:605-614.
 - 43 Behrens, C.B. 1970. A survey of platyhelminth and acanthocephalan parasites in Sturnus vulgaris L. of Nacogdoches County, Texas with emphasis on Cestoda and Acanthocephala. M.S. Thesis, Stephen F. Austin University.
 - 44 Bell, R.R., R.D. Turk, and T.J. Galvin. 1959. Incidence of certain gastrointestinal parasites of cattle in Texas. American Journal of Veterinary Research 20:766-767.
 - 45 Benton, A.H. 1955. The taxonomy and biology of Epitedia wenmanni (Rothschild, 1904) and E. testor (Rothschild, 1915)(Hystrichopsyllidae: Siphonaptera). Journal of Parasitology 41:491-495.
 - 46 Bishopp, F.C. 1911. Some new North American Ixodidae with notes on other species. Proceedings of the Biological Society of Washington 24:197-208.
 - 47 Bishopp, F.C. 1923. The rat mite attacking man. United States Department of Agriculture Bureau of Entomology Circular 294.
 - 48 Blachut, K. 1973. Seasonal independence and relative frequency of motile stages of scab mites, Psoroptes ovis, on sheep. Annals of the Entomological Society of America 66:285-287.
 - 49 Block, E.F. 1968. Parasites of the cricket frog, Acris crepitans. M.S. Thesis, North Texas State University.
 - 50 Boring, G.D. 1964. The infectivity of Bunostomum phlebotomum to sheep, rabbits, and guinea pigs. M.S. Thesis, Texas A&M University.
 - 51 Box, E.D. 1966. Blood and tissue protozoa of the English sparrow (Passer domesticus domesticus) in Galveston, Texas. Journal of Protozoology 13:204-208.
 - 52 Box, E.D. 1967. Influence of Isospora infections on patency of avian Lankesterella (Atoxoplasma, Garnham, 1950). Journal of Parasitology 53:1140-1147.
 - 53 Boyd, M.F. 1921. A possible intermediate host of Fasciola hepatica L. 1758 in North America. Journal of Parasitology 7:39-42.
 - 54 Bradshaw, G.V.R., and A. Ross. 1961. Ectoparasites of Arizona bats. Journal of the Arizona Academy of Science 1:109-112.
 - 55 Brennan, J.M. 1945. Field investigations pertinent to Bullis fever. Preliminary report on the species of ticks and vertebrates occurring at Camp Bullis, Texas. Texas Reports on Biology and Medicine 3:112-121.
 - 56 Brennan, J.M. 1945. Field investigations pertinent to Bullis fever. The Lone Star tick, Amblyoma americanum (Linnaeus, 1758). Texas Reports on Biology and Medicine 3:204-226.
 - 57 Bristol, J.R., L.F. Mayberry, R. Bueno, and A.G.

- Canaris. 1981. Coccidia of surface-feeding ducks from the central flyway, Texas. *Journal of Parasitology* 67:276-277.
- 58 Broderson, D., A.G. Canaris, and J.R. Bristol. 1977. Parasites of waterfowl from Southwest Texas: II. The shoveler, Anas clypeata. *Journal of Wildlife Diseases* 13:435-439.
- 59 Brooks, D.L., and R.W. Strandtman. 1960. The nasal mites of some West Texas flycatchers (Tyrannidae). *Journal of Parasitology* 46:418-432.
- 60 Brown, E.E. 1940. Biology of two freshwater trematodes: Analocreadium armatum (MacCallum, 1895) and Microcreadium parvum (Simer, 1929). M.S. Thesis, Texas A&M University.
- 61 Brown, J.C. 1944. A probable agent for the transmission of fowl paralysis. *Proceedings and Transactions of the Texas Academy of Science* 28:97-98.
- 62 Buchanan, G.D. 1956. Occurrence of the cestode Mathevotaenia surinamensis (Cohn, 1902) Spasskii, 1951 in a North American armadillo. *Journal of Parasitology* 42:34-38.
- 63 Buchanan, G.D. 1957. Occurrence of Cruzia americana in Texas nine-banded armadillos. *Journal of Parasitology* 43:92.
- 64 Buckner, C.B. 1972. An intestinal cestode of the Colorado River squawfish, Ptychocheilus lucius Girard. M.S. Thesis, Arizona State University.
- 65 Buechner, H.K. 1944. Helminth parasites of the gray fox. *Journal of Mammalogy* 25:185-188.
- 66 Bullock, W.L. 1957. Octospiniferoides chandleri, n. gen., n. sp., a neoechinorhynchid acanthocephalan from Fundulus grandis Baird and Girard on the Texas coast. *Journal of Parasitology* 43:97-100.
- 67 Burham, G.L. 1972. Some helminth parasites of the sandhill crane in West Texas. *Southwestern Naturalist* 17:200-201.
- 68 Buscher, H.N. 1965. Seasonal dynamics of the intestinal helminth fauna in three species of ducks. *Dissertation Abstracts* 26:2926
- 69 Buscher, H.N. 1965. Dynamics of the intestinal helminth fauna in three species of ducks. *Journal of Wildlife Management* 29:272-281.
- 70 Butler, W.F., and G.S. Pfaffenberger. 1981. Notes on parasites of the long-billed curlew, Numenius americanus from Eastern New Mexico. *Journal of Wildlife Diseases* 17:537-538.
- 71 Cain, D. 1966. Helminth parasites of bats from Carlsbad Caverns, New Mexico. *Journal of Parasitology* 52:351-357.
- 72 Cain, G.D., and E.H. Studier. 1972. Plagiorchis (Plagiorchis) micracanthos Macy, 1931 (Trematoda: Digenea) from Myotis lucifugus occultus in Northeast New Mexico. *Southwestern Naturalist* 17:104-105.
- 73 Cain, G.D., and E.H. Studier. 1974. Parasitic helminths of bats from Southwestern United States and Mexico. *Proceedings of the Helminthological Society of Washington* 41:113-114.
- 74 Canaris, A.G., A.C. Mena, and J.R. Bristol. 1981. Parasites of waterfowl from Southwest Texas. III. The green-winged teal. *Journal of Wildlife Diseases* 17:57-64.
- 75 Carter, J.M., A.G. Canaris, and J.R. Bristol. 1973. A survey of the parasites of starlings (Sturnus vulgaris L.) in El Paso County, Texas. *Journal of Parasitology* 59:775.
- 76 Casto, S., and B. McDaniel. 1967. Helminth parasitism in gars from South Texas with a description of Dichelyne lepisasteus n. sp. (Nematoda: Cucullanidae). *Proceedings of the Helminthological Society of Washington* 34:187-194.
- 77 Chandler, A.C. 1921. Notes on the occurrence of Moniliformis sp. in rats in Texas. *Journal of Parasitology* 7:179-183.
- 78 Chandler, A.C. 1922. Species of Hymenolepis as human parasites. *Journal of the American Veterinary Association* 78:636-639.
- 79 Chandler, A.C., and C.L. Suttles. 1922. A new rat tapeworm, Schizotaenia sigmodontis, from North America. *Journal of Parasitology* 8:123-128.
- 80 Chandler, A.C. 1929. A new species of trematode worm belonging to the genus Hassitesia from rabbits in Texas. *Publications of the United States National Museum* 75, Article 21.
- 81 Chandler, A.C. 1930. Specific characters in the genus Trichuris, with a description of a new species, Trichuris tenuis, from a camel. *Journal of Parasitology* 16:198-206.
- 82 Chandler, A.C. 1931. New genera and species of nematode worms. *Publications of the United States National Museum* 78, Article 23.
- 83 Chandler, A.C. 1932. Notes on the helminth parasites of the opossum (Didelphis virginiana) of Southeast Texas, with descriptions of four new species. *Publications of the U.S. National Museum* 81, Article 16.
- 84 Chandler, A.C. 1932. A n. sp. of Longistriata from the cotton rat, Sigmodontis hispidus, with notes on the division of the Heligmosominae into genera. *Journal of Parasitology* 19:25-31.
- 85 Chandler, A.C. 1941. Helminths of muskrats in Southeast Texas. *Journal of Parasitology* 27:175-181.
- 86 Chandler, A.C. 1941. Two new trematodes from the bonito, Sarder sarder, in the Gulf of Mexico. *Journal of Parasitology* 27:183-184.
- 987 Chandler, A.C. 1941. The specific study of Moniliformis (Acanthocephala) of Texas rats, and a review of the species of this genus in the Western hemisphere. *Journal of Parasitology* 27:241-244.
- 88 Chandler, A.C. 1941. Yellow fever, plague, and

- typhus - smoldering threats. Proceedings and Transactions of the Texas Academy of Science 25:61-62.
- 89 Chandler, A.C. 1942. The morphology and life cycle of a new strigeid, Fibricola texensis, parasitic in raccoons. Transactions of the American Microscopical Society 61:156-167.
 - 90 Chandler, A.C. 1942. The guinea worm, Dracunculus insignis (Leidy, 1858), a common parasite of raccoons in East Texas. American Journal of Tropical Medicine 22:153-157.
 - 91 Chandler, A.C. 1942. Helminths of tree squirrels in Southeast Texas. Journal of Parasitology 28:135-140.
 - 92 Chandler, A.C. 1942. Mesocestoides manteri n. sp. from a Lynx, with notes on other North American species of Mesocestoides. Journal of Parasitology 28:227-231.
 - 93 Chandler, A.C. 1942. The helminths of raccoons in East Texas. Journal of Parasitology 28:255-268.
 - 94 Chandler, A.C. 1943. A case of canine strongyloidiasis in Texas. Journal of Parasitology 29:157-158.
 - 95 Chandler, A.C. 1946. Trichuris peromysci n. sp. from Peromyscus californicus, and further notes on T. perognathi Chandler, 1945. Journal of Parasitology 32:208.
 - 96 Chandler, A.C. 1946. Helminths of armadillos, Dasybus novemcinctus, in eastern Texas. Journal of Parasitology 32:237-241.
 - 97 Chandler, A.C. 1946. Observations on the anatomy of Mesocestoides. Journal of Parasitology 32:242-245.
 - 98 Chandler, A.C., and R.A. Rausch. 1947. The species of the genus Filaria Mueller, 1787, s. str. Journal of Parasitology 33:449-452.
 - 99 Chandler, A.C., and R.A. Rausch. 1948. A contribution to the study of certain avian strigeids (Trematoda). Journal of Parasitology 34:207-210.
 - 100 Chandler, A.C. 1950. Gongylonema pulchrum in the black bear, Euarctos americanus, and the probable synonymy of G. pulchrum Molin, 1857, with G. ursi (Rudolphi, 1819). Journal of Parasitology 36:86-87.
 - 101 Chandler, A.C. 1950. Trichostrongylus calcaratus in muskrat. Journal of Parasitology 36:87.
 - 102 Chandler, A.C. 1950. Mesostephorus longisaccus, a new cyathocotyloid trematode from a dog. Journal of Parasitology 36:91.
 - 103 Chandler, A.C. 1951. Trematodes from the man-o'-war bird, Fregata magnificens Rothschild, on the Texas coast, with the description of a new species, Schwartzitrema seamsteri. Texas Journal of Science 3:186-189.
 - 104 Chandler, A.C. 1958. Some considerations relative to the nature of immunity in Trypanosoma lewisi infections. Journal of Parasitology 44:129-135.
 - 105 Christensen, Z.D., and D.B. Pence. 1977. Helminths of the plain chachalaca, Ortalis vetula maccalli, from the South Rio Grande Valley. Journal of Parasitology 63:830.
 - 106 Clark, G.G. 1972. Trypanosomes from mule deer in New Mexico and Colorado. Journal of Wildlife Diseases 8:325-326.
 - 107 Clayton, N.J., and E.A. Schlueter. 1971. A report on freshwater monogenetic trematodes of Garza-Little Elm Reservoir. Texas Journal of Science 22:159-168.
 - 108 Colbenson, H.P. 1962. Spinose ear tick Ornithodoros (Otobius) megnini, from pronghorn antelope. Journal of Parasitology 48:630.
 - 109 Conner, V. 1966. Helminths of the digestive systems of garfishes (Pisces: Lepisosteidae) from East-central Texas, with a survey of the literature on gar parasites. Texas Journal of Science 18:130.
 - 110 Cook, A.H. 1940. Screwworms infest beaver in Texas. Journal of Mammalogy 21:93.
 - 111 Cooley, R.A., and G.M. Kohls. 1939. Amblyomma philipi - a new tick from Texas and New Mexico, with a key to the known species of Amblyomma in the United States. (Acarina: Ixodidae) United States Public Health Reports, United States Treasury Department 54:44-47.
 - 112 Cooley, R.A., and G.M. Kohls. 1941. Further new species of Ornithodoros from bats (Acarina: Argasidae). United States Public Health Reports 59:910-914.
 - 113 Coombs, D.W., and M.D. Springer. 1974. Parasites of feral pig X European wild boar hybrids in Southern Texas. Journal of Wildlife Diseases 10:436-441.
 - 114 Couch, A.B. 1952. Blood parasites of some common Texas birds. Protozoology Journal, Field and Laboratory 20:146-154.
 - 115 Couch, A.B. 1963. Notes on the biology of Microlynchia pusilla Speiser, a lousefly of mourning doves. Journal of Parasitology 49:140-146.
 - 116 Craig, T.M., and R.R. Bell. 1978. Seasonal transmission of liver flukes to cattle in the Texas gulf coast. Journal of the American Veterinary Medicine Association 173:104-107.
 - 117 Craig, T.M. 1979. Seasonal transmission of bovine gastrointestinal nematodes in the Texas gulf coast. Journal of the American Veterinary Medicine Association 174:844-847.
 - 118 Crossley, D.A. 1952. Two new nasal mites from columbriform birds. Journal of Parasitology 38:385-390.
 - 119 Curfman, B.E., and F.F. Davidson. 1966. Parasitic helminths of Natrix erythrogaster transversa (Hallowell), the blotched water snake, in the Waco, Texas area. Texas Journal of Science 18:121.
 - 120 Custer, J.W., and D.B. Pence. 1979. Ectoparasites of the ringtail, Bassariscus astutus, from West Texas. Journal of Medical

- Entomology 15:132-133.
- 121 Custer, J.W., and D.B. Pence. 1981. Dirofilariasis in wild canids from the gulf coastal prairie of Texas and Louisiana, USA. *Veterinary Parasitology* 8:71-82.
 - 122 Custer, J.W., and D.B. Pence. 1981. Ecological analyses of helminth populations of wild canids from the gulf coastal prairies of Texas and Louisiana. *Journal of Parasitology* 67:289-307.
 - 123 Dancak, K., D.B. Pence, F.A. Stormer, and S.L. Beasom. 1982. Helminths of the scaled quail, Callipepla squamata, from Northwest Texas. *Helminthological Society of Washington* 49:144-146.
 - 124 Davies, R.B., and G.G. Clark. 1974. Trypanosomes from elk and horse flies in New Mexico. *Journal of Wildlife Diseases* 10:63-65.
 - 125 Davis, D.E. 1951. Observations on rat ectoparasites and typhus fever in San Antonio, Texas. *United States Public Health Reports* 66:1717-1726.
 - 126 Davis, G.E. 1940. Ticks and relapsing fever in the United States. *United States Public Health Reports* 55:2347-2350.
 - 127 Davis, J.R., and D.G. Huffman. 1977. A comparison of the helminth parasites of Gambusia affinis and Gambusia geiser (Osteichthyes: Poeciliidae) from the upper San Marcos River. *Southwestern Naturalist* 22:359-366.
 - 128 Davis, J.R., and Huffman, D.E. 1978. Some factors associated with the distribution of helminths among individual mosquitofish, Gambusia affinis. *Texas Journal of Science* 30:43-53.
 - 129 Davis, J.W. 1973. Deer ked infestation on white-tailed deer in East Texas. *Journal of Wildlife Management* 37:183-186.
 - 130 Davis, R.M., and R.B. Loomis. 1971. The intranasal chigger Micotrombicula merrihewi (Acarina: Trombiculidae) in the North American free-tailed bat, Tadarida brasiliensis. *Southwestern Naturalist* 15:437-458.
 - 131 Davis, R.W. 1971. Parasite endangers elk. *BioScience* 21:1221.
 - 132 DeBruin, D., and G.S. Pfaffenberger. 1984. Helminths of desert cotton-tail rabbits (Sylvilagus auduboni (Baird)) inhabiting prairie dog towns in Eastern New Mexico. *Proceedings of the Helminthological Society of Washington* 51:369-370.
 - 133 Degenhardt, W.G., and P.B. Degenhardt. 1965. The host-parasite relationship between Elaphe subocularis (Reptilia: Colubridae) and Aponomma elaphensis (Acarina: Ixodidae). *Southwestern Naturalist* 10:167-178.
 - 134 Denton, J.F. 1942. Eurytrema procyonis n. sp. (Trematoda: Dicrocoeliidae), from the raccon, Procyon lotor. *Proceedings of the Helminthological Society of Washington* 9:31-32.
 - 135 Denton, J.F. 1944. The occurrence of Eurytrema allentoshi (Foster, 1939) in the opossum in Texas. *Proceedings of the Helminthological Society of Washington* 11:54-55.
 - 136 Denton, J.F. 1944. Studies on the life history of Eurytrema procyonis Denton, 1942. *Journal of Parasitology* 30:277-286.
 - 137 Denton, J.F. 1945. Studies on the life history of Brachylecithum americanum n. sp., a liver fluke of passerine birds. *Journal of Parasitology* 31:131-141.
 - 138 Dewhirst, L.W., R.J. Trautman, and J.W. Pistor. 1958. Preliminary report on helminths of beef/cattle in Arizona. *Journal of Parasitology* 44:30.
 - 139 Dikmans, G. 1937. A note on the members of the nematode genus Trichostrongylus occurring in rodents and lagomorphs, with descriptions of two new species. *Journal of the Washington Academy of Science* 27:203-209.
 - 140 Douthitt, H. 1915. Studies on the cestode family Anoplocephalidae. *Illinois Biological Monographs* 1:1-96.
 - 141 Dove, W.E. and B. Shellmire. 1931. Tropical rat mites, Liponyssus bacoti, vectors of typhus fever virus. *Journal of the American Medical Association* 97:1506-1510.
 - 142 Dowell, J.H., R.J. Warren, and D.B. Pence. 1983. Helminth fauna of ring-necked pheasants from the Texas high plains. *Journal of Wildlife Diseases* 19:152-153.
 - 143 Dronen, N.O. 1975. The life cycle of Haematoloechus coloradensis Cort, 1915 (Digenea: Plagiorchiidae), with emphasis on host susceptibility to infection. *Journal of Parasitology* 61:657-660.
 - 144 Dronen, N.O. 1977. Studies on the population structure of two species of Haematoloechus Looss, 1899 (Digenea: Plagiorchiidae) in ranid frogs in New Mexico. *Proceedings of the Helminthological Society of Washington* 44:68-72.
 - 145 Dronen, N.O., H.T. Underwood, and M.T. Sunderman. 1977. Textrema hopkinsi gen. et sp. n. (Digenea: Cryptogonimidae) from the largemouth bass of Central Texas. *Journal of Parasitology* 63:282-284.
 - 146 Dronen, N.O., and J.E. Badley. 1979. Helminths of shorebirds from the Texas gulfcoast. I. Digenetic trematodes from the long-billed curlew, Numenius americanus. *Journal of Parasitology* 65:645-649.
 - 147 Dronen, N.O., and H.T. Underwood. 1980. Pseudomagnivitellinum ictalurum gen. et sp.n. (Digenea: Macroderoidae) from the black bullhead of South-Central Texas. *Proceedings of the helminthological Society of Washington* 47:33-36.
 - 148 Drummond, R.O. 1959. Cattle grubs in gullets of cattle from Southwestern Texas. *Journal of*

- Economic Entomology 51:1221-1222.
- 149 Drummond, R.O., and J.G. Medley. 1964. Occurrence of Speleognathus australis Womersley (Acarina: Speleognathidae) in the nasal passages of bison. *Journal of Parasitology* 50:655.
 - 150 Drummond, R.O. 1966. The common cattle grub in cattle in Southwestern Texas. *Journal of Economic Entomology* 59:1105-1108.
 - 151 Drummond, R.O. 1966. Lipoptena mazamae Rondani (Diptera: Hippoboscidae), a louse fly of deer on cattle in Southwestern Texas. *Journal of Parasitology* 52:825.
 - 152 Drummond, R.O. 1967. Seasonal activity of ticks (Acarina: Metastigmata) on cattle in Southwestern Texas. *Annals of the Entomological Society of America* 60:439-447.
 - 153 Dunagen, T.T. 1957. Larval trematodes (Gorgoderidae) from Central Texas. *Proceedings of the Helminthological Society of Washington* 24:87-93.
 - 154 Duszynski, D.W., W.A. Riddle, D.R. Anderson, and R.W. Mead. 1972. Coccidia from the tiger salamander, Ambystoma tigrinum, in Northeastern Colorado and in Northern New Mexico. *Journal of Protozoology* 19:252-257.
 - 155 Duszynski, D.W., and K.L. Jones. 1973. The occurrence of intradermal mites (Acarina: Trombiculidae) in anurans in New Mexico, with a histological description of the tissue capsule. *International Journal of Parasitology* 3:531-538.
 - 156 Eads, R.B. 1946. A new species of Rhopalopsyllus Baker (Siphonaptera) from Texas. *Journal of Parasitology* 32:407-408.
 - 157 Eads, R.B. 1946. A new species of flea from the field mouse, Baiomys taylori. *Annals of the Entomological Society of America* 39:545-548.
 - 158 Eads, R.B. 1948. Ectoparasites from a series of Texas coyotes. *Journal of Mammalogy* 29:268-271.
 - 159 Eads, R.B., and G.C. Menzies. 1948. An undescribed Anomiopsyllus Baker from the pack rat, Neotoma micropus Baird. *Journal of the Kansas Entomological Society* 21:133-136.
 - 160 Eads, R.B. 1949. A preliminary list of Siphonaptera of Texas. *Texas Journal of Science* 1:33-39.
 - 161 Eads, R.B., and G.C. Menzies. 1949. Meringis bilsingi, a new ectoparasite of the Ord kangaroo rat, Dipodomys ordii Woodhouse. *Proceedings of the Entomological Society of Washington* 51:116-118.
 - 162 Eads, R.B., and G.C. Menzies. 1949. A new flea from the pocket gopher, Cratogeomys castanops Baird. *Journal of Parasitology* 35:171-174.
 - 163 Eads, R.B. 1950. An additional report of a reduviid bug other than Triatoma attacking man. *Journal of Parasitology* 36:87.
 - 164 Eads, R.B. 1950. The fleas of Texas. *Bulletin of the Texas State Department of Health, Austin, Texas*.
 - 165 Eads, R.B., and B.G. Hightower. 1950. Arthropods of possible medical significance collected in Terrell County, Texas. *Entomology News* 61:106-108.
 - 166 Eads, R.B., and G.C. Menzies. 1950. Fox ectoparasites collected incident to a rabies control program. *Journal of Mammalogy* 31:78-80.
 - 167 Eads, R.B. 1951. New mites of the genus Androlaelaps Berlese. *Journal of Parasitology* 37:212-216.
 - 168 Eads, R.B. 1951. A note on the ectoparasites of the javelina, or wild pig, Tayassu angulatus (Cope). *Journal of Parasitology* 37:317.
 - 169 Eads, R.B. 1951. A note on the javelina flea, Juxtapulex porcinus. *Journal of Parasitology* 37:328.
 - 170 Eads, R.B., and B.G. Hightower. 1951. A new mite from the nests of the wood rat, Neotoma micropus. *Entomology News*. 62:249-252.
 - 171 Eads, R.B., and B.G. Hightower. 1951. A new Neoichoronyssus from the pack rat, Neotoma micropus (Acarina: Laelaptidae). *Proceedings of the Entomological Society of Washington* 53:295-298.
 - 172 Eads, R.B., and B.G. Hightower. 1952. Blood parasites of Southwest Texas rodents. *Journal of Parasitology* 38:89-90.
 - 173 Eads, R.B., and B.G. Hightower. 1952. A new mite of the genus Neoichoronyssus. *Proceedings of the Entomological Society of Washington* 54:32-35.
 - 174 Eads, R.B., G.C. Menzies, and V.I. Miles. 1952. Acarina taken during West Texas plague studies. *Proceedings of the Entomological Society of Washington* 54:250-253.
 - 175 Eads, R.B., and O.L. Walker. 1952. Texas records of the tick, Ixodes tovari Colley. *Journal of Parasitology* 38:368.
 - 176 Eads, R.B., and W.W. Dalquest. 1954. An addition to the Texas flea fauna. *Journal of Parasitology* 40:102.
 - 177 Eads, R.B. 1956. Ectoparasites from swallow nests, with descriptions of a new ceratophyllid flea. *Journal of Parasitology* 42:73-76.
 - 178 Eads, R.B., G.C. Menzies, and B.G. Hightower. 1956. The ticks of Texas, with notes on their medical significance. *Texas Journal of Science* 8:7-24.
 - 179 Eads, R.B. 1960. Two new fleas (Hystriechopsyllidae) from the kangaroo rat and pocket mouse. *Journal of Parasitology* 46:213-218.
 - 180 Eads, R.B., H.A. Trevino, and E.G. Campos. 1965. Ectoparasites of the spiny pocket mouse, Liomys irroratus texensis. *Southwestern Naturalist* 10:17-21.
 - 181 Elbihari, S. 1968. Feather mites (Acarina: Analgesoidea) of some Texan and Mexican falconiform and strigeform birds. M.S. Thesis,

- Texas A&M University.
- 182 Eller, R.R. 1962. The physiological specificity of the spiny rat mite, Echinolaelaps echidninus, for Rattus norvegicus. M.S. Thesis, Texas Technological University.
 - 183 Emerson, H.R. 1969. A comparison of parasitic infestations of white-tailed deer (Odocoileus virginianus) from Central and East Texas. Bulletin of the Wildlife Disease Association 5:137-139.
 - 184 English, P.F. 1932. Some habits of the pocket gopher, Geomys breviceps breviceps. Journal of Mammalogy 13:126-132.
 - 185 Epstein, R.A. 1972. Larval trematodes of marine gastropods of Galveston Island, Texas. M.S. Thesis, Texas A&M University.
 - 186 Ewing, H.E., and I. Fox. 1943. The fleas of North America. United States Department of Agriculture Miscellaneous Publication 500.
 - 187 Farrell, C.E. 1956. Chiggers of the genus Euschongastia (Acarina: Trombiculidae) in North America. Publications of the United States Nature Museum 106:85-235.
 - 188 Ferris, G.F. 1942. Some North American rodent-infesting lice (Insecta: Anoplura). Microentomology 7:84-90.
 - 189 Fincher, G.T., and O.C. Marti. 1982. Onthophagus gazella as an intermediate host for spiruroids in Georgia and Texas. Southwestern Entomologist 7:125-129.
 - 190 Fisher, F.M. and J.D. Webster. 1954. A new strigeid trematode from the pectoral sandpiper. Journal of Parasitology 40:444.
 - 191 Fisher, W.F. 1960. On Acanthocephala of turtles, with the description of Neoechinorhynchus emyditoides, n. sp. Journal of Parasitology 46:257.
 - 192 Fisher, W.F. 1973. Natural transmission of Demodex bovis Stiles in cattle. Journal of Parasitology 59:223-224.
 - 193 Flook, J.M., and J.E. Ubelaker. 1972. A survey of metazoan parasites in unionid bivalves of Garza-Little Elm Reservoir, Denton County, Texas. Texas Journal of Science 23:381-392.
 - 194 Foreyt, W.J., and A.G. Todd. 1972. The occurrence of Fascioloides magna and Fasciola hepatica together in the livers of naturally infected cattle in South Texas, and the incidence of the flukes in cattle, white-tailed deer, and feral hogs. Journal of Parasitology 58:1010-1011.
 - 195 Foreyt, W.J., A.G. Todd, and K.M. Foreyt. 1975. Fascioloides magna (Bassi, 1875) in feral swine from Southern Texas. Journal of Wildlife Diseases 11:554-559.
 - 196 Foreyt, W.J., and K.M. Foreyt. 1979. Elaeophora schneideri in a white-tailed deer from Texas. Journal of Wildlife Diseases 15:55-56.
 - 197 Francis, M. 1897. The biology of the cattle ticks. Proceedings and Transactions of the Texas Academy of Science 3:95.
 - 198 Gaines, G.D., R.J. Warren, and D.B. Pence. 1984. Helminth fauna of sandhill crane populations in Texas. Journal of Wildlife Diseases 20:207-211.
 - 199 Galvin, T.J. 1957. Helminth parasites of the domestic cat. Southwestern Veterinarian 10:137-146.
 - 200 Gentner, H.W., and S.H. Hopkins. 1966. Changes in the trematode fauna of clams in the Little Brazos River, Texas. Journal of Parasitology 52:458-461.
 - 201 Gentner, H.W. 1968. Host parasite relationships of cottontail rabbits (Sylvilagus floridanus) and tapeworms of the genera Cittotaenia and Raillietina. M.S. Thesis, Texas A&M University.
 - 202 George, J.E. 1960. The nasal mites of the genus Ptilonyssus occurring in some North American passeriforme birds. M.S. Thesis, Texas Technological University.
 - 203 George, J.E. 1960. Notes on the parasitic mites of some West Texas reptiles. Southwestern Naturalist 5:105-106.
 - 204 George, J.E., and R.W. Strandtman. 1960. New records of ectoparasites on bats in West Texas. Southwestern Naturalist 5:228-229.
 - 205 George, R.R., and E.G. Bolen. 1975. Endoparasites of black-bellied whistling ducks in Southern Texas. Journal of Wildlife Diseases 11:17-22.
 - 206 Gilmore, R.E. 1955. The helminths parasitic in the lizard Cnemidophorus grahami, and a description of a new cestode from the lizard Holbrookia texana. M.S. Thesis, New Mexico State University.
 - 207 Gilmore, R.E., and R.W. Allen. 1960. Helminth parasites of pronghorn antelope (Antilocapra americana) in New Mexico with new host records. Proceedings of the Helminthological Society of Washington 27:69-73.
 - 208 Glazner, W.C., and F.F. Knowlton. 1967. Some endoparasites found in Welder Refuge deer. Journal of Wildlife Management 31:595-597.
 - 209 Goff, W.L., and N.C. Ronald. 1981. Certain aspects of the biology and life cycle of Heterobilharzia americana in East Central Texas. American Journal of Veterinary Research 42:1775-1777.
 - 210 Grogl, M., R.E. Kuhn, D.S. Davis, and G.E. Green. 1984. Antibodies to Trypanosoma cruzi in coyotes in Texas. Journal of Parasitology 70:189-191.
 - 211 Gruninger, T.L., C.E. Murphy, and J.C. Britton. 1977. Macroparasites of fish from Eagle Mountain Lake, Texas. Southwestern Naturalist 22:525-535.
 - 212 Guthery, F.S., S.L. Beasom, and L. Jones. 1979. Cerebrospinal nematodiasis caused by Parelaphosytrungylus tenuis in angora goats in Texas. Journal of Wildlife Diseases 15:37-41.

- 213 Haas, G.E., R.P. Martin, M. Swickard, and N. Wilson. 1972. Bird fleas (Siphonaptera) of New Mexico. *Canadian Entomologist* 104:881-883.
- 214 Harkema, R. 1942. Pharyngostomoides procyonis n.g., n. sp., (Strigeids), a trematode from the raccoon in North Carolina and Texas. *Journal of Parasitology* 28:117-122.
- 215 Harwood, P.D. 1931. A new species of Oxysomatium (Nematoda) with some remarks on the genera Oxysomatium and Aplectana and observations on the life history. *Journal of Parasitology* 17:61-73
- 216 Harwood, P.D. 1932. The helminths parasitic in the amphibia and reptilia of Houston, Texas, and vicinity. *Proceedings of the United States National Museum* 81, Article 17.
- 217 Hathaway, R.P. 1967. A survey of intestinal helminths in dogs from Albuquerque, New Mexico. *Journal of Parasitology* 53:1240.
- 218 Hays, L.M. 1950. Comparison of Mallophaga on related species of birds. M.S. Thesis, Texas Technological University.
- 219 Hedeon, R.A. 1953. A new species of Sarcoptes (Acarina: (Sarcoptidae) from the cave bat. *Journal of Parasitology* 39:334-335.
- 220 Hedeon, R.A. 1953. Ectoparasites occurring on mammals in the vicinity of Ft. Hood, Texas. *Texas Journal of Science* 5:125-129.
- 221 Hegner, R.W. 1926. Giardia beckeri n. sp. from the ground squirrel and Endamoeba dipodomysi from the kangaroo rat. *Journal of Parasitology* 12:203-206.
- 222 Hellgren, E.C., R.L. Lochmiller, and W.E. Grant. 1984. Infection of captive adult collared peccaries, Dicotyles tajacu (Woodburne, 1968), with the nematode Ascaris suum (Goeze, 1782). *Proceedings of the Helminthological Society of Washington* 51:160-161.
- 223 Henson, R.N., and S. Yilmaz. 1971. Incidence of Toxocara canis (Werner) and Ancylostoma caninum (Ercolani) from dogs in Brazos County, Texas. *Texas Journal of Science* 22:247.
- 224 Herrin, C.S., and C.E. Yunker. 1973. A new species of Hirstionyssus from Liomys irroratus in Texas (Acari: Mesostigmata). *Journal of Medical Entomology* 10:269-273.
- 225 Hibler, C.P., K.S. Samson, and R.W. Allen. 1963. Dictyocaulus filaria in Colorado and New Mexico sheep. *Journal of Parasitology* 49:652.
- 226 Hibler, C.P. 1965. Description of the microfilaria of Wehrdikmansia cervipedis (Wehr and Dikmans, 1935) and observations on its location in Arizona deer. *Bulletin of the Wildlife Disease Association* 1:44-48.
- 227 Hibler, C.P. 1966. Development of Stephanofilaria stilesi in the horn fly. *Journal of Parasitology* 52:890-898.
- 228 Hibler, C.P., G.H. Gates, R. White, and B.R. Donaldson. 1970. Observations on horseflies infected with larvae of Elaeophora schneideri. *Journal of Wildlife Diseases* 7:43-45.
- 229 Hightower, B.G., and R.B. Eads. 1951. New records of Texas Siphonaptera. *Journal of Parasitology* 37:328.
- 230 Hoff, C. C., and L.A. Williams. 1949. Notes on some fleas from rodent nests in New Mexico. *Journal of Parasitology* 35:545.
- 231 Hoffpauir, C., and E.O. Morrison. 1966. Rhabdias ranae from Bufo valliceps. *Southwestern Naturalist* 11:302.
- 232 Holdenreid, R., and H.G. Morlaw. 1956. A field study of wild mammals and fleas of Santa Fe County, New Mexico. *American Midland Naturalist* 55:369-381.
- 233 Hollis, P.D. 1972. A survey of parasites of the bullfrog, Rana catesbiana Shaw, in Central East Texas. *Southwestern Naturalist* 17:198-200.
- 234 Hopkins, S.H. 1937. A new type of allocreadiid cercaria: The cercariae of Anallocreadium and Microcreadium. *Journal of Parasitology* 23:94-97.
- 235 Hopkins, S.H. 1940. The excretory systems of Tergestia stossich 1899 and Cornucopula adunca (Linton 1905) (Trematoda). *Transactions of the American Microscopical Society* 59:281-284.
- 236 Hopkins, S.H. 1941. New genera and species of the family Monorchidae (Trematoda) with a discussion of the excretory systems. *Journal of Parasitology* 27:395-407.
- 237 Hopkins, S.H. 1951. Studies on larval marine bucephalids. *Journal of Parasitology* 37:13-14.
- 238 Hopkins, S.H. 1954. The american species of trematode confused with Bucephalus (Bucephalopsis) haimeanus. *Parasitology* 33:353-370.
- 239 Hopkins, S.H. 1958. Trematode parasites of Donax variabilis at Mustang Island, Texas. *Publications of the Institute of Marine Science, University of Texas* 5:301-311.
- 240 Hopkins, S.H. 1965. Haemogregarina and Haemoproteus (Protozoa: Sporozoa) in blood of Texas freshwater turtles. *Journal of Parasitology* 51:682-683.
- 241 Hopkins, S.H. 1966. A comparison of the parasites of two freshwater shads, Dorosoma (Dorosoma) cepedianum and Dorosoma (Signalosa) petenense, in Lake Texoma. *Journal of Parasitology* 52:843.
- 242 Hopkins, S.H., and H.W. Gentner. 1966. Changes in the trematode fauna of clams in the Little Brazos River, Texas. *Journal of Parasitology* 52:458-461.
- 243 Hopkins, S.H., J.W. Little, and F.G. Schict. 1966. Acanthoparyphium spinulosum (Trematoda: Echinostomatidae) in oysters at Port Isabel, *Journal of Parasitology* 52:663.
- 244 Hopkins, H. 1967. Rhipidocotyle lepisostei Hopkins, 1954, in gars caught in freshwater. *Journal of Parasitology* 53:491.
- 245 Hopkins, S.H., and J.W. Little. 1968. Neoechinorhynchus constrictus sp. n., an

- acanthocephalan from Texas turtles. Proceedings of the Helminthological Society of Washington 35:46-49.
- 246 Hopkins, S.H. 1969. Evidence from living tags that fish go eight hundred miles up Atchafalaya-Red River to Denison Dam. Texas Journal of Science 20:295.
- 247 Hopkins, S.H., and J.W. Little. 1969. New locality records for Taenia rileyi Loewen, 1929 and Taenia macrocystis Diesing, 1850, and a comparison of some hook measurements. Proceedings of the Helminthological Society of Washington 36:268-269.
- 248 Hopkins, S.H. 1971. On the taxonomic status and comparative morphology of species of the genus Neoapectana Steiner. Proceedings of the Helminthological Society of Washington 38:68-79.
- 249 Hubbard, C.A. 1949. Fleas of Western North America, Iowa State College Press.
- 250 Huggins, E.J. 1949. A survey of the helminths and ectoparasites of rats, Rattus rattus frugivorus Rafinesque and Sigmodon hispidus hispidus Say and Ord, in Brazos County, Texas. M.S. Thesis, Texas A&M University.
- 251 Huggins, E.J. 1951. A survey of helminths and ectoparasites of roof and cotton rats in Brazos County, Texas. American Midland Naturalist 46:230-244.
- 252 Ingham, R.E., and N.O. Dronen, 1980. Endohelminth parasites from largemouth bass, Micropterus salmoides, in Belton and Livingston Reservoirs, Central Texas. Proceedings of the Helminthological Society of Washington 47:140-142.
- 253 Jacobs, B. and E.O. Morrison. 1966. Ectopic lung fluke. Southwestern Naturalist 11:412.
- 254 Jameson, D.K. 1959. A survey of the parasites of five species of bats. Southwestern Naturalist 4:61-65.
- 255 Johnson, P.T. 1962. The species of Fahrenholzia Kellogg and Ferris, from spiny pocket mice (Anoplura: Hoplopleuridae). Annals of the Entomological Society of America 55:415-528.
- 256 Jones, A.W., B.W. Mounts, and G.B. Wolcott. 1944. A pronoccephalid trematode from the - Pseudemys. Proceedings and Transactions of the Texas Academy of Science 28:92-93.
- 257 Joy, J.E. 1970. Gryporhynchus tetrorchis Hill (Cestoda: Dilepididae) from a heron in Texas. Southwestern Naturalist 15:272.
- 258 Joy, J.E. 1971. Geographic distribution and host records for the genus Crassicutis (Trematoda: Allocreadiidae) with a new locality for Crassicutis archosargii. Folia Parasitologia 18:233-234.
- 259 Joy, J.E. 1971. Spirocamallanus pereirai (Nematoda: Camallanidae) from the croaker, Micropogon undulatus, in Texas. Journal of Parasitology 57:390.
- 260 Joy, J.E. 1972. A new species of Henneguya (Myxosporidae: Myxobolidae) from the black drum, Pogonias cromis, in Clear Lake, Texas. Journal of Protozoology 19:126-128.
- 261 Joy, J.E. 1972. A new species of Kudoa (Myxosporidae: Chloromyxidae) from the spot, Leiostomus xanthurus, in Clear Lake, Texas. Journal of Protozoology 19:264-265.
- 262 Joy, J.E. 1972. New locality records for Diplomonorchis leiostomi Hopkins, 1941 (Trematoda: Digenea) and Macrovalvitrematoides micropogoni Pearse, 1949 (Trematoda: Monogenea) with notes on their geographical distribution. Texas Journal of Science 23:553-554.
- 263 Karstad, L., and D.O. Trainer. 1969. Sarcocystis in white-tailed deer. Bulletin of the Wildlife Disease Association 5:25-26.
- 264 Kellogg, F.E., T.P. Kistner, R.K. Strickland, and R.R. Gerrish. 1971. Arthropod parasites collected from white-tailed deer. Journal of Medical Entomology 8:495-498.
- 265 Kemper, H.E. 1938. Filial dermatosis of sheep. North American Veterinarian 19:36-41.
- 266 Kirby, D.R. 1979. Prevalence of patent and occult filarial infections in stray dogs from the coastal bend area of Texas. Southwestern Veterinarian 32:121-123.
- 267 Knight, M.T., C.J. Barbay, and E.O. Morrison. 1965. Incidence and infection by lung-fluke (Haematoloechus) of the bullfrog, Rana catesbiana, in Jefferson County, Texas. Southwestern Naturalist 10:141-142.
- 268 Knight, R.A., and D.W. Tuff. 1984. Trichuris spp. (Nematoda: Trichuridae) in sika deer (Cervus nippon) in Texas. Proceedings of the Helminthological Society of Washington 51:161-162.
- 269 Kohls, G.M., and P.A. Cooley. 1936. Notes on the occurrence and host relationships of the tick Ornithodoros talaje in Arizona. United States Public Health Reports 51:512.
- 270 Kohls, G.M. 1940. Siphonaptera. A study of the species infecting wild hares and rabbits of North America, north of Mexico. United States Public Health Service Bulletin 175.
- 271 Kohls, G.M., and W.L. Jellison. 1948. Ectoparasites and other arthropods occurring in Texas bat caves. National Speleological Society Bulletin 10:116-117.
- 272 Kohls, G.M. 1950. Description of the males of Ixodes woodi Bishopp (Acarina: Ixodidae). Journal of Parasitology 36:65-66.
- 273 Kohls, G.M. 1958. Amblyoma imitator, a new species of tick from Texas and Mexico, and remarks on the synonymy of A. cajennense (Fabricus)
- 274 Kruidenier, F.J., and K.N. Mehra. 1958. Aspiculuris (Nematoda: Oxyuridae) from rats of the Grand Canyon National Park, Arizona. Journal of Parasitology 44:33.

- 275 Kruidenier, F.J., and K.N. Mehra. 1959. Aspicularis ackerti n. sp. (Nematoda: Oxyuridae) from the wood rats of Arizona. Proceedings of the Helminthological Society of Washington 26:147-150.
- 276 Kruidenier, F.J., and K.N. Mehra. 1959. Wellcomia perognathi n. sp. (Nematoda: Oxyuridae) from pocket mice, Perognathus intermedius (Meriam, 1889) of the Grand Canyon in Arizona. Transactions of the Illinois Academy of Science 51:20-25.
- 277 Kruidenier, F.J., and V. Gallicchio. 1959. The orthography of the Brachylaimidae (Joyeux and Foley, 1930): Brachylaima microti sp. nov.: B. raushii McIntosh, 1950; and an addendum to Dollfus (1935) list of Brachylaima (Trematoda: Digenea). Transactions of the American Microscopical Society 78:428-441.
- 278 Lang, E.M. 1957. Deer of New Mexico. New Mexico Game and Fish Bulletin 5
- 279 Lang, J.D. 1972. Ectoparasites found on the round-tailed ground squirrel near Tucson, Arizona. Journal of the Arizona Academy of Science 7:99-100.
- 280 Lawrence, J.L. and C.E. Murphy. 1967. Parasites of five species of fish from Benbrook Lake, Tarrant County, Texas. Texas Journal of Science 19:164-174.
- 281 LeCount, A. 1981. A survey of trichinosis among black bears of Arizona. Journal of Wildlife Diseases 17:349-351.
- 282 Lee, H. 1955. The mode of egg dispersal in Physaloptera phrynosoma (Spiruroidea), a gastric nematode of Texas horned toads, Phrynosonum cornutum. Journal of Parasitology 41:70-73.
- 283 Lee, S.H. 1957. The life cycle of Skrijabinoptera phrynosoma (Ortlepp) Schuly, 1927 (Nematoda: Spiruroidea), a gastric nematode of Texas horned toads, Phrynosoma cornutum. Journal of Parasitology 43:66-75.
- 284 Lee, T. 1971. Host-parasite relationships of the endoparasitic helminths of the eastern meadowlarks (Sturnella magna magna L.) in McLennan County, Texas. M.A. Thesis, Baylor University.
- 285 Lesser, E., K.F. Burns, and C.J. Farinacci. 1957. A blood parasite from the bat, Tadarida brasiliensis mexicana. Journal of Parasitology 43:380.
- 286 Levine, N.D., V. Ivens, and F.J. Kruidenier. 1955. Two new species of Klossia (Sporozoa: Adeleidae) from a deer mouse and a bat. Journal of Parasitology 41:623-629.
- 287 Leyva, T., A.G. Canaris, and J.R. Bristol. 1980. Helminths and ectoparasites of the common snipe (Capella gallinago L.) from Southwest Texas and Monte Vista Wildlife Refuge, Colorado. Journal of Wildlife Diseases 16:549-557.
- 288 Lindquist, A.W. 1934. Myiasis in wild animals in Southwestern Texas. Journal of Economic Entomology 30:735-740.
- 289 Little, J.W., S.H. Hopkins, and F.G. Schlicht. 1966. Acanthoparyphium spinulosum in oysters at Port Isabel, Texas. Journal of Parasitology 52:663.
- 290 Little, J.W., and S.H. Hopkins. 1968. Neoechinorhynchus constrictus n. sp., an acanthocephalan from Texas turtles. Proceedings of the Helminthological Society of Washington 35:46-49.
- 291 Little, J.W., J.P. Smith, F.F. Knowlton, and R.R. Bell. 1971. Incidence and geographic distribution of some nematodes in Texas bobcats. Texas Journal of Science 22:403-407.
- 292 Locke, L.N. and P. James. 1962. Trichomonad canker in the Inca dove, Scardafella inca. Journal of Parasitology 48:497.
- 293 Loomis, R.B., and D.A. Crossley. 1963. New species and new records of chiggers (Acarina: Trombiculidae) from Texas. Acarologia 5:371-383
- 294 Ludwid, R.G. 1983. Filaroides osleri in Texas canids. Southwestern Veterinarian 35:11-195.
- 295 Lumsden, R.D. 1963. Saccocoelioides sogandaresi sp. n., a new haploporid trematode from the sailfin molly, Mollienisia latipinna Le Sueur in Texas. Journal of Parasitology 49:281-284.
- 296 Lumsden, R.D. 1963. A new heterophyid trematode of the ascotyle complex of species encysted in poeciliid and cyprinodont fishes of Southwest Texas. Proceedings of the Helminthological Society of Washington 30:293-296.
- 297 Marchiondo, A.A., D.W. Duszynski, and G.O. Maupin. 1976. Prevalence of antibodies to Toxoplasma gondii in wild and domestic animals of New Mexico, Arizona and Colorado. Journal of Wildlife Diseases 12:226-232.
- 298 Martin, J.L., and D.G. Huffman. 1980. An analysis of the community and population dynamics of the helminths of Sigmodon hispidus (Rodentia: Cricetidae) from three Central Texas vegetation regions. Proceedings of the Helminthological Society of Washington 47:247-255.
- 299 Martin, L. 1933. Life history of the pigeon louse (Columbicola columbae Linnaeus). M.A. Thesis, Rice University.
- 300 Mayberry, L.F., J.R. Bristol, and D.W. Duszynski. 1980. Isospora californica (Protozoa: Eimeriidae) in Peromyscus maniculatus (Cricetidae) from White Sands National Monument, New Mexico. Southwestern Naturalist 25:125-126.
- 301 Mayberry, L.F., J.R. Bristol, D.W. Duszynski, and W.H. Reid. 1980. Eimeria macrotis sp.n. from Vulpes macrotis neomexicanus Merriam, 1902. Zeitschrift fur Parasitenkunde 61:197-200.
- 302 McDaniel, B., and R.F. Lawrence. 1962. A new

- genus and species of parasitic mite from Mexico (Acarina: Listrophoridae). *Journal of Parasitology* 48:463-466.
- 303 McDaniel, B. 1964. A redescription of the genus Olabidocarpus Lawrence, with a description of a new species from Texas (Acarina: Listrophoridae). *Acarologia* 6:163-169.
- 304 McDaniel, B. 1965. The subfamily Listrophorinae Gunther with a description of a new species of the genus Listrophorus pagenstecher from Texas (Acarina: Listrophoridae). *Acarologia* 7:704-712.
- 305 McDaniel, B., and I. Patterson. 1966. Nematode infestation of a white pelican found along the Gulf coast of Texas. *Southwestern Naturalist* 11:312.
- 306 McDaniel, B. 1967. A new distribution record and description of Speleognathopis benoiti Fain collected from the black-bellied plover in Texas. *Texas Journal of Science* 19:94-98.
- 307 McDaniel, B. 1968. A new species of Fahrenheitia (Anoplura: Hoplopleuridae) from Texas, with a key to the species of the genus and a list of their known hosts. *Annals of the Entomological Society of America* 61:402-405.
- 308 McDaniel, J.S. 1963. Parasites from the genus Lepomis (Centrarchidae) in Lake Texoma, Oklahoma. *Transactions of the American Microscopical Society* 82:423-425.
- 309 McGraw, J.L. 1964. Some internal helminth parasites of certain Centrarchidae of the Navasota and Little Rivers of Texas. M.S. Thesis, Texas A&M University.
- 310 McGraw, J.L., and T.C. Allison. 1967. Helminth parasites of Centrarchidae of the Little River system of Texas. *Southwestern Naturalist* 12:332-333.
- 311 McIntosh, A. 1932. Description of a tick, Dermacentor halli, from the Texas peccary, with a key to the North American species of Dermacentor. *Publications of the United States National Museum* 82, Article 4.
- 312 Meade, T.G., and C.A. Bedinger. 1967. Posthodiplostomum minimum (Trematoda: Diplostomidae) in fishes of Madison County, Eastern Texas. *Southwestern Naturalist* 12:334-335.
- 313 Meade, T.G. 1968. Helminth parasitism of fishes in East Texas. *American Philosophical Society Yearbook* pp. 312-313.
- 314 Meade, T.G., and J.S. Harvey. 1968. New record of numbers and sites of infection in fishes by the acanthocephalan, Eocollis arcanus Van Cleave. *Journal of Parasitology* 54:371.
- 315 Meade, T.G., and C.A. Bedinger. 1972. Helminth parasitism in some species of fresh water fishes of Eastern Texas. *Southwestern Naturalist* 16:281-295.
- 316 Meadows, J. 1968. Parasite community structure in three rodents, Sigmodon, Perognathus and Baiomys, in Central Texas. M.A. Thesis, Baylor University.
- 317 Meleney, W.P., and I.H. Roberts. 1970. Otobius megnini (Acarina: Argasidae) in the ears of pronghorn antelope (Antilocapra americana) in New Mexico. *Journal of Parasitology* 56:917.
- 318 Melvin, D.M., and A.C. Chandler. 1950. New helminth records from the cotton rat, Sigmodon hispidus, including a new species, Strongyloides sigmodontis. *Journal of Parasitology* 36:505-510.
- 319 Melvin, D.M. 1952. Studies on the life cycle and biology of Monoecocestus sigmodontis (Cestoda: Anoplocephalidae) from the cotton rat, Sigmodon hispidus. *Journal of Parasitology* 38:346-355.
- 320 Mendez, E., and G.E. Haas. 1972. A new flea of the genus Megarthroglossus Jordan and Rothschild from New Mexico (Siphonaptera: Hystrichopsyllidae: Anomiopsyllinae). *Journal of Medical Entomology* 9:285-288.
- 321 Menzies, G.C. 1949. Polypax serrata (Brumeister) and Linognathus setosus (Olfers) recorded from the house mouse, Mus musculus Linnaeus in Texas. *Journal of Parasitology* 35:435.
- 322 Menzies, G.C, R.B. Eads, and B.G. Hightower. 1951. List of Anoplura from Texas. *Proceedings of the Entomological Society of Washington* 53:150-152.
- 323 Menzies, G.C., and R.W. Strandtman. 1952. A new species of mite taken from a nest of armadillo (Acarina: Laelaptidae: Hypoaspinae). *Proceedings of the Entomological Society of Washington* 54:265-269.
- 324 Miller, W.L. 1940. A survey of hookworm infection in Brazos County, Texas. M.S. Thesis, Texas A&M University.
- 325 Mitchell, R.W., and W.L. Rhodes. 1960. New host records for the mesostigmatid nasal mite, Rhinonyssus rhinolethrum (Acarina: Rhinonyssidae). *Southwestern Naturalist* 5:107-108.
- 326 Mitchell, R.W. 1961. New avian host records for some mesostigmatid nasal mites. *Southwestern Naturalist* 6:103-105.
- 327 Mitchell, R.W. 1963. Comparative morphology of the life stages of the nasal mite Rhinonyssus rhinolethrum (Mesostigmata: Rhinonyssidae). *Journal of Parasitology* 49:506-515.
- 328 Mollhagan, T. 1978. Habitat influence on helminth parasitism of the cotton rat in Western Texas with remarks on some of the parasites. *Southwestern Naturalist* 23:401-408.
- 329 Monson, G., and W. Kessler. 1940. Life history notes on the banner-tailed kangaroo rat, Meriam's kangaroo rat, and the white-throated rat in Arizona and New Mexico. *Journal of Wildlife Management* 4:37-43.
- 330 Moore, D.V. 1946. Studies on the life history and development of Monoliformis dubius Meyer, 1933. *Journal of Parasitology* 32:257-271.

- 331 Moore, D.V. 1946. Studies on the life history and development of Macracanthorhynchus ingens Meyer, 1933, with a redescription of the adult worm. *Journal of Parasitology* 32:387-399.
- 332 Moore, D.V. 1962. Morphology, life history and development of the acanthocephalan Mediorhynchus grandis Van Cleave, 1916. *Journal of Parasitology* 48:76-86.
- 333 Morlan H.B. 1951. Notes on the genus Gigantolaelaps and a description of a new species, Gigantolaelaps cricelidorum (Acarina: Laelaptidae). *Journal of Parasitology* 37:273-279.
- 334 Morlan, H.B. 1954. Notes on the genus Megarhroglossus (Siphonaptera, Hystrichopsyllidae) in Santa Fe County, New Mexico. *Journal of Parasitology* 40:446-447.
- 335 Morrison, E.E., and H.T. Gier. 1979. Parasitic infection of Filaroides osleri, Capillaria aerophila and Spirocercia lupi in coyotes from the Southwestern U.S. *Journal of Wildlife Diseases* 15:557-560.
- 336 Mortan, H.B., and C.C. Hoff. 1957. Notes on some Anoplura from New Mexico and Mexico. *Journal of Parasitology* 43:347-351.
- 337 Morrison, E.O. 1966. Crayfish, possible secondary intermediate host for lung flukes (Haematoloechus) of bullfrogs in Jefferson County, Texas. *Yearbook of the American Philosophical Society* pp. 361-362.
- 338 Morrison, E.O. 1967. Human dipylidiasis: A case report from Vidor, Texas. *Texas Journal of Science* 19:231.
- 339 Morrison, E.O. 1968. Lungworms of Texas amphibia. *Yearbook of the American Philosophical Society* pp. 320-321.
- 340 Mpoame, M. 1984. Helminths of apache (Salmo apache), gila (S. gilae), and brown (S. trutta) trouts. *Southwestern Naturalist* 29:505-506.
- 341 Naderman, J., and D.B. Pence. 1980. Helminths of the common crow, Corvus brachyrhynchos Brehm, from West Texas. *Proceedings of the Helminthological Society of Washington* 47:100-105.
- 342 Najarian, H.H. 1961. The life cycle of Plagiorchis goodmani n. comb. (Trematoda: Plagiorchiidae). *Journal of Parasitology* 47:625-633.
- 343 Nowlin, W.C., C.E. Price, and E.A. Schlueter. 1967. First report on the fresh water monogenetic trematodes of Texas. *Texas Journal of Science* 19:110-111.
- 344 Olsen, O.W. 1947. Longevity of metacercariae of Fasciola hepatica on pastures in the upper coastal region of Texas and its relationship to liver fluke control. *Journal of Parasitology* 33:36-42.
- 345 Olsen, O.W. 1948. Wild rabbits as reservoir hosts of the common liver fluke, Fasciola hepatica, in Southern Texas. *Journal of Parasitology* 34:119-123.
- 346 Olsen, O.W. 1950. Pseudostertagia bullosa (Ransom and Hall, 1912) Orlov, 1933, from New Mexico sheep, with a note on the gubernaculum. *Veterinary Medicine* 45:163-164.
- 347 Owen, B.L. 1958. Records of nasal mites of the mourning dove. *Texas Journal of Science* 10:447.
- 348 Pacheco, G., and G.S. Tulloch. 1970. Microfilariae of Dirofilaria striata in a dog. *Journal of Parasitology* 56:248.
- 349 Packchanian, A. 1939. Natural infection of Triatoma gerstreckeri with Trypanosoma cruzi in Texas. *United States Public Health Reports* 54:1547-1554.
- 350 Packchanian, A. 1942. Reservoir hosts of Chagas' disease in the state of Texas. *American Journal of Tropical Medicine* 22:623-631.
- 351 Panigraphy, B., J.E. Grimes, S.E. Glass, S.A. Nagi, and C.F. Hall. 1982. Disease of pigeons and doves in Texas - clinical findings and recommendations for control. *Journal of the American Veterinary Medicine Association* 181:384-386.
- 352 Panitz, E. 1967. Thelandes salamandrae (Oxyuroidea) Schad, 1960 in Plethodon neomexicanus from Jemez Mountains, New Mexico. *Canadian Journal of Zoology* 45:1296-1297.
- 353 Parish, H.E., and C.S. Rude. 1946. DDT to control the winter horse tick. *Journal of Economic Entomology*. 39:92-93.
- 354 Parish, H.E. 1949. Recent studies on life history and habits of the ear tick. *Journal of Economic Entomology* 42:416-419.
- 355 Parmalee, P.W. and M.A. Price. 1953. Bruella illustris (Kellogg) and other ectoparasites from the bobwhite in Texas. *Journal of Parasitology* 39:222.
- 356 Pence, D.B. 1975. Eyeworms (Nematoda:Thelaziidae) from West Texas quail. *Journal of Parasitology* 42:181-183.
- 357 Pence, D.B. and S. Casto. 1975. Two new species and new records of nasal mites of the genus Sternostoma (Acarina: Rhinonyssinae) from birds in Texas. *Journal of Parasitology* 61:360-368.
- 358 Pence, D.B., and A.G. Canaris. 1976. Tinaminyssus juxtamelloi sp. n. (Acari: Dermanyssidae; Rhinonyssinae) from the nasal passages of the band-tailed pigeon, Columba fasciata, in New Mexico. *Journal of Parasitology* 62:116-118.
- 359 Pence, D.B., and S. Casto. 1976. Nasal mites of the subfamily Speleognathinae (Ereynetidae) from birds in Texas. *Journal of Parasitology* 62:466-469.
- 360 Pence, D.B., and J.E. Stone, 1977. Lungworms (Nematoda: Pneumospiruridae) from West Texas carnivores. *Journal of Parasitology* 63:979-991.

- 361 Pence, D.B., and S. Bickel. 1977. Helminths of wild turkeys in the West Texas. Proceedings of the Helminthological Society of Washington 44:104-105.
- 362 Pence, D.B., and K.D. Willis. 1978. Helminths of the ringtail, Bassariscus astutus, from West Texas. Journal of Parasitology 64:568-569.
- 363 Pence, D.B., and D.L. Sell. 1979. Helminths of the lesser prairie chicken, Tympanuchus pallidicinctus (Ridgway) (Tetraonidae), from the Texas panhandle. Proceedings of the Helminthological Society of Washington 46:146-149.
- 364 Pence, D.B., and W.P. Meinzer. 1979. Helminth parasitism in the coyote, Canis latrans, from the rolling plains of Texas. International Journal of Parasitology 9:339-344.
- 365 Pence, D.B., and S. Eason. 1980. Comparison of the helminth faunas of two sympatric top carnivores from the rolling plains of Texas. Journal of Parasitology 66:115-120.
- 366 Pence, D.B., V.E. Young, and F.S. Guthery. 1980. Helminths of the ring-necked pheasant, Phasianus colchicus (Gmelin) (Phasianidae) from the Texas panhandle. Proceedings of the Helminthological Society of Washington 47:144-147.
- 367 Pence, D.B., and G.G. Gray. 1981. Elaeophorosis in barbary sheep and mule deer from the Texas panhandle. Journal of Wildlife Diseases 17:49-56.
- 368 Pence, D.B., and M.J. Rhodes. 1982. Trichobilharzia physellae (Digenea: Schistosomatidae) from endemic waterfowl on the high plains of Texas. Journal of Wildlife Diseases 18:69-74.
- 369 Pence, D.B., J.T. Murphy, F.S. Guthery and T.B. Doerr. 1983. Indications of seasonal variations in the helminth fauna of the lesser prairie chicken, Tympanuchus pallidicinctus (Ridgway) (Tetraonidae) from Northwest Texas. Proceedings of the Helminthological Society of Washington 50:345-347.
- 370 Pence, D.B., and L.A. Windberg. 1984. Population dynamics across selected habitat variables of the helminth community in coyotes, Canis latrans, from South Texas. Journal of Parasitology 70:735-746.
- 371 Pfaffenberger, G.S., and W.F. Butler. 1981. Helminths recovered from the white-necked raven (Corvus cryptoleucus Couch) in Eastern New Mexico. Journal of Wildlife Diseases 17:563-566.
- 372 Pfaffenberger, G.S., B. Nygren, D. DeBruin, and C. Wilson. 1984. Parasites of the black-tailed prairie dog (Cynomys ludovicianus) from Eastern New Mexico. Proceedings of the Helminthological Society of Washington 51:241-244.
- 373 Phillips, C.J., J.K. Jones, and F.J. Radovsky. 1969. Macronyssid mites in oral mucosa of long-nosed bats: Occurrence and associated pathology. Science 165:1868-1869.
- 374 Pigott, R.S. 1978. Prevalence of Dirofilaria immitis infection in a pet dog from the Texas Gulf Coast. Southwestern Veterinarian 31:193-196.
- 375 Porter, J.C., and R.W. Strandtman. 1952. Nasal mites of the English sparrow. Texas Journal Science 4:393-399.
- 376 Pratt, H.D., and N.E. Good. 1954. Distribution of some common domestic rat ectoparasites in the United States. Journal of Parasitology 49:113-129.
- 377 Price, E.W. 1926. The occurrence and distribution of Cysticercus cellulosae in Texas swine. Journal of Parasitology 12:81-83.
- 378 Price, E.W. 1928. A list of helminth parasites occurring in Texas. Journal of Parasitology 14:200-201.
- 379 Price, M.A. 1958. A new species of tick from the Trans-Pecos region in Texas. Journal of Parasitology 44:649-651.
- 380 Price, R.W. 1943. A redescription of Heterobilharzia americana Price (Trematoda: Schistosomatidae). Proceedings of the Helminthological Society of Washington 10:85-86.
- 381 Price, W.W. 1928. The civet, Bassariscus astutus flavus, a new host for Uncinaria lotoris. Journal of Parasitology 14:197.
- 382 Pritchard, A.E. 1947. Haplopleura oenomydis Ferris a louse found on domestic rats in the United States. Journal of Parasitology 33:374-375.
- 383 Prudhoe, S., and B.R. Manger. 1966. Three species of hymenolepid cestodes from a charadriiform bird in North America. American Annual Magazine of Natural History 13:539-549.
- 384 Randolph, N.M., and R.B. Eads. 1946. Gross infection of the domestic rat with ectoparasites. Journal of Economic Entomology 39:538.
- 385 Randolph, N.M., and R.B. Eads. 1946. An ectoparasite survey of mammals from Lavaca County, Texas. Annals of the Entomological Society of America 39:597-601.
- 386 Read, C.P. 1946. Strigeids from Texas mink with notes on the genus Fibricola Dubois. Transactions of the American Microscopical Society 67:165-168.
- 387 Read, C.P. 1947. Opecoeloides polyfimbriatus n. sp., a new trematode from the lizard fish, Synodus foetens. Journal of Parasitology 33:231-233.
- 388 Read, C.P. 1948. Part I: Studies on the North American helminths of the genus Capillaria zeder, Nematoda. Part II: The life history and morphology of Rhopalias mercracanthus Chandler (Trematoda). M.A. Thesis, Rice University.
- 389 Read, C.P. 1948. Spirometra from Texas cats. Journal of Parasitology 34:71.

- 390 Read, C.P. 1948. A new capillarid nematode from the spiny dogfish, Squalus acanthius. *Journal of Parasitology* 34:163.
- 391 Read, C.P. 1949. Studies of North American helminths of the genus Capillaria Zeder, 1800 (Nematoda): I. Capillarids from mammals. *Journal of Parasitology* 35:223-230.
- 392 Read, C.P. 1949. Studies of North American helminths of the genus Capillaria Zeder, 1800 (Nematoda): III. Capillarids from the lower digestive tract of North American birds. *Journal of Parasitology* 35:240-249.
- 393 Read, C.P. 1952. Human sparganosis in South Texas. *Journal of Parasitology* 38:29-31.
- 394 Read, C.P. 1956. Trichuris from kangaroo rats. *Proceedings of the Helminthological Society of Washington* 23:2.
- 395 Read, C.P. 1956. Trichuris dipodomis n. sp. from Ord's kangaroo rats. *Proceedings of the Helminthological Society of Washington* 23:119.
- 396 Rigdon, R.H., and J.W. Hendricks. 1955. Myxosporidia in fish in waters emptying into the Gulf of Mexico. *Journal of Parasitology* 41:511-518.
- 397 Roberts, R.A. 1932. Myiasis in jack rabbits, Lepus californicus texanus. *Journal of Parasitology* 18:102-104.
- 398 Robinson, R.W., K.L. Kuttler, J.W. Thomas, and R.G. Marburger. 1967. Theileriosis in Texas white-tailed deer. *Journal of Wildlife Management* 31:455-459.
- 399 Robinson, R.W., K.L. Kuttler, H.R. Emerson, L.P. Jones, and R.G. Marburger. 1968. Blood parasites in Texas deer. *Transaction of the North American Wildlife Conference* 33:359-364.
- 400 Rodriguez, P.H. 1963. Ectoparasites of rabbits collected in Grant County, New Mexico. M.S. Thesis, University of New Mexico.
- 401 Russo, J.P. 1956. The Desert Bighorn Sheep in Arizona. Arizona Game and Fish Department, Phoenix, Arizona.
- 402 Ryckman, R.E. 1956. Parasitic and non-parasitic arthropods from bat caves in Texas and Mexico. *American Midland Naturalist* 56:186-190.
- 403 Samson, K.S., R.W. Allen, and G.A. Schad. 1964. Comparative pathogenicity in Ovis aries of homologous and heterologous strains of Haemonchus (Nematoda: Trichostrongylidae) from domestic and wild sheep. *Journal of Parasitology* 50:421-426.
- 404 Samson, K.S. 1968. Helminths of the black-tailed jackrabbits in New Mexico. *Bulletin of the Wildlife Disease Association* 4:130.
- 405 Samson, K.S., and B.R. Donaldson. 1968. Parasites of the javelina in New Mexico. *Bulletin of the Wildlife Disease Association* 4:131.
- 406 Samson, K.S. 1970. Experimental infection of domestic sheep with Trichostrongylus colubriformis from natural infections in black-tailed jackrabbits. *Journal of Parasitology* 56:1037.
- 407 Samuel, W.M., and D.O. Trainer. 1970. Amblyomma (Acarina: Ixodidae) on white-tailed deer, Odocoileus virginianus (Zimmermann), from South Texas with implications for theileriosis. *Journal of Medical Entomology* 7:567-574.
- 408 Samuel, W.M., and D.O. Trainer. 1970. Pulex porcinus Jordan and Rothschild, 1923 (Siphonaptera: Pulicidae), an occasional parasite of white-tailed deer from Welder Refuge in Southern Texas. *Journal of Wildlife Diseases* 6:182-183.
- 409 Samuel, W.M., and W.A. Low. 1970. Parasites of the collared peccary from Texas. *Journal of Wildlife Diseases* 6:16-23.
- 410 Samuel, W.M. 1972. Lipoptena mazamae Rondani, 1878 (Diptera: Hippoboscidae) on white-tailed deer in Southern Texas. *Journal of Medical Entomology* 9:104-106.
- 411 Schad, G.A. 1958. First report of Ornithodoros (Otobius) megnini from the pronghorn antelope, Antilocapra americana. *Journal of Parasitology* 44:514.
- 412 Schad, G.A., and R.W. Raught. 1958. Thelazia californiensis from a mule deer, Odocoileus hemionus crooki (Mearns, 1879) in New Mexico. *Journal of Parasitology* 44:483.
- 413 Schad, G.A. 1959. A revision of the North American species of the genus Skirjabinema (Nematoda: Oxyuridae). *Proceedings of the Helminthological Society of Washington* 26:138-147.
- 414 Schuck, B.R. 1945. A new locality for Trypanosoma cruzi in Arizona. *Journal of Parasitology* 31:151.
- 415 Schwartz, B., and J.E. Alicata. 1933. Description of two parasitic nematodes from the Texas peccary. *Publications of the United States National Museum* 82, Article 15.
- 416 Scott, J.A., and E. Blynn. 1952. Observations on infections in the cotton rat with Longistriata adunca and Strongylidae sigmodontis. *Journal of Parasitology* 38:88.
- 417 Scott, J.A., E.M. MacDonald, and L.J. Olson. 1957. Comparative experiments with a Florida strain of Litomosoides carlinii in eastern and Texas cotton rats. *Experimental Parasitology* 6:594-598.
- 418 Seamster, A., and L.H. Monaco. 1956. A new species of Rhamnocercinae. *American Midland Naturalist* 55:180-183.
- 419 Self, J.T. 1954. Parasites of the goldeye, Hiodon alosoides (Raf.) in Lake Texoma. *Journal of Parasitology* 40:386-389.
- 420 Self, J.T., and H.F. Timmons. 1955. The parasites of the river carpsucker (Carpoides carpio Raf.) in Lake Texoma. *Transactions of the American Microscopical Society* 74:350-352.
- 421 Self, J.T., and J.W. Campbell. 1956. A study

- of the helminth parasites of the buffalo fishes of Lake Texoma with a description of Lissorchis gullaris n. sp. (Trematoda: Lissorchiidae). Transactions of the American Microscopical Society 75:397-401.
- 422 Short, J.A., L.F. Mayberry, and J.R. Bristol. 1980. Eimeria chihuahuaensis sp.n. and other coccidia from Dipodomys spp. in El Paso County, Texas. Journal of Protozoology 27:361-364.
- 423 Simpson, D.T., and J.L. McGraw. 1979. Two new genera of Aspidogasteridae from Pogonis cromis (L.) from the Texas Gulf Coast. Southwestern Naturalist 24:557-561.
- 424 Slade, M.V. 1954. A study of Texas Gulf Coast parasites of livestock. M.S. Thesis, University of Houston.
- 425 Slagle, W.G. 1966. A survey of the helminths parasitic in Rana catesbiana Shaw and Rana pipiens Schreber from Brazos County, Texas and vicinity. M.S. Thesis, Texas A&M University.
- 426 Smit, F.G.A.M. 1958. A preliminary note on the occurrence of Pulex irritans L. and Pulex simulans Baker in North America. Journal of Parasitology 44:523-526.
- 427 Smith, A.J. 1908. Synopsis of studies in metazoan parasitology in McManus Laboratory of Pathology, University of Pennsylvania. University of Pennsylvania Medical Bulletin 20:267-282.
- 428 Smith, C.F. 1954. Four new species of cestodes of rodents from the high plains, central and southern rockies and notes on Catenotaenia dendritica. Journal of Parasitology 40:245-254.
- 429 Smith, J.P. 1967. Studies of transmission of a deer hemoprotozoan to domestic animals. M.S. Thesis, Texas A&M University.
- 430 Smith, J.P., and A. Seaton. 1981. Helminth infections of dogs in central Texas. Veterinary Medicine and Small Animal Clinician 76:1627-1629.
- 431 Smith, T., and F.L. Kilborne. 1893. Investigations into the nature, causation and prevention of Texas or southern cattle fever. United States Department of Agriculture Bureau of Animal Industry Bulletin 1.
- 432 Smith, W.N., and M.B. Chitwood. 1967. Trichospirura leptostoma gen. et sp. n. (Nematoda: Thelazoidea) from the pancreatic ducts of the white-eared marmoset, Callithrix jacchus. Journal of Parasitology 53:1270.
- 433 Sogandares-Bernal, F., and R.D. Lumsden. 1964. The heterophyid trematode Ascocotyle (A.) leighi Burton, 1956, from the hearts of certain poeciliid and cyprinodont fishes. Zeitschrift fur Parasitenkunde 24:3-12.
- 434 Sparks, A.K. 1951. Some helminth parasites of the largemouth bass in Texas. Transactions of the American Microscopical Society 70:351-358.
- 435 Specian, R.D., and J.E. Ubelaker. 1974. Two new species of Pharyngodon Diesing, 1861 (Nematoda: Oxyuridae) from lizards in West Texas. Proceedings of the Helminthological Society of Washington 41:46-51.
- 436 Specian, R.D., and J.E. Ubelaker. 1974. Parathelandros texanus n.sp. (Nematoda: Oxyuridae) from lizards in West Texas. Transactions of the American Microscopical Society 93:413-415.
- 437 Spindler, L.A., R.W. Allen, L.S. Diamond, and J.C. Lotze. 1958. Babesia in a white-tailed deer. Journal of Protozoology 5:(Suppl.)8.
- 438 Stewart, G.L., J.J. Reddington, and W.G. Smyth. 1979. Intestinal parasites of dogs from Tarrant County, Texas. Southwestern Veterinarian 32:29-32.
- 439 Stone, J.E., and D.B. Pence. 1977. Ectoparasites of the bobcat from West Texas. Journal of Parasitology 63:463.
- 440 Stone, J.E., and D.B. Pence. 1978. Ecology of helminth parasitism in the bobcat from West Texas. Journal of Parasitology 64:295-302.
- 441 Strandman, R.W. 1946. Atricholaelaps sigmodoni, a new species of mite parasitic on the cotton rat, and notes on the genera Atricholaelaps and Ischnolaelaps (Acarina: Laelaptidae). Journal of Parasitology 32:164-169.
- 442 Strandman, R.W. 1947. Atricholaelaps megaventralis, a new species of parasitic mite. Proceedings of the Entomological Society of Washington 49:112-114.
- 443 Strandman, R.W., and R.B. Eads. 1947. A new species of mite, Ichoronyssus dentipes (Acarina: Liponyssinae), from the cotton rat. Journal of Parasitology 33:51-56.
- 444 Strandman, R.W. 1948. The mesostigmatic nasal mites of birds. I. Two new genera from shore and marsh birds. Journal of Parasitology 34:505-514.
- 445 Strandman, R.W. 1949. The blood-sucking mites of the genus Haemolaelaps (Acarina: Laelaptidae) in the United States. Journal of Parasitology 35:325-352.
- 446 Strandman, R.W. 1951. The mesostigmatic nasal mites of birds. II. New and poorly known species of Rhinonyssidae. Journal of Parasitology 37:129-140.
- 447 Strandman, R.W., and O.E. Hunt. 1951. Two new species of Macronyssidae, with notes on some established genera (Acarina). Journal of Parasitology 37:460-470.
- 448 Strandman, R.W., and D.J. Eben. 1953. A survey of typhus in rats and rat ectoparasites in Galveston. Texas Reports on Biology and Medicine 11:144-151.
- 449 Strandman, R.W., and H.B. Morlan. 1953. A new species of Hirstionyssus and a key to the known species of the world. Texas Reports on Biology and Medicine 11:627-637.
- 450 Strandman, R.W. 1956. The mesostigmatic

- nasal mites of birds. IV. The species and hosts of the genus Rhinonyssus (Acarina, Rhinonyssidae). Proceedings of the Entomological Society of Washington 58:129-142.
- 451 Strandtman, R.W., and D.M. Allred. 1956. Mites of the genus Brevisgerna Keegan, 1949 (Acarina: Haemogamasidae). Journal of the Kansas Entomological Society 29:113-132.
- 452 Strandtman, R.W., and D.P. Fuman. 1956. A new species of mite, Paraneonyssus icteridius, from the nasal cavities of blackbirds (Acarina: Rhinonyssidae). Pan-Pacific Entomologist 32:167-173.
- 453 Strandtman, R.W. 1959. New records for Rhinonyssus himantopus and notes on other species of the genus. Journal of the Kansas Entomological Society 32:133-136.
- 454 Strandtman, R.W. 1961. Neonyssus triangulus n. sp., nasal mite (Acarina: Mesostigmata) from the white-winged dove (Aves: Columbiformes) and key to the species of the genus Neonyssus. Journal of Parasitology 47:323-328.
- 455 Strandtman, R.W. 1962. Nycteriglyphus bifolium n. sp., a new cavernicolous mite associated with bats (Chiroptera) (Acarina: Glycyphagidae). Acarologia 4:623-631.
- 456 Sundman, J.A., and E.O. Morrison. 1965. A new host record for Rhipicephalus sanguineus (Acarina: Ixodidae). Journal of the Kansas Entomological Society 39:354.
- 457 Swank, W.G. 1958. The mule deer in Arizona chaparral and an analysis of other important deer herds. Arizona Game and Fish Department Wildlife Bulletin 3.
- 458 Swickard, M., G.E. Haas, and R.P. Martin. 1971 (1972). Notes on small mammals infrequently recorded from the Jemez Mountains, New Mexico. Bulletin of the New Mexico Academy of Science 12:10-14.
- 459 Taylor, W.P. 1935. Ecology and life history of the porcupine (Erethizon epixanthum) as related to the forests of Arizona and the Southwestern United States. University of Arizona Bulletin 3:5-178.
- 460 Telford, S.R. 1970. Plasmodium chiricahuae sp. nov. from Arizona lizards. Journal of Protozoology 17:400-405.
- 461 Telford, S.R. 1978. Saurian malaria in Texas. Journal of Parasitology 64:553-554.
- 462 Thatcher, V.E., and J. Scott. 1962. The life cycle of Trichostrongylus sigmodontis Baylis, 1945, and the susceptibility of various laboratory animals to this nematode. Journal of Parasitology 48:558-561.
- 463 Thornton, J. 1972. Parasites of blackbuck antelope (Antilope cervicapra) in Texas and experimental transmission of gastrointestinal helminths from blackbuck to domestic ruminants. M.S. Thesis, Texas A&M University.
- 464 Thornton, J.E., R.R. Bell, and M.J. Reardon. 1974. Internal parasites of coyotes in Southern Texas. Journal of Wildlife Diseases 10:232-236.
- 465 Tibbitts, T. 1957. Description of a new genus, Periglischrus, from a bat, Marmoops megalophylla senicula Rehn, together with a key to the species of Periglischrus (Acarina, Spinturnicidae). Journal of the Kansas Entomological Society 30:13-19.
- 466 Tiner, J.D. 1946. Helminth parasites of skunks in Texas. Journal of Mammalogy 27:82-83.
- 467 Tiner, J.D. 1948. Rictularia dipodomis n. sp. (Nematoda: Thelaziidae) from the kangaroo rat Dipodomys sp. Journal of Parasitology 34:332-334.
- 468 Toepfer, E.W., L.N. Locke, and L.H. Blankenship. 1966. The occurrence of Trichomonas gallinae in white-winged doves in Arizona. Bulletin of the Wildlife Disease Association 2:13.
- 469 Tomlinson, T.H., C.S. Smith, and N.H. Fogg. 1948. A piroplasm, Babesia wrighti n. sp., from the rock squirrel (Citellus variegatus buckleyi). Journal of the National Malaria Society 7:254-258.
- 470 Toweill, D.E., and M.A. Price. 1976. Ectoparasites of ringtails collected from Kerr County, Texas. Southwestern Entomologist 1:20.
- 471 Trembly, H.L., and F.C. Bishopp. 1940. Distribution and hosts of some fleas of economic importance. Journal of Economic Entomology 33:701-703.
- 472 Trimmier, B.L. 1964. The life history of Haemogregarina crotali Laveran, 1902, in the prairie rattlesnake Crotalis viridis viridis Rafinesque, 1818. M.S. Thesis, New Mexico State University.
- 473 Tuff, D.W. 1963. The Mallophaga of the ciconiformes of America north of Mexico. Ph.D. Dissertation, Texas A&M University.
- 474 Tuff, D.W. 1965. A new species and record of the genus Ardeiphilus in North America (Mallophaga: Menoponida). Texas Journal of Science 17:110-112.
- 475 Tuff, D.W. 1975. Dirofilaria scapiceps (Leidy) from the eastern cotton-tail Sylvilagus floridanus alacer (Bangs) in Texas. Texas Journal of Science 26:607.
- 476 Turk, R.D. 1949. Liver flukes from the cat. Journal of the American Veterinary Medical Association 115:23.
- 477 Turk, R.D. 1950. Guinea-worm (Dracunculus insignis, Leidy, 1958) infection in a dog. Journal of the American Veterinary Medical Association 117:215-216.
- 478 Underwood, H.T., and N.O. Dronen. 1977. The molluscan intermediate hosts for species of Haematoloechus Looss 1899 (Digenea: Plagiorchiidae) from raniid frogs of Texas. Journal of Parasitology 63:122.

- 479 Underwood, H.T., and N.O. Dronen. 1984. Endohelminths of fishes from the upper San Marcos River, Texas. *Southwestern Naturalist* 29:377-385.
- 480 Van Cleave, H.J. 1925. A critical study of the Acanthocephala described and identified by Joseph Leidy. *Proceedings of the National Academy of Science Philadelphia* 76:279-334.
- 481 Van Cleave, H.J., and H.F. Timmons. 1952. An additional new species of the acanthocephalan genus Neoechinorhynchus. *Journal of Parasitology* 38:53-56.
- 482 Vincent, I. 1946. Studies on the endoparasites of the Texas horned lizard, Phrynosoma cornutum (Harlan). *Proceedings and Transactions of the Texas Academy of Science* 30:250-252.
- 483 Von Wicklen, J.H. 1946. The trematode genus Opecoeloides and related genera, with a description of Opecoeloides polynemi n. sp. *Journal of Parasitology* 32:156-163.
- 484 Vorhies, C.T., and W.P. Taylor. 1940. Life history and ecology of the white-throated woodrat, Neotoma albigula albigula Hartley. *Technical Bulletin 86, Arizona Agricultural Experiment Station*, pp. 379-459.
- 485 Voss, W.J. 1961. A survey of ixodorhynchid mites on snakes. M.S. Thesis, Texas Technological University.
- 486 Walker, K.A. 1973. Helminths of some Northern Arizona lizards. *Proceedings of the Helminthological Society of Washington* 40:168-169.
- 487 Wang, C.C. 1955. A study of Haemogregarina and Haemoproteus in the blood of Texas turtles. M.S. Thesis, Texas A&M University.
- 488 Wardle, W.J. 1980. On the life cycle stages of Proctoeces maculatus (Digenea: Fellodistomidae) in mussels and fishes from Galveston Bay, Texas. *Bulletin of Marine Science* 30:737-743.
- 489 Wardle, W.J. 1983. Two new non-ocellate trichocercous cercariae (Digenea: Fellodistomidae) from estuarine bivalved molluscs in Galveston Bay, Texas. *Contributions in Marine Science* 26:15-22.
- 490 Webb, J.E., J.C. Clark, and A.L. Adams. 1956. Studies on the lone star tick. I. Preliminary population studies at Camp Bullis, Texas. *Texas Journal of Science* 8:399-404.
- 491 Webster, J.D., and C.J. Addis. 1945. Helminths from the bob-white quail in Texas. *Journal of Parasitology* 31:286-287.
- 492 Webster, J.D. 1947. Helminths from the bob-white in Texas, with descriptions of two new cestodes. *Transactions of the American Microscopical Society* 66:339-343.
- 493 Webster, J.D. 1948. A new acanthocephalan from the bob-white. *Journal of Parasitology* 34:84-86.
- 494 Webster, J.D. 1948. A new acanthocephalan from the sanderling. *Transactions of the American Microscopical Society* 67:66-69.
- 495 Webster, J.D. 1948. Two cestodes from the nighthawk. *Journal of Parasitology* 34:93-95.
- 496 Webster, J.D. 1951. Additional notes on the helminth parasites of the bob-white in Texas. *Journal of Parasitology* 37:322.
- 497 Weir, W.L., L.F. Mayberry, H.G. Kinzer, and P.R. Turner. 1983. Parasites of fishes in the Gila River drainage in Southwestern New Mexico. *Journal of Wildlife Diseases* 19:59-60.
- 498 Wheeler, J.R., D.L. Forcum, and C.D. Rael. 1970. Two records of bird fleas from New Mexico. *Southwestern Naturalist* 14:371.
- 499 Wilkinson, J.N., A.G. Canaris, and D. Broderson. 1977. Parasites of waterfowl from Southwest Texas: II. The shoveler, Anas clypeata. *Journal of Wildlife Diseases* 13:62-63.
- 500 Williams, L.A. 1949. A study of fleas of rodents in the Albuquerque, New Mexico area. M.A. Thesis, University of New Mexico.
- 501 Wilson, G.I. 1969. Some parasites of elk in New Mexico. *Bulletin of the Wildlife Disease Association* 5:23-24.
- 502 Wilson, G.I., and K.S. Samson. 1971. The incidence of fascioliasis in sheep and cattle of the Southwest with observations on snail vectors. *Proceedings of the Helminthological Society of Washington* 38:52-56.
- 503 Wilson, N., and G.B. Oliver. 1978. Noteworthy records of two ectoparasites (Cimicidae and Hippoboscidae) from the turkey vulture in Texas. *Southwestern Naturalist* 23:305-307.
- 504 Wiseman, J.S. 1959. The genera of Mallophaga of North America north of Mexico with special reference to Texas species. Ph.D. Dissertation, Texas A&M University.
- 505 Wood, D.E. 1963. Some intestinal helminths of Ambystoma tigrinum. M.S. Thesis, New Mexico State University.
- 506 Wood, S.F., and M.C. Herman. 1943. The occurrence of blood parasites in birds from Southwestern United States. *Journal of Parasitology* 29:187.
- 507 Wood, S.F. 1952. Mammal blood parasite records from Southwestern United States and Mexico. *Journal of Parasitology* 38:85-86.
- 508 Word, B.A. 1967. A study of external and internal parasites of beef cattle in the highland hereford area of Texas. M.A. Thesis, Sul Ross University.
- 509 Yeager, P.A. 1964. Incidence of human and swine trichinosis in a limited North-central Texas area. M.A. Thesis, Texas Christian University.

THESES AND DISSERTATIONS NOT REFERENCED

- Allison, T.C. 1964. Gill trematodes of certain Centrarchidae from the Little and Navasota Rivers. M.S. Thesis, Texas A & M University.
- Amin, O.M. 1968. Helminth fauna of suckers (Catostomidae) of the Gila River System, Arizona. Ph.D. Dissertation, Arizona State University.
- Anderson, R.D. 1969. The distribution and relative abundance of Nematopsis spp., as found in Crassostrea virginica (Gmelin) in the Galveston Bay area. M.S. Thesis, Texas A & M University.
- Anderson, R.R. 1968. Helminth and copepod parasites of the largemouth bass, Micropterus salmoides (Lacepede) from Lake Whitney, Bosque County, Texas. M.A. Thesis, Baylor University.
- Armstrong, H.W. 1969. A survey of the helminth parasites of certain Cyprinodontidae and Poeciliidae of the Texas coast. M.S. Thesis, Texas A & M University. Bedinger, C.A. 1967. Helminth parasites of East Texas fishes. M.S. Thesis, Sam Houston State University.
- Behrens, C.B. 1970. A survey of platyhelminth and acanthocephalan parasites in Sturnus vulgaris L. of Nacodoches County, Texas with emphasis on Cestoda and Acanthocephala. M.S. Thesis, Stephen F. Austin University.
- Block, E.F. 1968. Parasites of the cricket frog, Arcis crepitans. M.S. Thesis, North Texas State University.
- Bloebaum, A.P. 1971. The biology of Azygia angusticauda (Stafford, 1904) Manter, 1926 (Trematoda: Azygiidae). Ph.D. Dissertation, New Mexico State University.
- Boring, G.D. 1964. The infectivity of Bunostomum phlebotomum to sheep, rabbit, and guinea pigs. M.S. Thesis, Texas A & M University.
- Bornstein, A.C. 1974. Seasonal variation in metagon parasites of Dipodomys ordi in the northwest quadrant of the Chihuahuan Desert. M.S. Thesis, University of Texas at El Paso.
- Bowen, R.C. 1962. A comparison of intestinal protozoa among Peromyscus and other small rodents of Bernalillo County, New Mexico. M.S. Thesis University of New Mexico. Brooks, D.L. 1959. The nasal mites of some West Texas fly catchers. M.S. Thesis, Texas Technological University.
- Brown, E.E. 1940. Biology of two freshwater trematodes: Analocreadium armatum (MacCallum, 1895) and Microcreadium parvum (Simer, 1929). M.S. Thesis, Texas A & M University.
- Buckner, C.B. 1972. An intestinal cestode of the Colorado River squawfish, Ptychocheilus lucius Girard. M.S. Thesis, Arizona State University.
- Bueno, R. 1979. A survey for coccidia in migratory waterfowl along the Central Flyway and cross-transmission experiments to the pekin duck, Anas platyrhynchos domestica. M.S. Thesis, University of Texas at El Paso.
- Burke, E.P. 1967. Bodo alternata sp. nov.: Studies in physiological ecology and natural history. M.S. Thesis, Texas A & M University.
- Caskey, M.W. 1922. Morphology of Gigantorhynchus hirudinaceus. M.S. Thesis, Baylor University. Chan, B.S. 1962. Parasites in the digestive tract of the pocket gopher (Geomys bursarius) in the Waco area. Masters Thesis, Baylor University.
- Corkern, C.C. 1970. Investigations on helminths of the brown shrimp, Penaeus aztecus Ives, from Galveston Bay, Texas. M.S. Thesis, Texas A & M University.
- Crossley, D.A. 1951. Nasal mites of some columbiform birds. M.S. Thesis, Texas Technological University.
- Curfman, B.E. 1965. Parasitic helminths of Natrix erythrogaster transversa (Hallowell), the blotched water snake in the Waco area. M.S. Thesis, Baylor University.
- Davis, J.W. 1968. Infestation of parasitic keds (Diptera: Hippoboscidae) on white-tailed deer in East Texas. M.F. (Forestry) Thesis, Stephen F. Austin State University.
- Denton, J.F. 1941. Studies on the morphology, taxonomy and life histories of trematodes of the subfamily Dicrocoelinae Looss, 1899. Ph.D. Dissertation, Rice University.
- Dooley, T.J. 1974. Bats of El Paso County, Texas, with notes on habitat, behavior, and ectoparasites. M.S. Thesis, University of Texas at El Paso.
- Duckenfield, H. 1943. A mite of the genus Galumna and its relation to anoplocephalid cestodes. M.S. Thesis, Texas A & M University.
- Dunagen, T.T. 1955. Studies on cercariae in Brazos County, Texas. M.S. Thesis, Texas A & M University. Eads, R.B. 1949. A taxonomic study of the Siphonaptera, or fleas, of Texas and an evaluation of their importance in public health. Ph.D. Dissertation, Texas A & M University.
- El Bihari, S. 1968. Feather mites (Acarina: Analgesoidea) of some Texan and Mexican falconiform and strigiform birds. M.S. Thesis, Texas A & M University.
- Eller, R.R. 1962. The physiological specificity of the spiny rat mite, Echinolaelaps echidninus, for Rattus norvegicus. M.S. Thesis, Texas Technological University.
- Epstein, R.A. 1972. Larval trematodes of marine gastropods of Galveston Island, Texas. M.S. Thesis, Texas A & M University.
- Esslinger, J.H. 1958. Host-parasite relations in the screw-worm, Callitroga hominivorax. Ph.D. Dissertation, Rice University.
- Farias, J.D. 1983. Parasites of the Mexican duck, Anas platyrhynchos diazi, from Southwestern United States and Mexico. M.S. Thesis, University of Texas at El Paso.
- Gentner, H.W. 1968. Host-parasite relationships of

- cottontail rabbits (*Sylvilagus floridanus*) and tapeworms of the genera *Cittotaenia* and *Raillietina*. M.S. Thesis, Texas A & M University.
- George, J.E. 1960. The nasal mites of the genus *Ptilonyssus* occurring in some North American passeriforme birds. M.S. Thesis, Texas Technological University.
- Gilmore, R.E. 1955. The helminths parasitic in the lizard *Cnemidophorus grahami*, and a description of a new cestode from the lizard *Holbrookia texana*. M.S. Thesis, New Mexico State University.
- Harwood, P.D. 1931. The helminths parasitic in the amphibia and reptilia of Houston, Texas and vicinity. Ph.D. Dissertation, Rice University.
- Hathaway, R.P. 1966. A survey of intestinal helminths occurring in dogs from Albuquerque, M.S. Thesis, University of New Mexico.
- Hays, L.M. 1950. Comparison of Mallophaga on related species of birds. M.S. Thesis, Texas Technological University.
- Henson, R.N. 1966. A survey of cestodes in some shallow water elasmobranchs of Texas. M.S. Thesis, Texas A & M University.
- Hinojos, J.G. 1984. Metazoan parasites of the black-necked stilt, *Himantopus mexicanus*, from El Paso, Texas. M.S. Thesis, University of Texas at El Paso.
- Hollis, P.D. 1966. A survey of platyhelminth parasites in *Rana catesbiana* of Central-east Texas and notes on other parasites. M.S. Thesis, Stehen F. Austin State University.
- Howell, F.G. 1970. Responses of *Argas cooleyi* (Acarina: Argasidae) to relative humidity. M.S. Thesis, Texas Technological University.
- Hughins, E.J. 1949. A survey of the helminths and ectoparasites of rats, *Rattus rattus frugivorus* Rafinesque and *Sigmodon hispidus hispidus* Say and Ord, in Brazos County, Texas. M.S. Thesis, Texas A & M University.
- Jakeman, L.R. 1960. The internal anatomy of the spiny rat mite, *Echinolaelaps echidninus*. M.S. Thesis, Texas Technological University.
- Jameson, D.K. 1956. A survey of five species of bats and their parasites. M.S. Thesis, Texas Technological University.
- Joy, S.K. 1971. Relationships between the cotton rat, *Sigmodon hispidus texianus* (Audubon and Bachman) and its ectoparasites. M.S. Thesis, Texas A & M University.
- Koontz, A.C. 1965. The helminth fauna of the raccoon, *Procyon lotor* (Linnaeus), in South and Central Texas. M.A. Thesis, Baylor University.
- Koratha, K.J. 1953. A study of Monogenetic trematodes from marine fishes of Port Aransas, Texas, with a general review of the group. M.S. Thesis, Texas A & M University.
- Landwer, V.E. 1956. Laboratory observations on feeding, longevity and sensory behavior of *Echinolaelaps echnidninus*. M.S. Thesis, Texas Technological University.
- Lawrence, J.L. 1965. Parasites of five species of fish from Benbrook Lake, Tarrant County, Texas. M.A. Thesis, Texas Christian University.
- Laws, W.L. 1966. Photactic behavior of the spriny rat mite, *Echinolaelaps echidninus*. M.S. Thesis, Texas Technological University.
- Lee, D. 1966. The neotropical mite, Genus *Gigantolaelaps*, Fonseca, 1939. M.S. Thesis, Texas Technological University.
- Lee, T. 1971. Host-parasite relationships of the endoparasitic helminths of the eastern meadowlarks (*Sturnella magna magna* L.) in McLennan County, Texas. M.A. Thesis, Baylor University.
- Lushbaugh, W.B. 1969. Some biochemical aspects of the systematics of *Proterometra* spp. (Trematoda: Azygiidae). M.S. Thesis, New Mexico State University.
- Martin, M.L. 1933. Life history of the pigeon louse (*Columbicola columbae* Linnaeus). M.A. Thesis, Rice University.
- Matthes, H.C. 1931. A contribution to the study of bubonic plaque and the seasonal abundance of *Rhopalopsyllus sigmodoni* (Siphonaptera). M.A. Thesis, Rice University.
- Matthes, H.C. 1934. A study of the seasonal distribution of *Anopheles* in Houston, Texas. Ph.D. Dissertation, Rice University.
- McGraw, J.L. 1964. Some internal helminth parasites of certain Centrarchidae of the Navasota and Little Rivers of Texas. M.S. Thesis, Texas A & M University.
- McGuire, S. 1962. Two new furcocysticerous cercariae from *Goniobasis livescens* (Menke). M.S. Thesis, New Mexico State University.
- Meadow, J. 1968. Parasite community structure in three rodents, *Sigmodon*, *Perognathus*, and *Baiomys* in Central Texas. M.A. Thesis, Baylor University.
- Miller, W.L. 1940. A survey of hookworm infection in Brazos County, Texas. M.S. Thesis, Texas A & M University.
- Montgomery, D.F. 1966. A taxonomic study of the lizard mites (Pterygosomidae) occurring in the Gulf of California area. M.S. Thesis, Texas Technological University.
- Moore, D.V. 1942. Studies on the life history and development of certain Acanthocephala of the order Archiacanthocephala. Ph.D. Dissertation, Rice University.
- Norris, J.B. 1950. Studies on gametogenesis of the digenetic trematode, *Proterometra dickermani* Anderson. 1. Spermatogenesis. M.S. Thesis, New Mexico State University.
- Philpott, C.W. 1958. The reproductive response of the spiny rat mite, *Echinolaelaps echidninus*, to various bloods and blood fractions. M.S. Thesis, Texas Technological University.
- Pittard, S.K. 1970. Comparative external morphology of the life stages of *Cryptocellus*

- pelaezi (Arachnida: Ricinulei). M.S. Thesis, Texas Technological University.
- Ray, J.P. 1970. Developmental cycle for a new marine chytrid, Phlyctochytridium mackini sp. nov., (Chytridiales: Phlyctochytridiaceae) from the Gulf of Mexico. M.S. Thesis, Texas A & M University.
- Read, C.P. 1948. Part I: Studies on the North American helminths of the genus Capillaria zedar, Nematoda. Part II: The life history and morphology of Rhopalias mercracanthus Chandler (Trematoda). M.A. Thesis, Rice University.
- Riggs, V.C. 1956. Observations on the food preferences and reproductive potentiality of the spiny rat mite, Echinolaelaps echidninus. M.S. Thesis, Texas Technological University.
- Rodriguez, P.H. 1963. Ectoparasites of rabbits collected in Grant County, New Mexico. M.S. Thesis, University of New Mexico.
- Samson, K.S. 1966. On the synonymy of Nematodirus oiratianus Raevskaia, 1929, and Nematodirus lanceolatus Ault, 1944, nematode parasites of ruminants. M.S. Thesis, New Mexico State University.
- Schicht, F.G. 1963. The relationship of Hexamita sp. to mortalities of the Texas commercial oyster, Crassostrea virginica Gmelin, maintained under laboratory conditions. M.S. Thesis, Texas A & M University.
- Shaikh, H. 1963. Larval survival in mixed infections of Trichstrongylidae in calves. M.S. Thesis, Texas A & M University.
- Shoemaker, R.R. 1963. The nasal mites of the genus Paraneonyssus occurring in some North American passeriform birds. M.S. Thesis, Texas Technological University.
- Sileo, L. 1970. The incidence of virulence of Trichomonas gallinae (Rivolta) in mourning dove (Zenaidura macroura) (Linnaeus) populations in Southern Arizona. Ph.D. Dissertation, University of Arizona.
- Slade, M.V. 1954. A study of Texas Gulf Coast parasites of livestock. M.S. Thesis, University of Houston.
- Slagle, W.G. 1966. A survey of the helminths parasitic in Rana catesbiana Shaw and Rana pipiens Schreber from Brazos County, Texas and vicinity. M.S. Thesis, Texas A & M University.
- Smith, J.P. 1967. Studies of transmission of a deer hemoprotozoan to domestic animals. M.S. Thesis, Texas A & M University.
- Sparks, A.K. 1957. A study of the geographic distribution of digenetic trematodes of shallow-water fishes of the Gulf of Mexico. Ph.D. Dissertation, Texas A & M University.
- Strickland, R. 1972. The susceptibility of three species of mosquitoes to Reesimermis nielsenii Tsai and Grundmann, 1969 (Nematoda: Mermithidae). M.S. Thesis, Lamar University.
- Thornton, J.E. 1972. Parasites of blackbuck antelope (Antelope cervicapra) in Texas and experimental transmission of gastrointestinal helminths from blackbuck to domestic ruminants. M.S. Thesis, Texas A & M University.
- Trimmier, B. 1964. The life history of Haemogregarina crotali Laveran, 1902, in the prairie rattlesnake Crotalis viridis viridis Rafinesque, 1818. M.S. Thesis, New Mexico State University.
- Tringas, P. 1972. A taxonomic and biological study of nematode associates of five species of insects. M.S. Thesis, Lamar University.
- Tuff, D.W. 1963. The Mallophaga of the ciconiformes of America north of Mexico. Ph.D. Dissertation, Texas A & M University.
- Voss, W.J. 1961. A survey of ixodorhynchid mites on snakes. M.S. Thesis, Texas Technological University.
- Wang, C.C. 1955. A study of Haemogregarina and Haemoproteus in the blood of Texas turtles. M.S. Thesis, Texas A & M University.
- Weir, W. 1979. Parasites of fish in the Gila River drainage, New Mexico. M.S. Thesis, New Mexico State University.
- Williams, L.A. 1949. A study of fleas of rodents in the Albuquerque, New Mexico area. M.A. Thesis, University of New Mexico. Wiseman, J.S. 1959. The genera of Mallophaga of North America north of Mexico with special reference to Texas species. Ph.D. Dissertation, Texas A & M University.
- Wood, D.E. 1963. Some intestinal helminths of Ambystoma tigrinum. M.S. Thesis, New Mexico State University.
- Word, B.A. 1967. A study of external and internal parasites of beef cattle in the Highland Hereford area of Texas. M.A. Thesis, Sul Ross University.
- Yeager, P.A. 1964. Incidence of human and swine trichinosis in a limited North-central Texas area. M.A. Thesis, Texas Christian University.
- Yilmaz, S.M. 1971. Studies on the biology of Toxoplasma gondii (Nicolle and Manceaux, 1908), an important protozoan parasite of man and many animals. Ph.D. Dissertation, Texas A & M University.

ACKNOWLEDGEMENTS

We would like to thank Dr. A.H. Harris and Dr. R.G. Webb for their help in updating the taxonomic names, the numerous students who have helped through the years in the preparation of this bibliography and to express deep appreciation to Ms. Elena M. Cook for typing the manuscript.