University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Manure Matters (newsletter)

Biological Systems Engineering

1-1-2001

Manure Matters, Volume 7, Number 4

Raymond Ward Ward Laboratories, Inc.

Follow this and additional works at: http://digitalcommons.unl.edu/manurematters



Part of the Agronomy and Crop Sciences Commons

Ward, Raymond, "Manure Matters, Volume 7, Number 4" (2001). Manure Matters (newsletter). Paper 4. http://digitalcommons.unl.edu/manurematters/4

This Article is brought to you for free and open access by the Biological Systems Engineering at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Manure Matters (newsletter) by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln. UNL's Livestock
Environmental
Issues
Committee
Includes
representation from
UNL, Nebraska
Department of
Environmental
Quality, Natural
Resources
Conservation Service,
Natural Resources
Districts, Center for
Rural Affairs,
Nebraska Cattlemen,
USDA Ag Research
Services, and
Nebraska Pork
Producers

Contact: Chris Henry 217 LW Chase Hall University of NE Lincoln, NE 68583 (402) 472-6529 chenry@.unl.edu



How to Read a Manure Sample Report

Raymond C. Ward Ward Laboratories, Inc.

Correctly sampling manure can be a challenge. Try to obtain a representative sample and get the sample to the lab as close as possible to the time when manure is going to be applied. It is preferable to obtain sample results before land application so that appropriate application rates can be determined.

Slurry samples should be taken at the time the slurry is being applied, so that you have a good mix of the manure storage. The amount of material is usually recorded in gallons per acre and once the analysis is received, the amount of nutrients can be calculated. If you want to take a slurry sample before application, the manure storage would have to be mixed very well. It is difficult to get a good representative sample from a manure storage by just reaching in and grabbing a sample.

The container, preferably plastic, of 1/2 to 1 pint size can be used for the sample. Fill the bottle completely full so there is no head space. Seal the lid

with tape to ensure the lid does not come off during shipment. The sample should be chilled and preferably be shipped on ice if possible. We prefer a cooled sample more than a frozen sample. For those individuals close to the laboratory, hand delivered is best. Samples are considered a slurry any time they are fluid. If the sample is non-fluid, then we would call it a manure sample. Slurry samples are analyzed on an as is basis, without drying.

Manure samples are handled differently in the laboratory, because they are dried and ground before an analysis is done. Manure samples can be taken in a random method across the feedyard. If sampling stockpiled manure, use a soil probe and take several random samples from the pile for a composite sample. Be sure to get into the pile past the dry outside layer before collecting the sample for analysis. Poultry manure could be sampled as it is delivered from the houses into a stockpile or to a spreader, depending on the

UNL's Livestock Environmental Issues Committee Includes representation from UNL, Nebraska Department of Environmental Quality, Natural Resources Conservation Service Natural Affairs, Nebraska Cattlemen, USDA Ag Research Services, and Nebraska Pork Producers

Contact: Chris Henry 217 LW Chase Hall University of NE Lincoln, NE 68583 (402) 472-6529 chenry@.unl.edu operation. Again, hand grabbing several random samples to make a composite sample is the ideal way to collect these types of samples.

When taking a random sample, the preferred number of sub-samples is fifteen. Often we say take 10-15 sub-samples to make one composite sample. For manure samples, they should be sealed in zip-lock bags to retain moisture so that the moisture content can be determined to accurately calculate total nutrients applied.

In our laboratory, manure samples and slurry samples are analyzed the day after receipt. Ideally, it would be best to send the samples to the laboratory early in the week so that the samples do not sit over the weekend before analysis.

As mentioned above, figure on 1/2 to 1 pint of slurry for analysis, and the same could be said for the manure samples. About a pint of material is needed, but it is more important to get a good composite sample. The slurry analysis is reported on an as is basis, and the amount of the dry matter content is also reported in a report. The total amount of nutrients is reported in lbs/1000 gal and in lbs/acre inch. An estimated availability the first year is also reported. However, for someone that is using slurry on the same land year after year, they should calculate application based on the total nutrients in the slurry and not just what is available the first year. The first year calculations are important for the first year of slurry use, but in future years, the total should be used.

The manure samples are reported on a

dry basis, and then calculated to lbs/ton on an "as is" basis. The amount available the first year per ton is also shown on the report. Again, when manure is applied for the first time, you should use the amount available the first year. In future years, it is best to look at the total analysis to determine the actual amount of plant nutrients being applied so that the land does not become excessively high in nutrients.

Below is a description and explanation of what is shown in a manure and slurry sample report.

- * Organic Nitrogen is the total amount of nitrogen less ammonium and nitrate. The availability of organic nitrogen is less than 100% and dependent on many factors, but the organic nitrogen in slurries would be released at about 50% and about 25% from manures.
- * **Ammonium** is 100% available if incorporated within 2 days. Ammonium loss could be 90% if left on dry surfaces for more than 15 days.

Nitrate is 100% available.

Total Nitrogen is the sum of organic nitrogen, ammonium-N and nitrate-N.

Phosphorus Total phosphorus is reported as P_2O_5 . Approximately 50% would be available the first year in slurries, and 35% in manures.

Potassium is reported as K₂O. All potassium is in the mineral form, so it is 100% available the first year.

Sulfur is usually in organic form and it would be about 50% available the

UNL's Livestock Environmental Issues Committee Includes representation from UNL, Nebraska Department of Environmental Quality, Natural Resources Conservation Service, Natural Resources Districts, Center for Rural Affairs, Nebraska Cattlemen, USDA Ag Research Services, and Nebraska Pork Producers

Contact: Chris Henry 217 LW Chase Hall University of NE Lincoln, NE 68583 (402) 472-6529 chenry@.unl.edu first year in slurries and 25% available in manures.

Calcium is mainly in an organic form and would be available at about 50% in the first year.

Magnesium would be similar to calcium.

Sodium is a non-plant nutrient that can cause problems in soils. Sodium disperses the clays and slows water infiltration. The sodium content may become a problem when more than 500 lbs of sodium is applied.

Zinc, Iron, Manganese & Copper are available at about 50% the first year.

Soluble Salts is a reading that indicates the total amount of salts in the slurry or manure. Soluble salts are actually plant nutrients. When more than 5000 lbs of Soluble Salts are applied, a salinity condition could develop.

pH is acidity or alkalinity of the manure or slurry and is usually slightly alkaline.

Dry Matter is the amount of solid material in the slurry or manure.

* More information on nutrient availability factors and their use can be found at:

http://www.ianr.unl.edu/pubs/wastem gt/g1335.htm

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperative with the U.S. Department of Agriculture. Kenneth R. Bolen, Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources. University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska Lincoln and the United States Department of Agriculture.