

*Chemical and Biomolecular Engineering Research and
Publications*

Papers in Molecular Chemistry

University of Nebraska - Lincoln

Year 1992

Kinetic Analysis of Competing
Intramolecular and Intermolecular
Polymerization Reactions

Hossein Nouredini*

Delmar C. Timm[†]

*Department of Chemical Engineering, University of Nebraska-Lincoln,
hnoured@unlnotes.unl.edu

[†]Department of Chemical Engineering, University of Nebraska-Lincoln,
dtimm@unlserve.unl.edu

This paper is posted at DigitalCommons@University of Nebraska - Lincoln.

<http://digitalcommons.unl.edu/chemengmolecular/5>

H. Nouredini and D. C. Timm, *Kinetic Analysis of Competing Intramolecular and Intermolecular Polymerization Reactions*, *Macromolecules*; 1992; 25(6); 1725-1730.

Kinetic Analysis of Competing Intramolecular and Intermolecular Polymerization Reactions

H. Nouredini and D. C. Timm

Department of Chemical Engineering, University of Nebraska, Lincoln, Nebraska

Received August 21, 1990; Revised Manuscript Received December 8, 1991

Macromolecules; 1992; 25(6); 1725-1730.

Copyright © 1992 American Chemical Society.

The American Chemical Society allows the posting of only the title, abstract, tables, and figures from articles appearing in the *Macromolecules*.


This article is available at published online at:

<http://pubs.acs.org/cgi-bin/archive.cgi/mamobx/1992/25/i06/pdf/ma00032a016.pdf>

ABSTRACT: Kinetic reaction theory was used to model a step-growth, the most polymerization of a monomer of f functional groups. Intermolecular additions were represented by second-order and intramolecular reactions were expressed by first-order rate expressions. All functional groups were assumed to react with equal reactivity. Independent variables are degree of polymerization i , extent of cross-linking j , and conversion ρ . The normalized rate constant for intramolecular reactions is c . The solution for the normalized population density distribution is

$$P_{ij} = \frac{f(f-1)!(f-2)!/2!}{2^{j-1}(f-2)^{j-1} + 1} \frac{c^j \rho^{f-j} [(1-\rho + c)(1-\rho)]^{(f-1)j-1} \rho^{(f-1)j}}{(1+c)^{f-1}}$$

subject to $0 \leq i \leq (f-2)j/2 - 1$ and $c \geq 0$. Formulas for the number-, mass-, and cross-link-average molecular weights were derived.






Figure 2. Calculated vs actual degree of cure for a 2 K/min scan by DSC analysis.