

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Great Plains Wildlife Damage Control Workshop
Proceedings

Wildlife Damage Management, Internet Center for

April 1991

RELATIVE EFFECTIVENESS OF REPELLENTS FOR REDUCING DAMAGE BY ELK

William F. Andelt
Colorado State University

Dan L. Baker
Colorado Division of Wildlife,

Kenneth P. Burnham
Colorado State University

Follow this and additional works at: <http://digitalcommons.unl.edu/gpwdcwp>



Part of the [Environmental Health and Protection Commons](#)

Andelt, William F.; Baker, Dan L.; and Burnham, Kenneth P., "RELATIVE EFFECTIVENESS OF REPELLENTS FOR REDUCING DAMAGE BY ELK" (1991). *Great Plains Wildlife Damage Control Workshop Proceedings*. 5.
<http://digitalcommons.unl.edu/gpwdcwp/5>

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Great Plains Wildlife Damage Control Workshop Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

RELATIVE EFFECTIVENESS OF REPELLENTS FOR REDUCING DAMAGE BY ELK

WILLIAM F. ANDELT, *Department of Fishery and Wildlife Biology, Colorado State University, Fort Collins, CO 80523*

DAN L. BAKER, *Colorado Division of Wildlife, 317 W. Prospect, Fort Collins, CO 80526*

KENNETH P. BURNHAM, *Colorado Cooperative Fish and Wildlife Research Unit, Colorado State University, Fort Collins, CO 80523*

Proceedings 10th Great Plains Wildlife Damage Conference (S.E. Hygnstrom, R.M. Case, and R.J. Johnson, eds.) Published at the University of Nebraska-Lincoln, 1991.

Abstract: We tested the repellency of chicken eggs, MGK' Big Game Repellent (BGR), coyote (*Canis latrans*) urine, thiram, Hinder', Hot Sauce Animal Repellent' (Hot Sauce), and Ropey on captive tame elk (*Cervus elaphus*) in Colorado from December 1990 to February 1991. Coyote urine and BGR performed better than the other repellents for deterring elk from feeding on cubed alfalfa rations. Consumption of cubed rations treated with odor repellents increased from Day 1 through Day 5 of the trial, but consumption

of thiram-treated rations decreased. During the second phase of the study, hungry elk consumed significantly more apple twigs that were treated with coyote urine, Hot Sauce, and thiram than did elk that were full fed. The repellency of Hot Sauce increased significantly in the second phase when the concentration was increased 100 times above the labeled recommendations and was the only repellent treatment tested that consistently deterred hungry elk from feeding on apple twigs.