

January 2006

Nebraska Vine Lines, Volume IX, Issue 1. January/ February 2006.

Follow this and additional works at: <http://digitalcommons.unl.edu/viticulturepubs>



Part of the [Fruit Science Commons](#)

"Nebraska Vine Lines, Volume IX, Issue 1. January/February 2006." (2006). *Viticulture Publications and Research*. 5.
<http://digitalcommons.unl.edu/viticulturepubs/5>

This Article is brought to you for free and open access by the Viticulture Program at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Viticulture Publications and Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

UNIVERSITY OF
Nebraska
Lincoln



January/February

Volume IX, Issue 1

Nebraska Vine Lines

*Editors: Dr. Paul Read, Professor of Horticulture & Viticulture
Donna Michel, University of Nebraska Viticulture Program*

Be Sure To Put The 9th Annual Forum on Your Calendar

March 3 and 4, 2006. Outstanding speakers, growers roundtables, local speakers and an outstanding trade show await you at the 9th Annual Nebraska Winery and Grape Growers Forum and Trade Show to be held at the Kearney Holiday Inn March 3 and 4, 2006. Read more about the 2 key-note speakers in this issue.

For further reading of Anna Katharine's treatment of 'Frontenac', see article on p2.

See page 5 for a note and greetings from Paul Read.



Anna Katharine Mansfield
Senior Research Fellow
Horticultural Science
Department of Horticultural Science
University of Minnesota
Enology Lab

Enologist

A native of North Carolina, [Anna Katharine Mansfield](#) became the first Enology Project Leader at the University of Minnesota in 2001. She received a MS in Food Science from Virginia Tech, where she researched wine aroma and flavor precursors. Anna Katharine oversees the enological evaluation and optimization of new cultivars and is responsible for wine aroma/flavor characterization and enology outreach.

Terry Bates

Dr. [Terry Bates](#) is working on root biology related projects at the Vineyard Lab. From low-tech root trenches to high-tech minirhizotrons, researchers are exploring the world of root growth, root distribution, and resource storage in Concord grapevines.

Refining the Concord Diet -- Terry Bates is studying the effect of soil pH and nutrient availability on Concord root growth and nutrient uptake.

Viticulture Research Associate
Cornell University
New York State Agricultural
Experiment Station



Treessentials Company

Corporate Offices
60 E. Plato Boulevard
Suite 130

St. Paul, MN 55107

800-634-2843



Visit our WEB site: MidwestVineyardSupply.com

**Midwest
Vineyard
Supply, Inc.**

2300 S. Twin Bridge Road
Decatur, Illinois 62521

Phone: 217-864-9896
FAX: 217-422-7793

**SERVING ALL YOUR VINEYARD AND ORCHARD
SUPPLY NEEDS!**

EMAIL: Sales@Midwestvineyardsupply.com

REAMS SPRINKLER SUPPLY

Irrigation Experts For Over 30 Years

We have been working in the irrigation field for over 30 years. Our background includes low-volume irrigation (drip), residential/commercial, sports-fields/golf courses. We have fully stocked locations in Lincoln, Omaha, Grand Island, and Sioux City. Our CAD design staff includes an *Irrigation Association* Certified Irrigation Designer (C.I.D.) to provide professional design. We have the staff and the products to allow us the opportunity to EARN your business.

Let us Use Our Experience to Assist You in Your needs:

OMAHA 3625 s. 149 TH St Omaha, NE 68144 877-838-1600 402-896-1600	LINCOLN 6001 s. 57 TH Lincoln, NE 68516 800-798-2473 402-423-0120	GRAND ISLAND 453 Industrial Ln Grand Island, NE 68803 888-381-6677 308-381-6000	SIOUX CITY 1010 Gordon Dr Sioux City, IA 51101 800-373-2009 712-258-8225
---	---	--	--

www.reams.com

Making Wine With Frontenac By: Anna Katharine Mansfield UM Enology Project Leader

As the Frontenac plantings introduced to northern growing regions in the past 10 years mature, an ever growing group of winemakers are faced with the unusual winemaking parameters of this new cultivar. Traits derived from Frontenac's V. riparia ancestry help it grow and thrive in cold climates, but also make it a very different beast than its V. vinifera cousins. Consequently, putting it through the paces that would make a good Cabernet Sauvignon, Merlot or even Marechal Foch often result in an inky, unbalanced wine.

Handling Frontenac isn't difficult; it's just different. Fortunately, these differences can be manipulated to produce a variety of wine styles, making the grape more flexible than many. This allows the production of several marketable styles from one crop, and gives the winemaker options in years when the growing season has been less than optimal. Modifications in winemaking method are necessary to take advantage of Frontenac's four variations from traditional wine grapes: it is more highly colored than most V. vinifera, it has higher soluble solids and acid content, and it tends to have low tannin.

Like the fruit of its V. riparia ancestors, Frontenac berries are small, have high skin-to-pulp ratios, and tend to have colored pulp. These traits result in intense juice color. For rose production, this means that immediate crushing and pressing, without the few hours of skin time allowed in traditional rose production, results in a rich, bright rose-colored juice. Frontenac rose pigments have proven to be very stable, with little or no color loss during fermentation, and no fading as the wine ages. Tavel enthusiasts may dislike the color intensity, but attempts to lighten it by fining have so far resulted in faded wines with hues of sickly salmon. Until research reveals the exact nature of the grape's pigmentation, it may be best to accept the color as a unique and impressive feature.

Continue on page 3

www.vine@doublevineyards.com



10277 Christy Road
Fredonia, NY 14063
Phone: 716-672-8493

For red wine production, 2-5 days of ambient skin time, with caps punched thrice daily, generally produce appropriate color. (It's important to note that pre-fermentation pectinase addition is not recommended, as it tends to turn the must into a slimy mess and inhibits good cap formation.)

In experimental and commercial production, skin contact has been extended for as long as 7 days, but the resulting deepening of color and increase of herbaceous aroma and flavor compounds has been deemed unacceptable by some taste panels when evaluation a year or two post bottling. Later tastings suggest that herbaceous characteristics come back into balance by 3 or 4 years, but additional work is needed to determine the validity of this assessment. In informal aging trials, red Frontenac wines aged up to 10 years have shown no discernable decrease in color intensity, suggesting that phenolic polymerization is minimal. Research is currently underway to determine the nature of phenolic compounds in the grape, so further information regarding this phenomenon should be available in the future.

In addition to high color, Frontenac carries the riparia traits of high sugar and high acid. Soluble solids at harvest normally range from 24-28 °Brix, though measurements as high as 30 °Brix have been reported in some Minnesota vineyards. Since this sugar level is often coupled with titratable acidity (TA) ranging from 9-14 g/L (0.9-1.4%), winemakers may choose to ameliorate with water either before or after fermentation. This method can result in a lighter-bodied wine with less pronounced fruit characteristics. To emphasize the black cherry notes typical to Frontenac, some winemakers choose to stop fermentation before dryness, or sweeten the finished wine. The resulting residual sugar adds body and mouthfeel, enhances the perception of fruit in the finished wine, and reduces the perception of acid. Since palate balance is based on the equation

$$\text{Sweetness} \approx \text{Acidity} + \text{Bitterness}$$

(sugar) (acid) (tannins)

and Frontenac is naturally low in tannins, palate balance in Frontenac wine hinges primarily on the sugar/acid balance. The high acid in this variety allows the winemaker to add enough sugar to boost fruit perception without breaching the threshold of perceptible sweetness. If a sweeter product is desired, however, the high acidity and big fruity notes dominating the palate are capable of balancing a fairly high residual sugar without becoming cloying or over sweet.

To many winemakers, the most frightening aspect of Frontenac is the high TA. At harvest, TA's of 12.0 to 15.0 g/L are typical, though appropriate cluster thinning and canopy management techniques can result in significantly lower numbers. Though these numbers are initially daunting, harvest pH is generally low, ranging from 2.9 to 3.1. For this reason, malolactic fermentation is encouraged, and further deacidification with potassium bicarbonate is occasionally performed.

At the research winery, standard practice to decrease acid includes inducing malolactic fermentation with a strong culture when the primary fermentation is almost, but not quite, finished. Since high acid may stress lactic acid bacteria (LAB) culture, other fermentation parameters are optimized as much as possible—MLF is induced in the primary fermenter without racking, for instance, and yeast lees are stirred thoroughly to re-suspend them in the wine at inoculation, providing nutrients for the LAB. The fermenter is kept as close to 70° F as possible, and LAB nutrient (not DAP) is used at a rate of 1.5g/gal if MLF appears sluggish.

Fermenter's Supply & Equipment

8410 'K' Plaza, Suite #10, Omaha, NE 68127, ph 402-593-9171

www.fermenterssupply.com

Your local source for winemaking supplies and equipment.

We stock a complete line of supplies and equipment, including:

Sodium Meta-bisulfite	Potassium Meta-bisulfite	Grape Concentrate	Fruit Bases
Potassium Sorbate	Acid Blend	Tartaric Acid	Malic Acid
Citric Acid	Ascorbic Acid	Calcium Carbonate	Grape Tannin
White Oak Chips	Toasted Oak Chips	Yeast Energizer/Fermax	Yeast Nutrient
Sparkliod	Bentonite	singlass	LC finings
Phenothalein	Sodium Hydroxide	Red Star Yeast—7 varieties	Crushers
Lalvin Yeast – 5 varieties	Liquid Yeasts	White Labs Liquid Yeast	Presses
Carboys – 3,5,6,6.5 & 14 gallon		Crusher/Stemmers	Drilled Soppers
Airlocks	Corkers	Refractometers	pH Meters
Wine Bottles – clear, green, cobalt blue, dead leaf green	Books-many authors	Brushes	Vinometers
Corks - #9, #8 & #7 Labels	Shrink Capsules - 7 colors	Shrink Capsules—7 colors	
Books – many authors		Hydrometers	

We care about the wine you make as much as you do. We have been making wine for over 26 years from grapes and fruit.

Champagne Corks & Wires Special Order Stainless Steel Tanks & Much More.

Now ordering FROZEN vinifera grapes from California, Washington and Oregon. Call for varieties and prices after August 15, 2005. Delivery will be November or December 2005. Red grapes will be stemmed and crushed, then frozen in buckets. White grapes will be pressed juice, then frozen in buckets.

We are located in Omaha, NE. Coming through town on I-80, take the 84th Street exit, and go six blocks south to 84th & 'J' Sts. On the west side of 84th St. is the 'Just Good Meat' butcher shop. Turn into the butcher shop's parking lot and go down the left side of their building. There is another building behind them across the parking lot. We are door #10 of the 14 bays. We are physically lower on the hill than the butcher shop, so you cannot see us from 84th St. If you go down 'K' St., you will not see us either. We can mail order most items to you.

REGISTRATION FORM

**Ninth Annual Nebraska Winery & Grape Growers Forum and Trade Show
March 3 and 4, 2006
Holiday Inn, Kearney, NE**

Name _____.

(Price includes Friday and Saturday Sessions; Breaks;

Friday afternoon Wine Tasting and Reception; **\$110.00 Member** \$ _____Saturday morning "Early Bird" coffee, Saturday
Luncheon) **\$125.00 Non-Member**

Name _____ \$ _____

(Additional attendee)

Company Name _____

Address _____ City _____, State _____ Zip _____

E-mail: _____ Telephone: _____

I will attend the **Saturday morning** "Early Bird" coffee; (head count is necessary for caterer planning).
with the Exhibitors (cost included in registration) _____

Additional attendee (cost included in registration) _____

Friday Night BanquetName _____ **\$45.00** \$ _____Name (Additional Attendee) _____ **\$45.00** \$ _____

Entrée Choice _____ Prime Rib _____ Salmon

Additional Attendee _____ Prime Rib _____ Salmon

TOTAL \$ _____**Registration Deadline February 15, 2006****Refunds: \$15 handling fee; no refunds after February 15, 2006.****Price after deadline: \$125.00 members \$140.00 non-members****Walk-ins — \$150.00**Make your check payable to: **University of Nebraska Viticulture Program****And send to:**

University of Nebraska Viticulture Program

377 Plant Science Hall

Lincoln, NE 68583-0724

Attn: Donna Michel

GREETINGS FROM PAUL READ IN TASMANIA

Happy New Year and greetings from "Way Down Under" to all in the Nebraska grape and wine industry. It may seem a bit odd to you that I am sweating in the vineyards here in Tasmania, while you are dealing with sub-zero temperatures and snow and ice. The Australians have a map that shows Australia and Southern Hemisphere geography at the top and the Northern Hemisphere at the bottom, and at times it seems indeed that the world here is sort of "upside down". They, like their British ancestors, even drive on the left side of the road. That takes a bit of getting used to, but we've adapted quite well, since it seems a bit like a mirror image - driver on the right side of the car, so it seems natural to drive on the left side of the road.

However, the grapes grow beautifully here, although a bit more slowly than ours, because the spring is long and cool. It is now summer (think of January down here as being like July would be in the Midwest) and the grape berries range from peppercorn size to pea-size (near the end of December as I write this). The canopies are full and healthy-looking, with little disease in evidence. Most growers have applied several sprays for disease control, with sulfur being the most common early-season spray and a product called "Scala" (generic name: pyrimethanil) being used later in the season.

I have been working on several projects, including a "source-sink" study with Richard Smart and a PhD student (Fiona Chopping). This involves removing the growing tip of the shoot and second clusters ("sinks") and removing fully expanded leaves associated with the developing first cluster ("sources", that is, producing carbohydrate via photosynthesis). To eliminate another contributor of carbohydrate, another treatment involves girdling of the shoots below the cluster to prevent carbohydrates and other metabolites to be translocated via the phloem to the developing cluster. Each of these treatments was applied individually and in all possible combinations, with "control" vines receiving none of the treatments. Although this sounds a bit complicated, the concept is relatively simple: can various pruning treatments applied during the growing season influence flowering and fruit development and thus affect subsequent wine quality?

Another interesting project is an attempt to influence bunch architecture in a fashion similar to the approach used in the table grape industry, that is, application of gibberellic acid (GA3) to the bunches to increase the length of the rachis (bunch-stem) and that of the pedicels (stem that bears the individual berries). The purpose of this for wine grapes is to open up the cluster so that there is better air movement through the cluster and thus reduce potential bunch rot of tight-clustered cultivars ('Vignoles' would be an example of such a tight-clustered grape grown in the Midwest). Another potential benefit is better light penetration into the interior of the bunch, therefore enhancing development of color and other components affecting wine quality. I am working with another PhD student (Mark Robertson) on this project and a couple of others. Because of my background and expertise in working with growth regulating chemicals, I helped design this experiment and another in which we physically remove parts of the cluster. Again, the goal is to determine if the better air movement through the bunch will reduce disease and if the better light penetration will have a positive effect on fruit and wine quality. We are already seeing a response in the GA3 experiment, with at least one treatment causing an elongation of the bunch rachis. These and other projects with which I'm working involve a lot of time-consuming hand operations. This causes me to remind myself (and the students) that research is a lot like Edison's characterization of invention: 5% inspiration and 95% perspiration! I'll comment on these experiments and others when I speak at the 9th Annual Nebraska Winery and Grape Growers Forum and Trade Show in March.

Although Tasmania has only an area a little larger than Iowa, it has very diverse topography, climate and soils. It has high mountains and rugged wilderness, shallow bays and beautiful beaches (it hasn't been warm enough yet to swim in the ocean or bays, however), rolling hills and fertile volcanic-derived soils. Many grape growers like to situate their vineyards on north-facing slopes to take advantage of the sun's angle, much like the German industry, for example, where they select land with a south-facing aspect. Much of the vegetable production is in the northwestern part of the island, where the rich volcanic soils are located. There are also some interesting "alternative" crops being grown, including poppies for the pharmaceutical industry, boronia for the food/flavorings industry, pyrethrum (a daisy-like flower) from which an insecticide is produced, and tulip and lily bulb production. However, as I mentioned in an earlier note, sheep production is one of the most important agricultural enterprises, and along with forestry, constitute a large part of Tasmania's total industrial output. Their other major industry is tourism, and now that their school year is over, I've been told to expect to see a lot more cars with "mainland" licenses. They have also progressed greatly in combining tourism with vineyard and winery enterprises. They aim not just to sell wine, but want to sell the **experience** to give the visitor more than just a look at grapes and wine. I'll look forward to discussing various aspects of marrying the tourism and vineyard/winery experiences, as practiced in Tasmania. There could be some ideas for our industry to consider.

Although the time here has just seemed to have flown by, I'm excited about approaching another great year in the Nebraska grape and wine industry. I hope that your new year of 2006 is off to a wonderful start and that you are preparing for the coming growing season. If you haven't already done so, be sure to clean up your equipment, destroy any prunings and debris left in the vineyard, order necessary chemicals, planting stock, tools and equipment and of course register to attend the Forum in March. Let's hope for another great year in 2006!

Cheers,

Paul Read

continued from Page 3

Once MLF is complete, the TA of the wine measured again, and wines that are still deemed to be too acidic are treated with a conservative amount of potassium bicarbonate. The wine is then put into cold stabilization, and TA checked in a couple weeks—if it's still high at that point, another small bicarbonate addition is made, with the total amount still kept below the 15g/gal recommended on the label. It's important to note that bicarbonate additions are made incrementally, rather than all at once, to insure that the acid reduction can be fine-tuned to prevent excessive deacidification or production of off-flavors. Final acids higher than those acceptable in traditional wines can work well in Frontenac; because it is naturally low in tannin, the wine can support higher finished TA's and still present a balanced palate structure. Lack of tannins makes acid the primary wine component giving Frontenac aging potential, so a TA of 9g/L or so is appropriate for reds meant for aging. Calcium carbonate additions for deacidification are generally not recommended, primarily because the off-notes produced are usually deemed unpleasant by consumer panels.

Both increased aging potential and improved flavor characteristics can be achieved through barrel aging or contact with oak chips or staves. Oak-derived aroma and flavor compounds can round out and soften the aggressive one-note cherry typical to Frontenac, producing a more complex wine suitable for consumption with heavier dishes. Enological tannins, which can be added before or after fermentation, are now widely available, but determining the correct rate of addition is a matter of trial and error. All tannin additions require extra aging to allow components to integrate; even so, many tannin-enhanced wines exhibit a disharmonious mouthfeel, where the tannins seem divorced from the rest of the palate, and the whole wine a bit clunky. As more tannin products are available and more winemakers experiment with them, this option may become more approachable, but the inexperienced should approach it with caution.

The final legacy of Frontenac's riparia genetics is the occurrence in some wines of herbaceous notes, described as 'tarry' or 'resinous.' These are usually prominent when the vines have been overcropped, but in properly ripened fruit they should be faint and just enhance complexity. Yeasts that claim to 'enhance varietal character', however, should be avoided; they tend to enhance vegetative character as well. In fermentation trials performed at the Research Enology Lab, trained tasters preferred Frontenac reds produced with RC212, Pasteur Red, and BM 45, and L2226 yeast strains. Wines produced from Fermirouge and Barolo 97 were deemed to lack fruit and/or body.

As a general rule, aroma and flavor characteristics in Frontenac are dominated by a bold cherry note, with lesser hints of black current and general red fruit. Trained sensory panels have also identified notes of grass, green bean, evergreen, tar, and in some cases, chocolate.

Synopsis: Wine styles produced from Frontenac

Commercial production efforts hint at Frontenac's flexibility in various styles. As mentioned above, rose'-style whites have been successfully produced from must allowed 0-8 hours skin contact time, and have proven popular in the commercial market. Though darker than traditional rose's, the wine color is jewel toned and strikingly deep, with no hint of tapering to an insipid salmon. Cool (55°F) fermentation with an aromatic yeast, like Cotes de Blancs, is recommended. The nose and palate showcase a bright, Bing cherry note that is enhanced by an off-dry finish and moderate acidity. Depending on the fruit, sugar levels from bone dry to moderately sweet have shown appropriate balance and customer acceptance.

As a red wine, Frontenac shows a deep garnet color, sometimes tinted with purple. Cherry notes predominate, but are typical of black or sweet cherry rather than the brighter note of the rose. Earthy and slightly herbaceous notes round the palate, supported by a moderate acid backbone. Two to five days of skin contact produce optimum color, and some producers advocate producing separate lots of wine (a 2 day and a 7 day, for instance) and blending prior to cold stabilization to increase complexity. Barrel aging adds welcome structure and a synergistic hint of vanilla. While not as complex as a hearty vinifera, Frontenac packs a solid punch, and stands up well to heavier dishes.

A few creative producers have used Frontenac to produce port-style wines of outstanding quality. In port production, fermentation is stopped through the addition of grape neutral spirits while sugar content is still high, resulting in a product with higher sugar and 15-20% alcohol. The higher acid levels balance the increased sugar beautifully, deepening the typical fruit notes into lush shades of cherry, raspberry, black current, and stewed fruits. Some Frontenac ports exhibit pronounced chocolate notes, which seems dependent on vineyard microclimate. This dessert wine is a showstopper; a Frontenac port won a consensus gold at the 2004 Indy Wine Festival. It wouldn't be surprising to see an increase in commercial production in the coming season.

PROGRAM

DRAFT—Subject to Change

Ninth Annual Nebraska Winery and Grape Growers Forum and Trade Show

March 3rd and 4th, 2006

Holiday Inn, Kearney, NE

Friday—March 3, 2006

8:00a	Registration Opens—Commons Registration Area
8:30a-10:00a	NWGGA Annual Meeting
10:00a-6:30pm	Trade Show opens—Silent Auction opens—Loper Room
11:30a-1:00p	Lunch - On Your Own
12:00-1:00p	NWGGA Board Meeting
1:00p-1:15p	Welcome— Dr. Mark Lagrimini , Head, Dept of Agronomy and Horticulture, University of Nebraska-Lincoln
1:15p-2:15p	Anna Katharine Mansfield , Enologist, University of Minnesota; “Stubbornness, Persistence, and Flexibility: Cold-Climate Cultivars and the New Wine Frontier”
2:15p-3:00p	Terry Bates , Viticulture Research Associate, Cornell University; Soils
3:00-3:30p	Break—visit Tradeshow—Loper Room
3:30-4:15p	Paul Read , Professor, University of Nebraska Viticulture Program—Vineyard experiences in Tasmania
4:15-6:00p	Visit Trade Show—Loper Room
6:00-6:30p	Wine Tasting— Loper Room
7:00p-9:00p	Annual Banquet-Ballroom—Silent Auction Winners

Saturday—March 4, 2006

7:00a-8:30a	Early Bird Trade Show—Coffee and Rolls for attendees
Session I	Advanced Viticulture
8:30a-9:15a	Terry Bates —Soils and Fertilizers
9:15a-9:45a	Growers Panel—Moderated by Terry Bates
9:45a-10:30a	Break—Visit Trade Show—Loper Room
10:30a-11:00a	Mark Steele , UNL CALMIT- “Geospatial Technologies for Nebraska Vineyards.” Conservation Survey Division
11:00a-11:30a	Chelsey Wasem , “Are Insects a Problem in your Vineyard?” Dept of Entomology, University of Nebraska Lincoln
11:30a-1:00p	Grazing Lunch in Trade Show Area
1:30p	Trade Show Closes
1:30p-2:30p	Tim Creger —“Pesticide Regulatory Issues for the Nebraska Grape Industry”, Department of Agriculture
2:30p-5:00	Tourism Partners Panel Part I and II
Session II	Enology and the Commercial Winery
8:30a-9:05a	Anna Katharine Mansfield , “Making Wine from New Cold Hardy University of Minnesota Grapes.”
9:05a-9:45a	Wine Varietal Panel—Moderated by Anna Katharine Mansfield
9:45a-10:30a	Break—Visit Trade Show—Loper Room
10:30a-11:00a	Anna Katharine Mansfield — “Sensory evaluation in the commercial winery—How to establish a tasting panel for small or mid-sized operations.”
11:00a-11:30a	TTB —What to expect in an on-site audit..
11:30a-1:00p	Grazing Lunch in Trade Show Area
1:30p	Trade Show Closes
1:30p-2:30p	Quality Assurance (Winery Council)
2:30p-5:00p	Growers Chemical Application Certification—(Tentative)
Session III	Fundamentals of Viticulture
8:30a-9:15a	Steve Gamet —“Vineyard Environments Across Nebraska”, University of Nebraska Lincoln Viticulture Program
9:15a-9:45a	Growers Panel—Beginning Concerns
9:45a-10:30a	Break—Visit Trade Show—Loper Room
10:30a-11:00a	Terry Bates —“Soils and Site Selection”
11:00a-11:30a	State Level Marketing
11:30a-1:00p	Grazing Lunch in Trade Show Area
1:30p	Trade Show Closes
1:30p-2:30p	Kris Sperry , “Basic Winery Design”
2:30p-5:00p	Beginning Vineyard Concepts. Growers Panel & Growers Panel on Successful Varieties/Contracts
5:00p	Adjourn—Have a safe trip home! See you next year!



377 Plant Science Hall
 Department of Agronomy
 & Horticulture
 Lincoln, NE 68583-0724

It is the policy of the University of Nebraska-Lincoln not to discriminate on the basis of gender, age, disability, race, color, religion and marital status, veterans status, national or ethnic origin or sexual orientation

REMEMBER—Future Forums

2006—March 3 and 4—Holiday Inn, Kearney, NE
 2007—March 2 and 3—Holiday Inn, Kearney, NE
 2008—February 29 and March 1—Holiday Inn, Kearney, NE

FOR YOUR CALENDAR

January 27-28, 2006 (Friday and Saturday) = Iowa Wine Growers Conference, Hotel Ft. Des Moines, Des Moines, IA. Contact Paul Domoto, 515-294-0035 or email: domoto@iastate.edu: **Friday Session (Member Prices: \$75, Wine Tasting \$35; Saturday Sessions—\$95—Evening Banquet—\$50;** Contact 515/262-8323 or 800-383-1682

February 4-6, 2006—Mid-America Wine and Grape Conference; TanTara, MO; Conference Fees—\$125.00—Pre-conference viticulture and enology \$25—MEALS—Sunday Super Bowl Dinner—\$45. Monday Breakfast—\$12; Monday Luncheon—\$14.00; Monday Grand Banquet—\$70. (Prices are with conference fees) Higher pricing without conference fees. Information 573-437-2416.

February 3-4, 2006—Minnesota Grape Growers Cold Climate Grape & Wine Conference, Kahler Grand Hotel, Rochester, MN; **Friday Afternoon Sessions \$50; Saturday Sessions—\$125; Saturday Banquet—\$45.**

March 24, 2006—Pruning and Advanced Viticulture Workshop/Field Day — Kimmel Education and Research Center, Nebraska City, NE

April 20, 2006 —Western Nebraska Pruning and Advanced Viticulture Workshop/Field Day—Scottsbluff, NE

June 24, 2006 —Trellis Systems Field Day—Heritage Vines Vineyard and Czechland Vineyards, Crete and Wilber, NE

July 15, 2006—Multi-State Field Day hosted by the South Dakota vineyard group, probable location—Devils Nest, NE and Yankton, SD area

November 11, 2006—Fall UNL Viticulture Program Workshop, Lincoln, NE, University of Nebraska-Lincoln East Campus Union

Further details of these programs will be announced in the Nebraska VineLines and on the University of Nebraska Viticulture Program website. <http://agronomy.unl.edu/viticulture>

**9th ANNUAL NEBRASKA WINERY AND GRAPE GROWERS FORUM
 AND TRADE SHOW REGISTRATION MATERIALS ENCLOSED**

