University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Studies from the Zoological Laboratory: The University of Nebraska

Parasitology, Harold W. Manter Laboratory of

8-1-1894

Some Notes on the Biological Relations of the Fish Parasites of the Great Lakes

Henry B. Ward University of Nebraska - Lincoln

Follow this and additional works at: http://digitalcommons.unl.edu/zoolabstud



Part of the Zoology Commons

Ward, Henry B., "Some Notes on the Biological Relations of the Fish Parasites of the Great Lakes" (1894). Studies from the Zoological Laboratory: The University of Nebraska. Paper 6. http://digitalcommons.unl.edu/zoolabstud/6

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at Digital Commons@University of Nebraska - Lincoln. It has been accepted for inclusion in Studies from the Zoological Laboratory: The University of Nebraska by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

SOME NOTES ON THE BIOLOGICAL RELATIONS OF THE FISH PARASITES OF THE GREAT LAKES.

(ABSTRACT)

BY H. B. WARD.

A pond or small lake furnishes few variations in depth, temperature, current, or food, and hence the life in it is comparatively uniform. Nowhere else in the world is there found such a continuous body of fresh water as in the Great Lakes. They afford, in themselves alone, an area of water one-fourth greater than that of the state of Nebraska; or, considering the streams and minor lakes within easy reach, no doubt the area is twice as great. Furthermore, there are found here conditions of temperature, depth, light and food more nearly like those prevailing in the ocean. As a result the development of life is greater than in fresh water elsewhere. The existence of a peculiar deep-water fauna has been known for some time, but its extent and the general biological condition of the lakes have never been the object of extended observations. The state of Michigan is bordered by all of the chain except Lake Ontario, and its fisheries are of a magnitude to warrant the large yearly outlay for fish culture under the auspices of the State Fish Commission, by which are planted every year, 100,000,000 fry of the white fish alone. The importance of full and accurate information concerning the entire biological conditions of the lakes has long been felt; and after some tentative work in previous years, the Commission fitted out last summer a temporary laboratory on Lake Saint Clair. Professor J. E. Reighard, the scientific expert of the commission, was appointed director and under his charge work was carried on for nearly three months. To each of the party under his supervision was assigned a particular group; the work on the worms was done by the writer.

This group is represented in fresh water by a few Annelida and free Flat Worms, but most prominently by parasites, including the Trematodes (Flukes), Cestodes (Tape-worms), Nematodes (Round-worms), and Echinorhynchi. Fish of every possible variety were obtained and carefully examined. In the course of the summer one hundred and two individuals belonging to twenty species were dissected and all the parasites found were preserved for future study. The appended table will show the number of each species of fish examined and the number of parasites found in each.

The parasites occur on the gills, in the alimentary canal, in the liver, in the air-bladder, or in the body cavity, so that they are all removed in cleaning the fish and are not to be classed as dangerous to man. Their volume, even in the extreme case, is so small in comparison with that of the fish that, during the summer, a season of abundant nourishment, they can hardly affect at all the health of the host and between those affected by a few parasites and those with many, no difference could be found either in general appearance or in any particular organ so far as could be seen. Five species, Nos. 8, 10, 11, 13 and 16 (see table), were affected by only one kind of parasite and all except the last in small numbers. The other species harboured from two to twenty kinds each.

The Nematodes were the least common of the parasites, being wanting in fourteen species; Acanthocephali were not found in seven species; Trematodes and Cestodes in six species; but while the latter were never very plentiful, the former occurred several times in countless numbers. Some other peculiarities will appear from a study of the appended table.

No. of Nema- toda	10 32 32 0 0 0 0 0 0 0 0 0 0 0 0 0	13
No. of Acan- thocephala.	0 226 226 226 357 226 0 0 0 0 1 1194 154 130	7 44
No. of Cesto-	0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6 45
No. of Trema- toda.	75 11 17 17 26 26 0 0 15 15 52 27 2 0 0 0 9 8 8 8 8 2 2 2 2 0 0 0 0 0 0 0 0 0 0 0 0	5 41
Total No.	88 88 119 30 30 30 30 30 263 263 12 12 12 12 12 1489 154 105 227	
Total No. ex- benima.		
No. free.	00000000000000000000000000000000000000	n
No. Infected.	221-421 010221-422 0221-42 0	olum he co
Common Name.	,	arasite at the head of the
Scientific Name.	Acipenser rubicundus Le Su Ambloplites rupestris (Raf.). Amia calva L. Aplodinotus grunniens (Raf.) Catostomus teres (Mitch.). Coregonus clupeiformis (Mitch.). Coregonus carpio L. Esox lucius L. Esox masquinongy (Mitch.). Hiodon tergisus Le Sueur. Lepidosteus osseus (L.) Lepidosteus osseus (L.) Lepomis gibbosus (L.) Lepomis gibbosus (L.) Micropterus dolomieu (Laccimon aureolum (Le Su Moxostoma aureolum (Le Su Roccus chrysops (Raf.). Stizostedion canadense (C.) Stizostedion vitreum (Mitch.). Otal number of species, 20.	Total number of species free from the parasite at the head of the column Total number of individuals free from the parasite at the head of the column.
	No. of Near-thouse, of No. of Near-thouse, or Cesto. No. of Cesto. No. of Cesto. No. of Acsn. No. of Acsn. No. of Acsn. No. of Acsn.	Scientific Name. Common Name. extended Common Name. Comm

TABLE No. 1. [CONTINUED.]

PARASITES FOUND	No. of Nema- toda.	0 0 0 0 0 0
	No. of Acan- thoceaphala.	0000
	No. of Cesto-da.	01100
	No. of Trema- toda.	f 7 28 0 i 16 k
	Total No.	g 51 h250 i 16 k
ALSO EXAMINED.	Total No. ex- amined.	131 20 20 20 20 20 20 20 20 20 20 20 20 20
	No. free	0 0 0 177
	No. infected	13 0 13 0 2 0 1 0 h 33 h17
	Common Name	Robin. Bonaparte's Gull. Snapping Turtle. Mud Turtle.
	Scientific Name.	Merrula migratoria (L.) Larus philadelphia Ord Chelydra serpentina (L.) Chrysemys marginata (Agassiz). Anodonta ovata Barnes Unio gracilis Lea

Names after Jordan, Manual of the Vertebrates, fifth edition, 1890.

^a The number of Distomata in one specimen was estimated at 3,000; only one-third was counted with a

b Also an indefinite number of cysts in the liver. result of 1,023.

c The total number is too small since there was also an unknown (large) number of a very small Distoma, of which only a few were preserved and included in the figures given.

d Intermediate between A. natalis and A. nebulosus, having 23 rays in the anal fin.

e From gills of two specimens only.

Distoma macrourum Rudolphi.

Including 12 which cannot be assigned to any order at present.

Estimated.

i Monostoma n. sp. k The foot was filled with encyst Cercariea which have not been identified as yet. The number present in each specimen was almost incalculable. About two-thirds of the mussels were infected.