

8-27-2007

Antarctica, the southern ocean, and climate evolution: Insights from drilling, coring, and geophysical surveys

Timothy R. Naish
Victoria University of Wellington

Ross Powell
Northern Illinois University

Follow this and additional works at: <http://digitalcommons.unl.edu/andrillrespub>

 Part of the [Environmental Indicators and Impact Assessment Commons](#)

Naish, Timothy R. and Powell, Ross, "Antarctica, the southern ocean, and climate evolution: Insights from drilling, coring, and geophysical surveys" (2007). *ANDRILL Research and Publications*. 9.
<http://digitalcommons.unl.edu/andrillrespub/9>

This Article is brought to you for free and open access by the Antarctic Drilling Program at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in ANDRILL Research and Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Citation: Cooper, A.K. et al. (2007), Program book for the 10th International Symposium on Antarctic Earth Sciences: in Antarctica: A Keystone in a Changing World-Online Proceedings of the 10th ISAES, edited by A. K. Cooper and C. R. Raymond et al., USGS Open-File Report 2007-1047, 141 p.

10th International Symposium on Antarctic Earth Sciences



August 26 - August 31, 2007



1.P2.A: Antarctica, the southern ocean, and climate evolution: Insights from drilling, coring, and geophysical surveys ORAL

1.P2.A-1: The diatom record of the ANDRILL – McMurdo Ice Shelf project drillcore

Reed Scherer¹, Diane Winter², Charlotte Sjunneskog³ and Paola Maffioli⁴, (1)Department of Geology and Environmental Geosciences, Northern Illinois University, DeKalb, IL 60115, (2)University of Nebraska - Lincoln, Department of Geosciences, Lincoln, NE 68588, (3)Department of Geology and Geophysics, Louisiana State University, Baton Rouge, LA 70803, (4)Department of Geological Sciences and Geotechnologies, University of Milano-Bicocca, Milan, Italy

The inaugural drilling season of ANDRILL, recovered a 1,285 m core, AND-1B, with ~99% recovery. The core contains a superb record of Antarctic continental shelf sediments, providing an unparalleled record of climate change through a critical interval in Earth history. The upper c. 600m of core, reflecting Pliocene and early Pleistocene deposition, is composed of alternating glacial diamictites and diatomites, with episodic volcanic facies. The diatomites document extended periods of open marine conditions with reduced ice, in an area currently covered by a thick ice shelf. The diatomites reflect high biosiliceous productivity, and most reflect warmer than present conditions with variable sea ice and ice rafting. Many likely represent an absence of a large ice shelf, whereas diamictites reflect glacial advances. Analysis of the diatom assemblages will result in a new biostratigraphic zonation and high resolution paleoenvironmental reconstructions.

1.P2.A-2: Preliminary ⁴⁰Ar/³⁹Ar results from the AND-1B core

Jake I. Ross, William C. McIntosh and Nelia W. Dunbar, Department of Earth and Environmental Sciences, New Mexico Tech, 801 Leroy Pl, Socorro, NM 87801

Preliminary ⁴⁰Ar/³⁹Ar dating of 13 volcanic samples from four stratigraphic intervals within the AND-1B core provide key age constraints for the development of an accurate age-model. To date seven analyses have yielded statistically robust and stratigraphically meaningful ages. The four different stratigraphic intervals represented by the seven successfully dated samples are: 1) 85.53-85.85 mbsf felsic tephra (1.014±0.004 Ma), 2) ~112-145 mbsf basaltic tephra (1.65±0.05 to 1.67±0.05 Ma), 3) 646.30-649.34 mbsf basaltic lava flow (6.48±0.13 Ma), and 4) ~1280 mbsf volcanic clasts (maximum depositional age 13.57±0.13 Ma).

1.P2.A-3: Preliminary chronostratigraphy for the upper 700 m (upper Miocene–Pleistocene) of the AND-1B drillcore recovered from beneath the McMurdo Ice Shelf, Antarctica

Gary S. Wilson¹, Richard H. Levy², Greg Browne³, Fabio Florindo⁴, Stuart A. Henrys³, Ian Graham³, William C. McIntosh⁵, R. M. McKay⁶, T. R. Naish⁶, Christian Ohneiser¹, Ross D. Powell⁷, Jake I. Ross⁵, Leonardo Sagnotti⁴, Reed Scherer⁷, Charlotte Sjunneskog⁸, C. Percy Strong³, Marco Taviani⁹ and Diane Winter¹⁰, (1)Department of Geology, University of Otago, PO Box 56, Dunedin, New Zealand, (2)ANDRILL Science Management Office, University of Nebraska - Lincoln, 126 Bessey Hall, Lincoln, NE 68588-0341, (3)GNS Science, 1 Fairway Drive, Avalon, Lower Hutt, New Zealand, (4)Istituto Nazionale di Geofisica e Vulcanologia, Via di Vigna Murata, 605, Rome, I-00143, Italy, (5)Department of Earth and Environmental Sciences, New Mexico Tech, 801 Leroy Pl, Socorro, NM 87801, (6)Antarctic Research Centre, Victoria University of Wellington, Wellington, New Zealand, (7)Department of Geology and Environmental Geosciences, Northern Illinois University, DeKalb, IL 60115, (8)Department of Geology and Geophysics, Louisiana State University, Baton Rouge, LA 70803, (9)ISMAR-CNR, Via Gobetti 101, Bologna, Italy, (10)University of Nebraska - Lincoln, Department of Geosciences, Lincoln, NE 68588

Chronostratigraphic data available for the preliminary age model for the upper 700 m for the AND-1B drill core include diatom biostratigraphy, magnetostratigraphy, ⁴⁰Ar/³⁹Ar ages, ⁸⁷Sr/⁸⁶Sr ages and surfaces of erosion identified from physical appearance in the drill core. The age data allow a relatively well-constrained age model to be constructed. ~70% of the AND-1B magnetic polarity stratigraphy can be correlated with the Geomagnetic Polarity Time Scale (GPTS). Unique correlation is not possible in several coarse diamictite intervals with closely spaced glacial surfaces of erosion and sparse microflora. However, the age model indicates relatively rapid (up to 1m / k.y.) accumulation punctuated by several half to million year hiatuses representing more than half of the last 7 m.y. in the drillcore. The mid-late Pleistocene is represented by superimposed diamictite units separated from late Pliocene alternating diamictites / diatomites by a ~ 1 m.y. hiatus.

1.P2.A-4: Future geological drilling in Antarctica – a discussion paper on ANDRILL and beyond

Peter Barrett¹, P.N. Webb², Dieter Futterer³, Claudio Ghezzi⁴, M. R. A Thomson⁵, A. R. Pyne¹ and F. Rack⁶, (1) Antarctic Research Centre, Victoria University of Wellington, Wellington, New Zealand, (2)The Ohio State University, (3)Alfred Wegener Institute, (4)Dipartimento di Scienze della Terra, Università di Siena, Via Laterina, 8, Siena, 53100, Italy, (5)Stone House, Thornton Rust, (6)ANDRILL Science Management Office

Geological drilling technology developed over the last quarter century has provided access to critical knowledge on the climatic and tectonic history of the Victoria Land margin of Antarctica, giving us a new understanding of the history and behaviour of the Antarctic ice sheet in this region over the last 34 million years. The challenge now is to develop a framework within the Antarctic science and logistics communities and other relevant groups for further projects to extend this technology to other areas both around the Antarctic margin and into the interior. This paper reviews some of the issues and offers a way forward.