

1-2013

Corn Cobs On Deck for Cellulosic Feedstock


Brian J. Wienhold

Soil and Water Conservation Research Unit, USDA-ARS, bwienhold1@unl.edu

Ann Perry

USDA-ARS, ann.perry@ars.usda.gov

Follow this and additional works at: <http://digitalcommons.unl.edu/usdaagresmag>

 Part of the [Agriculture Commons](#), [Animal Sciences Commons](#), [Food Science Commons](#), and the [Plant Sciences Commons](#)

Wienhold, Brian J. and Perry, Ann, "Corn Cobs On Deck for Cellulosic Feedstock" (2013). *Agricultural Research Magazine*. 10.
<http://digitalcommons.unl.edu/usdaagresmag/10>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agricultural Research Magazine by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Corn Cobs On Deck for Cellulosic Feedstock

When energy officials proposed using crop residues to produce cellulosic ethanol, concerned soil scientists took to the fields to learn more about how these residues protect soil from erosion and enhance soil quality. Agricultural Research Service soil scientist Brian Wienhold focused on a single component of residue—the corncob.

“We didn’t have data on how postharvest cob residues might protect soil quality,” says Wienhold, who works in the ARS Agroecosystem Management Research Unit in Lincoln, Nebraska. “But corncobs make up 20 percent of residue by weight, which means that the average U.S. production of corn could provide 40 to 50 million tons of cobs for feedstock every year.”

Wienhold led colleagues in studies that compared runoff from no-till corn fields where postharvest crop residues were either removed or retained. The scientists also removed the cobs from half of the test plots that were protected by the residues. Then they generated two simulated rainfall events; the first occurred when the fields were dry, and the next occurred 24 hours later when the soils were almost completely saturated.

During the first event, on plots where residue was removed, runoff began around 200 seconds after the “rain” began, whereas runoff in the residue-protected plots didn’t start until around 240 seconds after it started to “rain.” Runoff from the residue-free plots

contained 30 percent more sediment than runoff from all the residue-protected plots. The presence or absence of cobs on the residue-protected plots did not affect sediment loss rates.

Wienhold’s team concluded that even though cob residues did slightly delay the start of runoff, they did not affect rates of sediment loss. The results showed that the cobs could be removed from other residue and used for feedstock without significantly interfering with the role of crop residues in protecting soils.

In related studies, Wienhold examined how the removal of cob residues affected soil nutrient levels. He placed litter bags containing cob pieces on the surface of no-till fields or buried them 0 to 4 inches

deep in the soil. Every 2 months, he tested cob samples from the bags for levels of carbon, nitrogen, phosphorus, potassium, sulfur, calcium, manganese, iron, magnesium, copper, and zinc. Over the course of a year, his sampling indicated that cobs were a source of soil potassium, but that they weren’t a significant source of any other plant nutrients.

Wienhold believes these findings demonstrate that harvesting cobs for biofuel production would not result in any notable loss of soil quality. This means that Nebraska’s York County—where average corn production results in around 0.2 million tons of cobs—could potentially provide enough cob feedstock every year to keep two 10.5-million-gallon ethanol plants in business.

Results from this work have been published in *Agronomy Journal*.—By **Ann Perry**, ARS.

This research is part of Agricultural System Competitiveness and Sustainability, an ARS national program (#216) described at www.nps.ars.usda.gov.

*Brian Wienhold is in the USDA-ARS Agroecosystem Management Research Unit, Keim Hall, University of Nebraska, Lincoln, NE 68583-0937; (402) 472-1484, brian.wienhold@ars.usda.gov. **



PEGGY GREB (D2732-1)

Soil scientists Brian Wienhold (left) and Gary Varvel compare corncob residue in various stages of decomposition in no-till corn in Lincoln, Nebraska.