

9-1989

Biology and Characterization of *Prodiplosis Longifila* (Diptera: Cecidomyiidae) on Lime in Florida

J. E. Pena
University of Florida

Raymond Gagne
ARS

R. Duncan
University of Florida

Follow this and additional works at: <http://digitalcommons.unl.edu/systementomologyusda>



Part of the [Entomology Commons](#)

Pena, J. E.; Gagne, Raymond; and Duncan, R., "Biology and Characterization of *Prodiplosis Longifila* (Diptera: Cecidomyiidae) on Lime in Florida" (1989). *USDA Systematic Entomology Laboratory*. 15.
<http://digitalcommons.unl.edu/systementomologyusda/15>

This Article is brought to you for free and open access by the Entomology Collections, Miscellaneous at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA Systematic Entomology Laboratory by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

- HAVRANEK, D. 1987. *Melanagromyza tomaterae* (Diptera: Agromyzidae) a tomato pest in the States of Táchira and Mérida, Venezuela. Fl. Entomol. 70(2): 294-296.
- SPENCER, K. A. 1963. A synopsis of the neotropical Agromyzidae (Diptera). Trans. R. ent. Soc. London 115: 291-389.
- SPENCER, K. A. 1973. The Agromyzidae (Diptera) of Venezuela. Rev. Fac. Agron. (Maracay) 7(2): 5-107. (Venezuela).
- STEYSKAL, G. C. 1972. Two new species of *Melanagromyza* Hendel (Diptera: Agromyzidae) that bore in tomato stalks in Colombia and Ecuador. J. Washington Acad. Sci. 62: 265-267.

BIOLOGY AND CHARACTERIZATION OF *PRODIPLOSIS LONGIFILA* (DIPTERA: CECIDOMYIIDAE) ON LIME IN FLORIDA

J. E. PEÑA¹, RAYMOND J. GAGNÉ², AND R. DUNCAN¹

¹University of Florida, IFAS

Tropical Research and Education Center

18905 S.W. 280 Street, Homestead, Florida 33031

²Systematic Entomology Laboratory

Agricultural Research Service

USDA c/o U.S. National Museum NHB 168

Washington, D.C. 20560

ABSTRACT

Prodiplosis longifila Gagné (Diptera: Cecidomyiidae) is a polyphagous, Neotropical species with a known range that extends into southern Florida. This gall midge feeds on the flower ovaries of lime and can cause premature flower abscission. Details of its biology on lime are given, the life stages illustrated, and the three larval instars described in detail.

RESUMEN

Prodiplosis longifila Gagné (Diptera: Cecidomyiidae) es una especie polífaga y neotropical y su rango se extiende hasta el sur de la Florida. Esta mosquita destruye los ovarios de las flores de los limones y puede causar caída prematura de las flores. Se dan detalles sobre su biología en los limones, se hacen ilustraciones de los diferentes estados, y se describen los tres estadios larvarios.

Prodiplosis longifila Gagné was recently reported feeding in flowers of lime, *Citrus aurantifolia* (Christm.) Swingle (Rutaceae), in Dade and Collier Counties, Florida (Peña et al. 1987). Larvae completely destroy the ovaries of the flowers and may cause premature flower abscission. The only previous record of this species from the United States was from wild cotton, *Gossypium* sp. (Malvaceae), from Monroe Co., Florida (Gagné 1986). *Prodiplosis longifila* is otherwise known from Colombia and Peru, where it is known to be a pest of tomatoes, potatoes, alfalfa, and other commodities (Gagné 1986).

In this report we present the first known details of the life history of *P. longifila* on lime and describe the larval instars.

METHODS

The biology of *P. longifila* was studied in a 0.7 ha, 10-yr old 'Tahiti' lime orchard in Homestead, Florida and in the laboratory at the Tropical Research and Education Center, Homestead, Florida. Approximately 30 trees were used in this study. Observations were made over a period of 86 days (November 1987 through January 1988) for five generations of the gall midge.

Lime flowers in their first stages of development (less than 1 mm in diameter) were bagged with nylon mesh. Flowers were left exposed for 12 h and egg masses collected after females oviposited. Egg masses ($n = 14$) were transferred to the laboratory and eggs ($n = 258$) placed individually in petri dishes (5.2 cm in diameter) lined with filter paper. Petri dishes were held at 27°C and 84 ± 2 RH and observed daily. After eclosion each larva was removed and transferred to similar petri dishes with excised flowers. Larvae were removed two times during the day to determine the duration of each instar. Full-grown larvae from a set of infested flowers were allowed to drop into one of 12 360 ml carton containers, each containing 400 g of previously sterilized 'Rockdale' soil. Pupae were collected from the soil surface and from different soil depths (1.5-2.9 cm, 3.0-5.4 cm, and 5.5-8.0 cm). Adult emergence was determined in the field by placing infested flowers on plastic containers (43 x 59 x 26 cm) filled with 'Rockdale' soil. Clear-plastic circular plates (24 cm diam.) coated with Tanglefoot® were placed on top of the containers. Adults trapped on the plastic plates were counted every 2 h during a 24 h period on five separate days (November 19, 20, 21, 1987 and January 14, 15, 1988). Adults kept in the laboratory were maintained in a rearing chamber at 27°C and 84 ± 2 RH with ca. 12 h photoperiod. Each adult was kept in a petri dish lined with filter paper and provided with ca. 0.01 ml of honey and 0.01 ml water or left without food.

Larvae used for detailed descriptive study were mounted on glass slides in Hoyer's mounting medium or Canada balsam for study. Measurements were made of 20 specimens of each instar. Representative specimens have been deposited in the National Museum of Natural History, Washington, D.C.

RESULTS AND DISCUSSION

Eggs are transparent, elongate-ovoid, 0.265 ± 0.035 SE mm x 0.096 ± 0.002 SE mm (Fig. 2). They are deposited on stamens or styles, usually of flowers that are 0.46-0.65 cm in diameter [$\chi^2.05[8]$; $P < 0.05$] or in flowers in which a small opening in the corolla allows the style to protrude. One to 59 eggs (mean = 12.6 ± 2 SE) were found per infested flower under laboratory conditions. Larvae hatched in 1.4 ± 0.045 SE days.

Larvae are almost transparent when newly eclosed, turn white 1.2 days later and become yellowish to orange when full-grown (Fig. 1c). There are three instars. Major anatomical features common to all three (Figs. 5-13) are a small head capsule with one-segmented, conical antennae, a neck segment, three thoracic segments, and nine abdominal segments, each segment with papillae characteristic for the Cecidomyiini in number and position (Gagné in press). The spiracles of the eighth abdominal segment are situated posteriorly.

The first instar (Figs. 5-7) ranges from 0.40-0.92 mm in length. The head capsule is 0.045 ± 0.003 mm wide at the posterior end. This instar has only one pair of spiracles, on the eighth abdominal segment (Fig. 5), and that is the most conspicuous difference between this instar and the remaining two.

The second instar (Figs. 8-10) ranges from 0.76-1.85 mm in length. The head capsule






Fig. 1-4. *Prodiplosis longifila*. 1, adult. 2, eggs. 3, third instar. 4, pupa.

is 0.050 ± 0.005 mm wide at its posterior end. This instar has the full complement of spiracles for cecidomyiid larvae, one pair on the first thoracic segment and one pair on each of the first through eighth abdominal segments. It differs from the third instar in lacking a spatula and having the corniform pair of papillae on the terminal segment less strongly developed relative to the other three pairs.



Figs. 5-10. Larvae, *Prodiplosis longifila*. 5-7, First instar: 5, dorsal view; 6, head through second thoracic segment, ventral; 7, abdominal segments seven through nine, dorsal. 8-10, Second instar: 8, dorsolateral view; 9, head through second thoracic segment, ventrolateral; 10, abdominal segments seven through nine, dorsolateral. Line = 0.1 mm.

The third instar (Figs. 11-13) ranges from 1.15-1.90 mm in length. The head capsule is 0.050 ± 0.005 mm wide at its posterior end. The spiracular system is similar to that of the second instar. The distinguishing feature of this instar is the clove-shaped spatula on the venter of the first thoracic segment (Fig. 12). The papillar setae, except for those



Figs. 11-13. Third instar, *Prodidiplosis longifila*. 11, dorsal view with the ventral spatula dotted in to show relative size and position; 12, head through first thoracic segment, ventral; 13, abdominal segments seven through nine, dorsal. Line = 0.1 mm.

of the terminal segment, are shorter relative to the width of the papillar bases than in previous instars.

The head capsule width of cecidiomyiid larvae usually becomes significantly larger in each instar (Gagné and Hatchett in press, Roskam 1977, 1979, Wilson 1966), but in *P. longifila* the head of the second instar is as wide as that of the third. First instar head capsules are sometimes also as wide as those in following instars. Antennae are similarly shaped in each instar.

Two days after hatching, 17 of 20 larvae collected were first instar, three were second instars. Three days after hatching, 12 of 20 larvae were second instars and eight were third instars. Four days after hatching all larvae were third instars. The entire larval stage last 9 ± 1.63 days.

Larvae feed on the surface of the ovary, pistils, and stamens of lime flowers. One to 66 larvae were found per infested flower (24.26 ± 4.15 SE). Mature larvae drop to the ground, where they penetrate the soil more frequently at a depth of 1.5 cm than at




Fig. 14. Average daily catches of *Prodioplosis longifila* adults every two h.

other depths ($\chi^2_{0.005[3]} = 7.81$; $P < 0.005$). They then spin a white cocoon, which sometimes incorporates sand grains.

Pupae are 0.85-1.00 mm long and pale yellow when newly molted (Fig. 4). The head and thorax turn black 3.02 ± 0.92 SE days later. This stage lasted 4.11 ± 1.22 SE days unless parasitized.

Adults are about 1.5 mm in length (Fig. 1). Wing length is 1.42 ± 0.04 mm in males, 1.53 ± 0.02 mm in females. A technical description of this stage is given in Gagné (1986). Peak emergence occurred between 1700 and 2300 hours, but a few adults emerged at other times (Fig. 14). Temperatures recorded during periods of adult activity ranged from 17-20°C. Relative humidity ranged between 69-98%. The female to male sex ratio of emerging adults fluctuated between 70:30 and 50:50 ($n = 87-40$). Adults reared in the laboratory survived 1.06 ± 0.24 days if not fed; if held in vials with available honey and water, they survived 8.30 ± 1.21 days, with a maximum of 16 days.

Prodioplosis longifila is parasitized by *Synopeas* sp. (Platygasteridae), an egg-larval parasitoid. Parasites emerged 14-16 days after pupation of the gall midge.

ACKNOWLEDGMENTS

We thank Deborah Leather Roney, Arlington, Virginia, for the detailed larval draw-

ings and L. Stange, Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, for identifying the parasitoid of *P. longifila*. We are grateful to R. M. Baranowski, University of Florida, Homestead, P. E. Boldt, Agricultural Research Service, USDA, Temple, Texas, N. E. Woodley, Agricultural Research Service, USDA, Washington, D.C., H. C. Roskam, University of Leiden, The Netherlands and L. F. Wilson, East Lansing, Michigan for commenting on a draft of the ms. The work was supported by a grant from the Florida Lime and Avocado Committee. This paper is Florida Agricultural Experiment Station Journal Series No. 9573.

REFERENCES CITED

- GAGNÉ, R. J. 1986. Revision of *Prodiplosis* (Diptera: Cecidomyiidae) with descriptions of three new species. Ann. Entomol. Soc. Am. 79: 235-245.
- . In press. The plant feeding gall midges of North America. Cornell University Press, Ithaca, New York.
- GAGNÉ, R. J. AND J. H. HATCHETT. In press. Larval instars of the Hessian fly (Diptera: Cecidomyiidae). Ann. Entomol. Soc. Am.
- PEÑA, J. E., R. M. BARANOWSKI, AND R. T. McMILLAN, JR. 1987. *Prodiplosis longifila* (Diptera: Cecidomyiidae) a new pest of citrus in Florida. Florida Entomol. 70: 527-529.
- ROSKAM, H. C. 1977. Biosystematics of insects living in female birch catkins. I. Gall midges of the genus *Semudobia* Kieffer (Diptera: Cecidomyiidae). Tijdschr. Entomol. 120: 153-197.
- . 1979. Biosystematics of insects living in female birch catkins. II. Inquiline and predaceous gall midges belonging to various genera. Netherlands J. Zool. 29: 283-351.
- WILSON, L. F. 1966. Life history, habits, and damage of the boxelder leaf gall midge, *Contarinia negundifolia* Felt (Diptera: Cecidomyiidae) in Michigan. Canadian Entomol. 98: 777-784.

REVIEW OF THE NEW WORLD DIMARINI WITH THE
DESCRIPTION OF A NEW GENUS FROM PERU
(NEUROPTERA: MYRMELEONTIDAE)

LIONEL A. STANGE
Florida State Collection of Arthropods,
Division of Plant Industry,
P. O. Box 1269
Gainesville, Florida 32602

ABSTRACT

The genus *Dimares* Hagen is divided into two genera, *Dimares* with one species in Brazil and Argentina, and a new genus *Millerleon* with three species in the coastal desert of Peru. A world-wide key for the genera in the tribe Dimarini is presented based on adults and larvae. Also, a key for the species in the genus *Millerleon* is presented. A diagnosis of the tribe, the two American genera and the American species are provided with new records for the species. Data are provided about the biology and morphology of the larvae of *Dimares elegans* and *Millerleon bellulus*.

