University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Martin Centurion Publications

Research Papers in Physics and Astronomy

1-1-2006

Erratum: Dynamics of filament formation in a Kerr medium [Phys Rev. A 71, 063811 (2005)]

Martin Centurion University of Nebraska-Lincoln, mcenturion2@unl.edu

Ye Pu California Institute of Technology, yepu@sunoptics.caltech.edu

Mankei Tsang California Institute of Technology

Demetri Psaltis California Institute of Technology

Follow this and additional works at: http://digitalcommons.unl.edu/physicscenturion



Part of the Physics Commons

Centurion, Martin; Pu, Ye; Tsang, Mankei; and Psaltis, Demetri, "Erratum: Dynamics of filament formation in a Kerr medium [Phys Rev. A 71, 063811 (2005)]" (2006). Martin Centurion Publications. Paper 13. http://digitalcommons.unl.edu/physicscenturion/13

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at Digital Commons@University of Nebraska -Lincoln. It has been accepted for inclusion in Martin Centurion Publications by an authorized administrator of Digital Commons@University of Nebraska - Lincoln.

PHYSICAL REVIEW A 74, 069902(E) (2006)

Erratum: Dynamics of filament formation in a Kerr medium [Phys Rev. A 71, 063811 (2005)]

Martin Centurion, Ye Pu, Mankei Tsang, and Demetri Psaltis (Received 29 October 2006; published 18 December 2006)

DOI: 10.1103/PhysRevA.74.069902 PACS number(s): 42.65.Jx, 42.65.Tg, 89.75.Kd, 99.10.Cd

We have found an error in this paper. In our calculation of the intensity threshold for breakdown in CS_2 we erroneously used the ionization energy of molecular CS_2 (10.08 eV). The correct value should be 3.3 eV [1], the band gap energy of liquid CS_2 . Using the band gap energy, nonlinear absorption will be a three-photon process, and the calculated breakdown threshold becomes 8.5×10^{11} W/cm² (as opposed to the value of 2×10^{13} W/cm² reported previously). Multiphoton absorption and plasma defocusing could then become significant for the intensity of the filaments, $I_{fil} = (5.9 \pm 2.4) \times 10^{11}$ W/cm², that was measured experimentally.

In order to test our conclusion that a model including a fifth order nonlinearity is the best fit to the experimental data, we have numerically simulated the pulse propagation in CS_2 both with and without fifth order nonlinearity (n_4) and plasma effects. The numerical model is the same in this paper with the addition of plasma effects (negative index change and multiphoton absorption). The parameters for the plasma effects were calculated according to the method described in Ref. [2]. The numerical results show that for the case of only plasma effects (with n_4 =0) a filament forms with an intensity of 3 \times 10¹² W/cm², well above the experimental value. For both cases including n_4 (with and without plasma effects) the intensity peaks at about 10¹² W/cm², and the inclusion of plasma effects results in a loss of 15% after 4 mm of propagation. It should be noted that while the losses due to multiphoton absorption can be significant, they are not essential for filament formation. In addition, the simulation with only plasma effects resulted in very high plasma density (>10¹⁹ cm⁻³) which was not observed experimentally [3].

In conclusion, our numerical results show that the effect of plasma alone cannot reproduce the experimental results, while including a fifth order nonlinearity (with or without plasma) generates good agreement with the experiments. The main conclusion of the numerical section of the original paper then remains unchanged in that the fifth order nonlinearity provides good agreement with the experiment while plasma effects alone do not.

^[1] A. Brodeur et al., J. Opt. Soc. Am. B 16, 637 (1999).

^[2] P. K. Kennedy, IEEE J. Quantum Electron. 31, 2241 (1995).

^[3] M. Centurion, Y. Pu, and D. Psaltis, J. Appl. Phys. 100, 063104 (2006).