

January 2004

Residual Effects of Manure and Compost Applications on Corn Production and Soil Properties

Bahman Eghball
USDA-ARS

Daniel Ginting
University of Nebraska - Lincoln, dginting3@unl.edu

John E. Gilley
University of Nebraska - Lincoln, john.gilley@ars.usda.gov

Follow this and additional works at: <http://digitalcommons.unl.edu/biosysengfacpub>



Part of the [Biological Engineering Commons](#)

Eghball, Bahman; Ginting, Daniel; and Gilley, John E., "Residual Effects of Manure and Compost Applications on Corn Production and Soil Properties" (2004). *Biological Systems Engineering: Papers and Publications*. 14.
<http://digitalcommons.unl.edu/biosysengfacpub/14>

This Article is brought to you for free and open access by the Biological Systems Engineering at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Biological Systems Engineering: Papers and Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Residual Effects of Manure and Compost Applications on Corn Production and Soil Properties

Bahman Eghball,* Daniel Ginting, and John E. Gilley

ABSTRACT

Residual effects of manure or compost application on crop production and soil properties can last for several years. This study was conducted to evaluate residual effects of annual or biennial applications of N- and P-based composted and noncomposted beef cattle (*Bos taurus*) feedlot manure, chemical fertilizer, and no-treatment check on corn (*Zea mays* L.) production and soil properties. Manure and compost were applied from 1992 to 1995, and the residual effects were determined from 1997 to 1999. Residual effects of N- and P-based manure and compost applications on corn grain yield and N uptake lasted for at least one growing season while the effects on soil properties were longer lasting. Soil P can contribute to crop P uptake for >4 yr after N-based manure or compost application had ceased. The residual effects of manure and compost applications significantly increased soil electrical conductivity and pH levels and plant-available P and NO₃-N concentrations. Four years after the last application, P leaching to a soil depth of 45 to 60 cm was observed with N-based manure or compost application. No residual effects of manure and compost applications on soil NH₄-N were observed. Averaged across years, soil total C concentrations or quantities were not different among the treatments, indicating that total C was not a sensitive indicator. Residual effects of N- or P-based manure or compost application increased crop production for one year and influenced soil properties for several years.

APPLICATION OF MANURE or composted manure can result in increased soil concentrations of nutrients and organic matter (Chang et al., 1991; Eghball, 2002). The residual effects of increased nutrients and organic matter in soil following manure or compost application on crop yield and soil properties can last for several years (Mugwira, 1979; Wallingford et al., 1975). Significant residual effects of dairy manure application rates that ranged from 22.5 to 270 Mg dry weight ha⁻¹ (530–6400 kg N ha⁻¹) on crop yield increase were observed for the higher 180 and 270 Mg ha⁻¹ rates. These high rates also increased NO₃-N leaching into the soil in a study conducted in Alabama (Mugwira, 1979). Even the lowest application rate of 22.5 Mg ha⁻¹ would have provided nearly twice the N-based requirement for the millet (*Pennisetum americanum* L.) used (Binford et al., 2000). Four years after application, residual effects of one-time application of beef feedlot manure at rates varying from 123 to 590 Mg dry weight ha⁻¹ (1280–6140 kg N ha⁻¹) resulted in a quadratic increase in corn grain yield but also in increased leaching of NO₃-N and Na to a depth of at least 1 m (Wallingford et al., 1975).

B. Eghball and J.E. Gilley, USDA-ARS, 121 Keim Hall, Univ. of Nebraska–Lincoln, Lincoln, NE 68583; and D. Ginting, Dep. of Agron. and Hortic., Univ. of Nebraska–Lincoln, Lincoln, NE 68583. Joint contribution of USDA-ARS and Univ. of Nebraska Agric. Res. Div., Lincoln, NE, as Paper no. 14066. Received 22 Apr. 2003. *Corresponding author (BEGHBALL1@unl.edu).

Published in *Agron. J.* 96:442–447 (2004).
© American Society of Agronomy
677 S. Segoe Rd., Madison, WI 53711 USA

Residual effects of manure or compost application can maintain crop yield level for several years after manure or compost application ceases since only a fraction of the N and other nutrients in manure or compost become plant available in the first year after application (Motavalli et al., 1989; Eghball et al., 2002). Eghball and Power (1999) found that 40% of beef cattle feedlot manure N and 20% of compost N were plant available in the first year after application, indicating that about 60% of manure N and 80% of compost N became plant available in the succeeding years, assuming little or no loss of N due to NO₃-N leaching or denitrification.

Residual effects of organic materials on soil properties can contribute to improvement in soil quality for several years after application ceases (Ginting et al., 2003). Increased levels of soil N, P, K, pH, and C levels in the soil can increase crop yield beyond the application years. Soil pH, organic matter, total N, NO₃-N, and P levels were still elevated 4 yr after dairy manure application ceased (Mugwira, 1979; Lund and Doss, 1980). Eghball et al. (2003) found that the increased plant-available P level in soil following N-based manure or compost application can contribute to crop P uptake for up to 10 yr without any additional P addition. Ginting et al. (2003) did not find increased emission of greenhouse gasses (CO₂, CH₄, and N₂O) as a result of residual manure and compost applications that ceased 4 yr earlier.

Residual effects of manure application have been reported for studies where excessive rates of manure had been applied (Wallingford et al., 1975; Mugwira, 1979; Lund and Doss, 1980). Nitrogen- and P-based manure or compost application provides rates that are agronomically and environmentally sound. Nitrogen-based manure or compost application can increase soil P levels (Eghball and Power, 1999). However, in areas where the risk of P transport in runoff is not a concern, N-based applications can be made. The objective of this study was to determine the residual effects of N- and P-based manure and composted manure application strategies on corn production and soil properties.

MATERIALS AND METHODS

Manure or Compost Application

The experiment was initiated in 1992 on a Sharpsburg silty clay loam soil (fine, smectitic, mesic Typic Argiudolls) under rainfed conditions until 1996 at the University of Nebraska Agricultural Research Center near Mead, NE. The Sharpsburg series consists of deep, moderately well drained soils formed in loess on uplands and high benches. Permeability is moderately slow in the upper part of the soil profile and moderate in the lower part. The study area had a Bray and Kurtz no.1 soil test P concentration of 69 mg kg⁻¹, a pH of 6.2, and a soil

Abbreviation: EC, electrical conductivity.

Table 1. Composted and noncomposted manure dry weight and N and P applications in 4 yr in eastern Nebraska.

Treatment	Dry weight				Nitrogen					Phosphorus				
	1992	1993	1994	1995	1992	1993	1994	1995	Total	1992	1993	1994	1995	Total
	Mg ha ⁻¹				kg ha ⁻¹									
Manure for N	46.9	18.5	12.1	14.5	378	189	189	189	945	107	92	40	46	285
Manure for P	28.3	6.4	6.6	2.7	227 (60)†	66 (80)	103 (74)	35 (99)	744	64	32	21	8‡	125
Manure for N/2 yr	93.9	–	36.3	–	756	–	567	–	1323	214	–	119	–	333
Manure for P/2 yr	56.4	–	19.8	–	454	–(60)	308	–(67)	889	129	–	64	–	193
Compost for N	34.6	49.5	25.1	36.4	378	378	189	283	1228	144	156	102	111	513
Compost for P	15.5	10.3	5.3	2.9	168 (84)	78 (102)	40 (111)	22 (123)	728	64	32	21	9‡	126
Compost for N/2 yr	69.4	–	75.2	–	756	–	567	–	1323	289	–	305	–	594
Compost for P/2 yr	31.0	–	15.9	–	337 (16)	–(84)	120 (94)	–(110)	761	129	–	64	–	193
Fertilizer	–	–	–	–	151	151	151	151	604	26	26	26	26	104
Check	0	0	0	0	0	0	0	0	0	0	0	0	0	0

† Numbers in parentheses are kg N ha⁻¹ applied as NO₃NH₄ fertilizer.

‡ Phosphorus application rates for annual P-based manure and compost were about 7 kg ha⁻¹ less than the expected plant P uptake (26 kg ha⁻¹).

organic C content of 1.8 g kg⁻¹ in the top 15 cm before the initiation of the experiment in 1992.

The experimental design was a randomized complete block with four replications. The 10 treatments applied included annual or biennial manure or compost application based on N or P removal by corn (151 kg N ha⁻¹ and 25.8 kg P ha⁻¹ for an expected yield level of 9.4 Mg ha⁻¹) and fertilized and unfertilized checks (Table 1). Fertilizer was applied in the spring each year from 1993 to 1996. The inorganic fertilizer plots received N as NH₄NO₃ (34–0–0, N–P–K) and P as superphosphate (0–20–0, N–P–K) in 1993 and diammonium phosphate (18–20–0, N–P–K) in 1994, 1995, and 1996. If necessary, the P-based treatments (annual or biennial application) were supplemented with N fertilizer as NH₄NO₃ in the spring so that a total of 151 kg N ha⁻¹ was available to the crop (Table 1). Biennial manure or compost applications were made to provide 151 kg N ha⁻¹ for N-based and 25.8 kg P ha⁻¹ for P-based rates in the second year after application. Nitrogen and P were overapplied in the first year of application for the biennial manure and compost treatments.

Manure or compost was hand-applied to plots 12.2 m long and 4.6 m wide (six corn rows) in late autumn after corn harvest. The P application rates in 4 yr of applications are given in Table 1. Manure and compost were applied and incorporated into the top 10 cm soil by disking within 2 d after application. The experimental area was disked to 10-cm soil depth in the spring before planting. Additional information regarding manure and compost N and P availability assumptions, corn yield, N and P uptake, and soil properties during the application years from 1993 to 1996 is reported in Eghball and Power (1999) and Eghball (2002).

Residual Effects

For the 1997, 1998, and 1999 growing seasons, no manure, compost, or fertilizer (except N application for the fertilizer treatment only in 1999) was applied. The experiment was under linear-move pivot sprinkler irrigation from 1997 to 1999. The amount of water applied was 30.7 cm in 1997, 8.9 cm in 1998, and 17.0 cm in 1999. Lower amount of irrigation water was applied in 1998 than the other years because of a problem with the pivot system early in the growing season. In all 7 yr, corn (Pioneer brand hybrid '3394') was planted at a rate of 47 000 seeds ha⁻¹ at a row spacing of 0.76 m. Weed control was achieved by band application of herbicide in the corn rows at planting and by cultivation. Four randomly chosen plants were harvested at tasseling for plant biomass and N content measurements in 1997 and 1998. The middle two rows of corn (6.1 m long) were harvested, and grain yield was measured. Stover was harvested from one row 6.1 m long. Grain yields were adjusted to a water content of 155 g kg⁻¹,

and stover yield is reported on oven-dry (60°C)-weight basis. Plots were harvested on 7 Oct. 1997 and 1998 and on 8 Sept. 1999.

Soil samples at depth increments of 0 to 15, 15 to 30, 30 to 45, 45 to 60, and 60 to 90 cm were collected from all plots shortly after corn harvest in the autumn. Soil samples were air-dried, ground to pass a 1-mm sieve, and analyzed for selected soil parameters. Soil pH and electrical conductivity (EC) were determined on 1:1 soil/water ratio (Smith and Doran, 1996). Plant and soil total N and C were determined by dry combustion as reported by Schepers et al. (1989). Soil bulk density on this site was not significantly influenced by the 10 treatments used in the application years (Eghball, 2002). Therefore, soil bulk density from 1996 was used to calculate the total N and C quantities for each plot. Soil NO₃-N and NH₄-N concentrations were determined on samples extracted by KCl and using a Lachat system (Zellweger Analytics, Milwaukee, WI). Phosphorus was analyzed by the Bray and Kurtz no. 1 soil P test method.

Analysis of variance was used to determine differences among treatments across years using SAS PROC MIXED procedure (Littell et al., 1996). In these analyses, year and soil depth increments were considered repeated observations. A probability level ≤0.05 was considered significant.

RESULTS AND DISCUSSION

Biomass and Nitrogen Uptake at Tasseling

Plant weight at tasseling in 1997 was significantly greater for the annual P-based manure, N- and P-based compost, and commercial fertilizer treatments than those for the check plots (Table 2). The biennial manure or compost application did not result in greater biomass than the check plots (Table 2) as the last biennial applications were made in the autumn of 1994 for the 1995 and 1996 corn growing seasons. The last application time for the annual manure, compost, and fertilizer treatments was in the autumn of 1995 for 1996 corn growing season. These data indicate that the positive residual effects of N- and P-based manure and compost applications and N fertilizer on plant biomass at tasseling were still evident for one growing season after the intended crop. By 1998, however, the residual effects of manure, compost, and fertilizer applications did not increase plant biomass at tasseling even though plant N content was higher for the biennial P-based manure treatment than the check (Table 2). Plant N uptake in 1997, in general, followed a pattern similar to that for plant biomass (Table 2).

Table 2. Residual effects of beef cattle manure or composted manure application on plant weight and N uptake at tasseling in 2 yr in eastern Nebraska.

Treatment†	Dry matter		N content	
	1997	1998	1997	1998
	g plant ⁻¹			
Manure for N	143	126	1.92	1.33
Manure for P	163	118	2.29	1.13
Manure for N/2y	149	129	2.09	1.39
Manure for P/2y	154	140	2.23	1.57
Compost for N	161	126	2.25	1.37
Compost for P	167	130	2.33	1.35
Compost for N/2y	149	141	2.06	1.50
Compost for P/2y	157	122	2.13	1.21
Fertilizer‡	164	121	2.39	1.34
Check	136	116	1.59	1.19
LSD _{0.05}	22	NS§	0.61	0.35
CV, %	12.0	20.9	22.1	23.7

† N and P indicate applications to meet N or P needs of corn, respectively, and 2y indicates biennial application. Manure and compost were applied from 1992 to 1995.

‡ In 1999 only, N was applied to the fertilizer treatment only at a rate of 151 kg ha⁻¹.

§ NS = nonsignificant.

Grain and Stover Yields and Total Nitrogen Uptake

Residual effects of annual or biennial N- or P-based manure, compost, and fertilizer treatments resulted in greater grain yield than the check plots in 1997 (Table 3). The biennial treatments had similar plant biomass to the check plots at tasseling (Table 2), but by harvest time, manure and compost treatments resulted in greater grain yield than the check treatment (Table 3). It seems that N mineralization after tasseling contributed to increased yield for the biennial applications even though the applications were made about 3 yr earlier (Eghball, 2000). By 1998, all treatments, except biennial P-based manure, had grain yields similar to the check plots (Table 3). Lower grain yield in 1998 than the other years reflects less irrigation water applied in 1998. In 1999, when only the fertilizer treatment received NH₄NO₃, greater grain yield was found for the fertilizer than other treatments except the biennial N-based compost treat-

Table 3. Residual effects of beef cattle manure or composted manure application on corn grain yield, stover yield, and total N uptake in three years in eastern Nebraska.

Treatment†	Grain yield‡			Stover yield			Total N uptake		
	1997	1998	1999	1997	1998	1999	1997	1998	1999
	Mg ha ⁻¹			kg ha ⁻¹					
Manure for N	9.49	6.58	9.30	6.62	3.78	5.11	161	109	157
Manure for P	9.50	6.10	9.48	6.53	3.50	5.31	171	93	148
Manure for N/2y	9.88	6.81	9.00	6.77	3.55	5.66	184	107	147
Manure for P/2y	9.92	7.03	9.22	5.86	4.13	5.70	172	114	158
Compost for N	9.87	6.73	8.93	6.59	3.66	5.65	167	102	146
Compost for P	10.25	6.62	9.40	6.95	3.82	5.17	186	100	151
Compost for N/2y	10.20	6.73	9.73	6.56	3.85	5.40	178	106	159
Compost for P/2y	9.57	6.01	9.44	6.36	3.60	6.26	166	87	153
Fertilizer§	9.90	6.58	10.64	5.91	4.24	6.71	178	113	184
Check	7.08	5.63	8.85	5.54	3.41	5.29	108	91	142
LSD _{0.05}	0.97	1.25	0.94	1.11	0.78	1.09	23	26	19
CV, %	13.5	14.6	9.4	12.9	16.8	16.5	16.4	17.8	11.9

† N and P indicate applications to meet N or P needs of corn, respectively, and 2y indicates biennial application. Manure and compost were applied from 1992 to 1995.

‡ Grain yield at 155 g kg⁻¹ water content.

§ In 1999 only, N was applied to the fertilizer treatment only at a rate of 151 kg ha⁻¹.

ment. Residual effects of manure treatments resulted in corn grain yields that were 85 to 89% of that for the applied N fertilizer treatment while they were 84 to 89% for the compost treatments in 1999 (Table 3). The check plot produced a corn yield of 8.85 Mg ha⁻¹ (141 bu ac⁻¹) in 1999 even though it had not received any treatment since 1992. The higher organic matter in this Mollisol ($\approx 3.5\%$) seemed to have contributed enough N to produce 8.85 Mg ha⁻¹ corn yield, which was 83% of that for the applied fertilizer treatment.

Stover yield was not different among most of the treatments in 1997 (Table 3). It seems that higher corn yields resulting from the residual effects of manure, compost, and N fertilizer did not result in greater stover yields than the check, except biennial N-based manure and annual P-based compost treatments. In 1998 and 1999, only the fertilizer treatment produced greater amount of stover than the check plots (Table 3). The fertilizer treatment received N fertilizer in 1999 and was expected to have greater stover yield than the check treatment. Averaged across treatments, stover/grain yield ratio was 0.67 in 1997, 0.58 in 1998, and 0.60 in 1999. There was damage from corn rootworm (*Diabrotica barberi*) in 1998, and that might have contributed to reduced grain and stover yields.

Total N uptakes for all treatments were greater than the check plots in 1997 (Table 3). By 1998, however, all treatments had total N uptake similar to the check plots. As expected, the fertilizer treatment resulted in greater N uptake than the check plots in 1999 (Table 3).

Soil Properties

Averaged across treatments, soil surface (0 to 15 cm) pH decreased from 1997 to 1999 (Table 4). Eghball (1999)

Table 4. Residual effects of beef cattle manure or composted manure application on soil electrical conductivity (EC), pH, total C and N concentrations, and quantities at 0- to 15-cm soil depth in eastern Nebraska.

Variable	EC	pH	Total C	Total N	Total C	Total N
	dS m ⁻¹		g kg ⁻¹		Mg ha ⁻¹	
Year						
1997	0.263	6.58	20.7	1.83	40.29	3.55
1998	0.340	6.43	21.1	1.86	41.03	3.61
1999	0.334	6.30	20.3	1.81	39.34	3.49
LSD _{0.05}	0.011	0.04	0.4	0.04	0.81	0.08
Treatment†						
Manure for N	0.323	6.50	21.6	1.89	39.75	3.48
Manure for P	0.287	6.34	19.7	1.73	37.49	3.29
Manure for N/2y	0.323	6.53	21.3	1.90	40.41	3.60
Manure for P/2y	0.318	6.40	22.2	1.97	43.56	3.85
Compost for N	0.357	6.75	21.0	1.88	41.20	3.70
Compost for P	0.299	6.33	19.9	1.76	41.30	3.61
Compost for N/2y	0.353	6.70	22.2	2.05	41.68	3.84
Compost for P/2y	0.296	6.35	20.1	1.78	39.20	3.46
Fertilizer‡	0.277	6.02	19.9	1.71	39.94	3.43
Check	0.294	6.43	19.3	1.66	37.67	3.23
LSD _{0.05}	0.026	0.17	NS§	0.24	NS	NS
Analysis of variance	P > F					
Year	0.0001	0.0001	0.0003	0.0573	0.0005	0.0197
Treatment	0.0001	0.0001	0.5524	0.0507	0.9810	0.8114
Year × treatment	0.2456	0.6177	0.4559	0.3078	0.5192	0.2468

† N and P indicate applications to meet N or P needs of corn, respectively, and 2y indicates biennial application. Manure and compost were applied from 1992 to 1995.

‡ In 1999 only, N was applied to the fertilizer treatment only at a rate of 151 kg ha⁻¹.

§ NS = nonsignificant.

found that manure and composted manure provide a liming effect on soil since the cattle diet contains lime, which is subsequently excreted in manure. The liming effect seems to diminish with time after termination of manure and compost applications. There was a liming effect with both manure and compost when compared with commercial fertilizer application (Table 4). Across 3 yr, the N-based application of manure or compost resulted in higher soil surface pH than the P-based treatments, indicating the combination effects of greater liming influence of higher application rates associated with N-based treatments and the acidifying effect of supplemental commercial fertilizer added with the P-based treatments (Table 1). Soil surface EC was also higher for the residual effect of N-based than the P-based manure and compost applications (Table 4). The higher EC for the N-based treatments reflects the larger amounts of manure and compost applied.

Total C and N concentrations and quantities in the 0- to 15-cm soil were highest in 1998 and decreased in 1999 (Table 4). There were residual effects of manure and compost applications on soil N concentration ($P = 0.0507$; Table 4). The biennial N-based compost and P-based manure treatments had more soil N content than the check or the fertilizer treatments. The residual effects of annual N- or P-based treatments were not different from those of the fertilizer or check treatments. Total C and N quantities were not different among treatments, indicating that total C and N were not sensitive indicators of the C and N pools in this high organic matter Mollisol beyond the application years. However, Ginting et al. (2003) showed that 4 yr after applications had ceased, soil potentially mineralizable N and microbial biomass C were greater for manure and compost applications than those for the control or inorganic fertilizer treatment.

Similar to the 0- to 15-cm soil depth, pH in the 15- to 30-cm soil depth decreased with time when averaged across all treatments (Table 5). No residual effects of the treatments on EC, pH, and total C and N concentrations and quantities were observed in the 15- to 30-cm soil depth (Table 5). Manure was incorporated into the top 10 cm of soil, and it appears that the movement of salt, lime, C, and N did not occur to the 15- to 30-cm soil depth. During the application years (1992–1995), no difference among the treatments for soil properties was observed, except EC, at the 15- to 30-cm soil depth increment (Eghball, 2002).

Soil $\text{NO}_3\text{-N}$ and $\text{NH}_4\text{-N}$ concentrations were different among the soil depth increments, with the concentrations being generally higher in the surface 15 cm than deeper soil increments (Table 6). Across soil depths, $\text{NO}_3\text{-N}$ concentrations after corn harvest were higher for the annual N-based manure, biennial P-based manure, and biennial N-based compost treatments than the check plots (Table 6). By 1999, the fertilizer treatment, which received NH_4NO_3 fertilizer, had higher $\text{NO}_3\text{-N}$ concentration than the other treatments. Soil $\text{NH}_4\text{-N}$ concentrations were similar among treatments in all three residual years. Residual values of manure and compost application resulted in increased soil $\text{NO}_3\text{-N}$ for 3 yr.

Table 5. Residual effects of beef cattle manure or composted manure application on soil electrical conductivity (EC), pH, total C and N concentrations and quantities at 15- to 30-cm soil depth in eastern Nebraska.

Variable	EC	pH	Total C	Total N	Total C	Total N
	dS m ⁻¹		g kg ⁻¹		Mg ha ⁻¹	
Year						
1997	0.233	6.44	19.4	1.58	41.17	3.34
1998	0.340	6.38	18.7	1.58	39.81	3.35
1999	0.262	6.32	19.0	1.60	40.38	3.39
LSD _{0.05}	0.013	0.04	0.5	NS†	1.04	NS
Treatment‡						
Manure for N	0.272	6.33	19.3	1.59	40.78	3.37
Manure for P	0.267	6.40	18.5	1.56	38.50	3.23
Manure for N/2y	0.285	6.39	18.8	1.60	39.31	3.35
Manure for P/2y	0.283	6.31	20.6	1.67	43.51	3.54
Compost for N	0.285	6.59	17.6	1.52	36.81	3.17
Compost for P	0.279	6.40	18.6	1.55	41.13	3.38
Compost for N/2y	0.301	6.39	19.5	1.63	42.18	3.52
Compost for P/2y	0.260	6.30	19.0	1.57	41.77	3.39
Fertilizer§	0.283	6.34	19.4	1.59	40.65	3.34
Check	0.268	6.36	19.2	1.60	39.82	3.31
LSD _{0.05}	NS	NS	NS	NS	NS	NS
Analysis of variance	$P > F$					
Year	0.0001	0.0001	0.0058	0.4410	0.0384	0.4737
Treatment	0.3151	0.2416	0.8867	0.9517	0.9544	0.9867
Year × treatment	0.7365	0.2947	0.6546	0.9765	0.6936	0.9567

† NS = nonsignificant.

‡ N and P indicate applications to meet N or P needs of corn, respectively, and 2y indicates biennial application. Manure and compost were applied from 1992 to 1995.

§ In 1999 only, N was applied to the fertilizer treatment only at a rate of 151 kg ha⁻¹.

Soil test P levels from 1996 to 1999 in the 0- to 15- and 15- to 30-cm soil depth increments were reported by Eghball et al. (2003). Soil test P levels in 1999 (Table 7) can be used to determine P leaching 4 yr after manure

Table 6. Residual effects of beef cattle manure or composted manure application on soil nitrate and ammonium concentrations in eastern Nebraska.

Variable	1997		1998		1999	
	$\text{NO}_3\text{-N}$	$\text{NH}_4\text{-N}$	$\text{NO}_3\text{-N}$	$\text{NH}_4\text{-N}$	$\text{NO}_3\text{-N}$	$\text{NH}_4\text{-N}$
	mg kg ⁻¹					
Depth (m)						
0.00–0.15	1.72	2.32	3.09	2.59	5.64	1.51
0.15–0.30	1.47	1.88	4.71	1.94	2.23	1.29
0.30–0.60	1.90	1.38	1.47	1.50	1.28	1.12
0.60–0.90	1.16	1.42	–	–	0.63	1.36
0.90–1.20	–	–	–	–	0.85	2.02
LSD _{0.05}	0.28	0.29	0.59	0.29	0.41	0.13
Treatment‡						
Manure for N	1.92	1.59	4.00	2.05	1.53	1.38
Manure for P	1.37	1.81	2.43	2.18	1.87	1.53
Manure for N/2y	1.92	1.74	3.39	2.02	1.89	1.61
Manure for P/2y	2.71	1.79	3.65	2.05	1.80	1.34
Compost for N	0.88	1.61	2.97	1.66	2.01	1.20
Compost for P	1.18	1.66	2.64	2.09	1.78	1.51
Compost for N/2y	2.38	1.82	4.79	2.09	2.49	1.30
Compost for P/2y	1.20	1.89	2.38	2.21	1.44	1.55
Fertilizer‡	1.44	1.79	2.77	2.04	3.70	1.31
Check	0.61	1.81	1.90	1.73	1.36	1.33
LSD _{0.05}	1.21	NS§	1.74	NS	1.18	NS
Analysis of variance	$P > F$					
Depth	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
Treatment	0.0307	0.9647	0.0637	0.6287	0.0165	0.9913
Depth × treatment	0.1180	0.9981	0.9246	0.8939	0.5264	0.9567

‡ N and P indicate applications to meet N or P needs of corn, respectively, and 2y indicates biennial application. Manure and compost were applied from 1992 to 1995.

§ In 1999 only, N was applied to the fertilizer treatment only at a rate of 151 kg ha⁻¹.

§ NS = nonsignificant.

Table 7. Residual effects of beef cattle manure or composted manure application on soil test P (Bray and Kurtz no. 1) at various depth increments in 1999 in eastern Nebraska.

Treatment†	Soil depth (cm)				
	0-15	15-30	30-45	45-60	60-90
	mg kg ⁻¹				
Manure for N	77.5	22.0	11.8	10.2	23.9
Manure for P	50.3	21.4	11.8	8.8	15.9
Manure for N/2y	90.7	34.0	13.5	8.9	18.8
Manure for P/2y	69.7	21.5	12.1	6.9	12.9
Compost for N	109.1	34.8	14.2	10.6	20.0
Compost for P	58.0	24.4	11.9	8.8	22.1
Compost for N/2y	167.6	49.7	18.3	9.0	19.1
Compost for P/2y	58.2	42.3	12.0	6.9	15.5
Fertilizer‡	55.1	19.5	9.7	6.3	13.7
Check	49.2	27.1	11.0	7.3	17.8
LSD _{0.05}	25.9	17.5	3.0	3.1	NS§

† N and P indicate applications to meet N or P needs of corn, respectively, and 2y indicates biennial application. Manure and compost were applied from 1992 to 1995.

‡ In 1999 only, N was applied to the fertilizer treatment only at a rate of 151 kg ha⁻¹.

§ NS = nonsignificant.

or compost application had ceased. Soil P levels in 1999 were different among treatments in all soil depth increments except in the 60- to 90-cm depth increment (Table 7). This indicates leaching of P from applied manure and compost to the 45- to 60-cm soil depth increment. Eghball (2003) showed P leaching to the soil depth increment of 15 to 30 cm in 1996 when manure or compost application ceased. By 1999, however, P had leached deeper into the soil. In the 0- to 15-cm soil depth increment, the N-based treatments had higher soil P levels than the check, fertilizer, and P-based treatments (Table 7). At the 45- to 60-cm soil depth increment, annual N-based manure and compost applications had greater soil P levels than the fertilizer treatment. The biennial N-based compost treatment had higher soil P levels than the fertilizer or check treatments in the 15- to 30- and 30- to 45-cm soil depth increments. A greater amount of P was applied for the biennial N-based compost treatment (Table 1), resulting in deeper leaching of P in the soil profile. The amount of P applied was higher for the N-based compost than manure treatments since only 20% of compost N was considered plant available in the first year after application compared with 40% for noncomposted manure when applications were made.

The elevated plant available P concentration in the soil surface (0 to 15 cm) as a result of manure or compost application can remain for several years. This has both agronomic and environmental implications since it can contribute to crop P uptake and also can be available for transport by runoff. Eghball et al. (2003) found that 10 yr of crop removal was needed to reduce soil P level from 265 mg kg⁻¹ to the original 69 mg kg⁻¹ that existed before compost application. How long soil P continues to contribute to crop P uptake depends on soil characteristics. Sharpley (1996) found that the rate of P release from high-P soils decreased more rapidly as soil P sorption saturation increased and P sorption maxima decreased. Even though positive residual effects of manure or compost application on crop production were observed for at least 1 yr, P contribution to plant uptake could last much longer than 1 yr. That can be significant

in soils that are entirely P deficient or have P-deficient areas within the field.

CONCLUSIONS

Residual effects of N- and P-based manure and compost applications on corn yield and N uptake can last for at least one growing season. Residual effects of N-based manure and compost applications on corn production were greater than the P-based treatments since the amounts of manure or compost applied were greater for N- than P-based management systems. Residual effects of manure and compost applications on soil properties were longer lasting than those influencing corn production. Soil P, NO₃-N, EC, and pH levels were greater for the N-based manure or compost application than the check treatment 4 yr after the last applications were made. Phosphorus leaching to a soil depth increment of 15 to 30 cm was observed in 1996, 1 yr after the last manure and compost applications. By 1999 however, P from N-based manure or compost treatments had leached to the 45- to 60-cm soil depth increment. Leached P can reach the ground water when ground water is close to the soil surface or excess P is in soils with fluctuating ground water. Both corn production and soil properties were improved by the residual values of applied manure or compost. Phosphorus-based application was environmentally sound since it provided nutrients for the crop while maintaining the soil P level similar to the untreated check. Applications of manure and compost not only improved soil properties for several years after applications had ceased but also provided nutrients and liming effects for the growing corn.

REFERENCES

- Binford, G.D., G.W. Hergert, and J.M. Blumenthal. 2000. Millet. p. 135-137. In R.B. Ferguson and K.M. De Groot (ed.) Nutrient management for agronomic crops in Nebraska. Univ. of Nebraska Coop. Ext., Lincoln.
- Chang, C., T.G. Sommerfeldt, and T. Entz. 1991. Soil chemistry after eleven annual applications of cattle feedlot manure. *J. Environ. Qual.* 20:475-480.
- Eghball, B. 1999. Liming effects of beef cattle feedlot manure or compost. *Commun. Soil Sci. Plant Anal.* 30:2563-2570.
- Eghball, B. 2000. Nitrogen mineralization from field-applied beef cattle feedlot manure or compost. *Soil Sci. Soc. Am. J.* 64:2024-2030.
- Eghball, B. 2002. Soil properties as influenced by phosphorus- and nitrogen-based manure and compost applications. *Agron. J.* 94: 128-135.
- Eghball, B. 2003. Leaching of phosphorus fractions following manure and compost applications. *Commun. Soil Sci. Plant Anal.* 34:2803-2815.
- Eghball, B., and J.F. Power. 1999. Phosphorus and nitrogen-based manure and compost application: Corn production and soil phosphorus. *Soil Sci. Soc. Am. J.* 63:895-901.
- Eghball, B., J.F. Shanahan, G.E. Varvel, and J.E. Gilley. 2003. Reduction of high soil test phosphorus by corn and soybean varieties. *Agron. J.* 95:1233-1239.
- Eghball, B., B.J. Wienhold, J.E. Gilley, and R.A. Eigenberg. 2002. Mineralization of manure nutrients. *J. Soil Water Conserv.* 57: 470-473.
- Ginting, D., A. Kessavalou, B. Eghball, and J.W. Doran. 2003. Greenhouse gas emissions and soil indicators four years after manure and compost applications. *J. Environ. Qual.* 32:23-32.
- Littell, R.C., G.A. Milliken, W.W. Stroup, and R.D. Wolfinger. 1996. SAS system for mixed models. SAS Inst., Cary, NC.

- Lund, Z.F., and B.D. Doss. 1980. Residual effects of dairy cattle manure on plant growth and soil properties. *Agron. J.* 72:123–130.
- Motavalli, P.P., K.A. Kelling, and J.C. Converse. 1989. First-year nutrient availability from injected dairy manure. *J. Environ. Qual.* 18: 180–185.
- Mugwira, L.M. 1979. Residual effects of dairy manure on millet and rye forage and soil properties. *J. Environ. Qual.* 8:251–255.
- Schepers, J.S., D.D. Francis, and M.T. Thompson. 1989. Simultaneous determination of total C, total N, and ¹⁵N on soil and plant material. *Commun. Soil Sci. Plant Anal.* 20:949–959.
- Sharpley, A.N. 1996. Availability of residual phosphorus in manured soils. *Soil Sci. Soc. Am. J.* 60:1459–1466.
- Smith, J.L., and J.W. Doran. 1996. Measurement and use of pH and electrical conductivity for soil quality analysis. p. 169–185. *In* J.W. Doran and A.J. Jones (ed.) *Methods of assessing soil quality*. SSSA Spec. Publ. 49. SSSA, Madison, WI.
- Wallingford, G.W., L.S. Murphy, W.L. Powers, and H.L. Manges. 1975. Disposal of beef-feedlot manure: Effects of residual and yearly applications on corn and soil chemical properties. *J. Environ. Qual.* 4:526–531.