University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Proceedings of the Thirteenth Vertebrate Pest Conference (1988)

Vertebrate Pest Conference Proceedings collection

3-1-1988

FIELD EVALUATION OF QUINTOX (CHOLECALCIFEROL) FOR CONTROLLING COM-MENSAL RODENTS

L. Dawn Brown Bell Laboratories, Inc., Madison, Wisconsin

E. F. Marshall Bell Laboratories, Inc., Madison, Wisconsin

Follow this and additional works at: http://digitalcommons.unl.edu/vpcthirteen



Part of the Environmental Health and Protection Commons

Brown, L. Dawn and Marshall, E. F., "FIELD EVALUATION OF QUINTOX (CHOLECALCIFEROL) FOR CONTROLLING COM¬MENSAL RODENTS" (1988). Proceedings of the Thirteenth Vertebrate Pest Conference (1988). Paper 15. http://digitalcommons.unl.edu/vpcthirteen/15

This Article is brought to you for free and open access by the Vertebrate Pest Conference Proceedings collection at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Proceedings of the Thirteenth Vertebrate Pest Conference (1988) by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

FIELD EVALUATION OF QUINTOX (CHOLECALCIFEROL) FOR CONTROLLING COMMENSAL RODENTS

L. DAWN BROWN and E.F. MARSHALL, Bell Laboratories, Inc., Madison, Wisconsin 53704.

AB STRACT: Field efficacy studies were performed on Norway rats (Rattus norvegicus). roof rats (Rattus rattus). and house mice (Mus musculus) under a variety of conditions throughout the continental United States. Baits containing .075% (750 ppm) cholecalciferol yielded exceptional results in reducing commensal rodent populations under actual field conditions.

Proc. Vertebr. Pest Conf. (A.C. Crabb and R.E. Marsh, Eds.), Printed at Univ. of Calif., Davis. 13:70-74, 1988

INTRODUCTION

In the late seventies, Bell Laboratories, Inc. applied for registration of a rodenticide containing a totally new active compound, cholecalciferol. Bell was interested in this compound because of its different mode of action which was becoming increasingly more important in light of the occurrence of certain anticoagulant resistance. Additionally, cholecalciferol had a unique stop-feed action and low secondary hazards, making it quite appealing for use as a rodenticide. Laboratory trials showed QUINTOX to be highly effective on all three types of commensal rodents: Norway rat, roof rat, and house mouse. Seven to fourteen grams of QUINTOX bait was found to be sufficient to kill an adult rat, while 2-2.8 grams of QUINTOX would kill a mouse.

A Conditional Registration was granted by the U.S. Environmental Protection Agency to Bell Laboratories, Inc., for QUINTOX Rat & Mouse Bait (0.075% cholecalciferol) in November 1984. The issuance of U.S. clearance of several forms of QUINTOX (i.e., bait, place packs, and seeds) marked the culmination of seven years of research and development by Bell, but fell short of completing extensive field research (i.e., product performance data of cholecalciferol required by the EPA before a nonconditional registration could be granted). It should be noted that many field and laboratory studies were conducted on product performance; however, the EPA felt that though there was a need for a nontraditional chemical method of rodent control, the product needed to further prove itself in the field.

Starting in early 1985, an extensive Field Efficacy Testing Program was instituted in the five geographical areas of the United States, testing the effectiveness of QUINTOX on Norway rats, roof rats, and house mice under field conditions, both indoors and out, following EPA approved testing guidelines.

FIELD METHODS AND DESIGN

A field test protocol was developed for the efficacy testing of QUINTOX Rat and Mouse Bait. It was designed to be a practical flexible approach to studying the effectiveness of the bait in natural habitats.

For each species of rodent the following minimum regional studies were required: Norway rat-five indoor studies (one in each of the five major regions of the country),

and two outdoor studies (each in a different region); roof rattwo indoor studies (in different regions) and one outdoor study; house mouse-five indoor studies (one in each region) and one outdoor study.

An indoor study was defined as a study conducted in enclosed structures such as hotels, houses, barns, etc. Outdoor studies were conducted in sites comparable to the following: sheds (covered, non-enclosed structures); small dumps not being continually reinfested; small isolated stock yards (all placements in tamper-proof bait boxes or in burrows).

Field trials were conducted utilizing QUINTOX Rat and Mouse Bait (EPA Reg. No. 12455-39) which contained 0.075% cholecalciferol in a propriety grain base formula. (In most instances QUINTOX was placed in covered bait stations; however, place packs were used in some situations.) Bait was replenished as necessary.

Each test was set up with both a pre- and post-test census period of 3-5 days, a 1-3 day break period between the census period and test period, and a 10-20 day treatment (test) period. The test period may be extended if necessary.

3-5 d per. 1-3 d per. 10-20 d per. 1-3 d per. 3-5 d per. pre-test break test break post-test census census

The census methods included tracking powder patches, live trapping, sightings, feed consumption, scat counts, and a variety of other methods.

In all but two field trials, two census techniques were utilized in both the pre- and post-test census evaluation. At least a one- to three-day period was allowed between the census periods and the treatment period to avoid conditioning effects. When feed consumption was utilized as a census technique, care was taken to ensure that the test baits were not placed in the same area where non-toxic food was previously placed. QUINTOX bait was usually made available until all signs of feeding stopped. Reductions in rodent population were calculated based on the population differences found between the pre-test and post-test census.

RESULTS

Field test data indicated the OUINTOX Rat and Mouse

Bait was effective in controlling Norway rats, roof rats, and house mice under a variety of field conditions and geographical locations (see Table 1 and Figures 1,2, and 3). The raw data are found in Tables 2, 3, and 4. QUINTOX was well accepted, even when competitive foods were available. No bait shyness or aversion was noted in any of the field trials.

Species	Period of Treatment (days)		Popn. Reduction (avg.)
	Range	Mean	
Norway rat	3-34	13,4	91.9%
Roof rat	6-14	8.8	89,8%
House mouse	7-23	13.1	98.0%

Fig. 3. Quintox field efficacy testing program (indoor/outdoor results).




Table 1. Summary of QUINTOX efficacy results.




Fig. 1. Quintox field efficacy testing program geographical breakdown.




Fig. 2. Quintox field efficacy testing program overall results.

QUINTOX treatments ranged from 3-34 days with a mean of 13.4 days for Norway rats and an average reduction of 91.9%; from 6-14daysforroofratswithamean of 8.8 days and an average reduction of 89.8%; and 7-23 days for house mice with a mean of 13.1 days and an average of 98% reduction. Field tests received 90% or greater reduction in each of the five regions.

DISCUSSION

QUINTOX Rat and Mouse Bait containing cholecalciferol was given a rigorous series of indoor and outdoor tests in five different geographical regions of the U.S. on three species of commensal rodents; Norway rat, roof rat, and house mouse.

Five different regions of testing were chosen to ensure that QUINTOX would be effective under a variety of climate conditions. Both indoor and outdoor studies were conducted to challenge QUINTOX in a wide array of the difficult baiting situations that face today's pest control operator.

Under all circumstances, QUINTOX Rat and Mouse Bait did exceptionally well. Rodent population reduction control was very good in each of the five regions tested not only indoors, but also in the often harder-to-control outdoor settings.

QUINTOX surpassed the minimum of 70% population reduction in every trial.

CONCLUSION

0.075% cholecalciferol was shown to be an effective toxicant for commensal rodent control (Marshall 1982). With the conclusion of successful field trials 0.075% cholecalciferol in the QUINTOX formula has shown to yield excellent results under a variety of challenging field conditions. Based upon the final submission of field efficacy data, Unconditional Registration was granted for all formulas of

Table 2. Efficacy summary* of QUINTOX (0.075% cholecalciferol) against the Norway rat (Rattus norvegicus) Site Site Census % Census % Reduction Reduction Location Method #1 Method #2 Outdoor 82 Burrow activity City Block Maryland 100 100 Feed consumption Tracking patches Indoor Single Family Dwelling New Jersey Outdoor Burrow activity 100 Sightings 98 Hog Lot Tennessee 100 100 Outdoor Waste Transfer Station Burrow activity Sightings Tennessee Indoor 75 Burrow activity 75 Sightings Poultry Farm Louisiana Outdoor Hog Farm **Trapping** 90 Tracking patches 70 Nebraska Outdoor 100 100 Theme Park Burrow activity Scat count Illinois Outdoor Solid Waste Transfer Tracking patches 95 Feed consumption 95 Station, Michigan Outdoor Burrow activity 74 83 City Block Scat count Illinois Outdoor **Burrow** activity 100 Feed consumption 95 Restaurant Nebraska Outdoor Tracking patches 79 Trapping 100 Farm Nebraska Indoor Food Storage Tracking patches 100 Feed consumption 100 Nebraska Indoor Tracking patches 100 100 Feed consumption Barn Wisconsin Indoor Live trapping 87 Tracking patches 81 Farm House California Indoor Live trapping 88.9 Feed consumption 98 Poultry Farm

California

[■]Range 3-34 days, mean = 13.4 days, average % control = 91.9.

Table 3. Efficacy summary* of QUINTOX (0.075% cholecalciferol) against the roof rat (Rattus rattus).

Site	Site Location	Census Method #1	% Reduction	Census Method #2	% Reduction
Outdoor	Processing Plant California	Feed consumption	99	Tracking patches	99
Indoor	Farm House California	Live trapping	85	Tracking patches	75
Indoor	Restaurant Texas	Tracking patches	93	Sightings	67
Indoor	Hog Farrowing Facility Texas	Scat count	100	Tracking patches	100

^{*}Range 6-14 days, mean = 8.8 days, average & control = 89.8

Table 4. Efficacy summary* of QUINTOX (0.075% cholecalciferol) against the house mouse (Mus musculus).

Site	Site Location	Census Method #1	%. Reduction	Census Method #2	% Reduction
Outdoor	Restaurant New Jersey	Burrow activity	94	Scat count	100
Indoor	Office New Jersey	Tracking patches	100	Scat count	100
Indoor	Apt. Building New Jersey	Tracking patches	100	Scat count	100
Indoor	Office Tennessee	Sightings	100	Scat count	100
Indoor	Barn Wisconsin	Burrow activity	100	Scat count	100
Indoor	House Wisconsin	Tracking patches	100	Scat count	100
Indoor	Garage Wisconsin	Feed consumption	100	Tracking patches	100
Indoor	Egg Carton Storage Area, Indiana	Feed consumption	94.1	90	950
Indoor	Storage Bldg. California	Live trapping	100	Tracking patches	100
Outdoor	House Texas	Sightings	100	Scat count	100
Indoor	Feed storage California	Live trapping	71.4	Feed consumption	99
Indoor	Feed storage California	Live trapping	95.4	Feed consumption	100

[&]quot;Range 7-23 days, Mean=13.1 days, Average % control=98

QUINTOX by the Environmental Protection Agency on December 21,1987.

Further research of QUINTOX is warranted, as preliminary laboratory studies indicate that the bait is highly effective in controlling rock squirrels and pocket gophers.

ACKNOWLEDGMENTS

We wish to thank Rex E. Marsh, University of California, Davis; Rex O.Baker, California State Polytechnical University, Pomona; Dr. C. Douglas Mampe, DM Associates; Donna A. Johnson, City of Baltimore; Joseph K. Harris, Hill-Smith Termite & Pest Control; Dr. James F. Fowler, Louisiana Cooperative Extension Services; Dr. Robert M. Timm and Daryl D. Fisher, University of Nebraska-Lincoln; Patri-

cia Hottel, W. B. McCloud & Co.; Gene Meester, Presto-X Co.; Dr. Neil E. Pennington and Herb Zisser, Michigan Dept. of Public Health; John J. Halpin, City of Chicago; Dr. Rosalie P. Myers, Florida A&M University; Lee R. Martin, Wildlife Control Technology,; David Saunders, Herwald Pest Control; Dr. Theodore A. Granovsky, Granovsky Associates; Richard C. Berman, Waltham Chemical; Robert M. Corrigan, Purdue University and James Boren, Pest Control Services, Inc. for their assistance in developing field test data.

LITERATURE CITED

MARSHALL, E.F. 1982. Cholecalciferol, a promising new rodenticide. Proc. 10th Vert. Pest Conf. University of California, Davis.

