University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Related Publications from ANDRILL Affiliates | Antarctic Drilling Program

1-1-2007

Showing a strong link between climatic and pCO_2 changes: resolving discrepancies between oceanographic and Antarctic climate records for the Oligocene and early Miocene (34-16 Ma)

Stephan Pekar
Queens College, stephen.pekar@qc.cuny.edu

N. Christie-Blick
Columbia University, ncb@ldeo.columbia.edu

Follow this and additional works at: http://digitalcommons.unl.edu/andrillaffiliates

Part of the Environmental Indicators and Impact Assessment Commons

Pekar, Stephan and Christie-Blick, N., "Showing a strong link between climatic and pCO_2 changes: resolving discrepancies between oceanographic and Antarctic climate records for the Oligocene and early Miocene (34-16 Ma)" (2007). *Related Publications from ANDRILL Affiliates.* Paper 17.
http://digitalcommons.unl.edu/andrillaffiliates/17

This Article is brought to you for free and open access by the Antarctic Drilling Program at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Related Publications from ANDRILL Affiliates by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Showing a strong link between climatic and pCO_2 changes: resolving discrepancies between oceanographic and Antarctic climate records for the Oligocene and early Miocene (34-16 Ma)

S. F. Pekar,[1,2] and N. Christie-Blick[3]

[1] Queens College School of Earth and Environmental Sciences, CUNY, 65-30 Kissena Blvd., Flushing, New York 11367, USA
[2] Lamont-Doherty Earth Observatory of Columbia University, Palisades, New York 10964, USA (stephen.pekar@qc.cuny.edu)
[3] Department of Earth and Environmental Sciences and Lamont-Doherty Earth Observatory of Columbia University, Palisades, New York 10964, USA (ncb@ldeo.columbia.edu)

Summary An apparent mismatch between published oxygen isotope data and other paleoclimate proxies for the span from 26-16 Ma is resolved by calibration against eustatic estimates obtained from backstripped continental margin stratigraphy. Ice-volume estimates from calibrated oxygen isotope data compare favorably with stratigraphic and palynological data from Antarctica, and with estimates of atmospheric carbon dioxide for the early Oligocene through early Miocene (34-16 Ma). These isotopic data suggest that the East Antarctic Ice Sheet grew to as much as 30% greater than the present-day ice volume at glacial maxima. This conclusion is corroborated by seismic reflection and stratigraphic data from the Antarctic margin that suggest that the ice sheet may have covered much of the continental shelf at Oligocene and early Miocene glacial maxima. Palynological data suggest long-term cooling during the Oligocene, with near tundra environments developing along the coast at glacial minima by the late Oligocene.

Citiation: Pekar, S.F., N. Christie-Blick (2007), Showing a strong link between climatic and pCO_2 changes: resolving discrepancies between oceanographic and Antarctic climate records during the Oligocene and early Miocene (34-16 Ma), in Antarctica: A Keystone in a Changing World – Online Proceedings of the 10th ISAES, edited by A.K. Cooper and C.R. Raymond et al., USGS Open-File Report 2007-1047, Extended Abstract 024, 4 p.

Introduction

The history and mechanisms of late Paleogene and early Neogene (34-16 Ma) climate and ice-volume changes continue to be controversial, despite a plethora of new data, owing to apparent inconsistencies between available proxies. For example, distal proxies for Antarctic climate and ice volume, which include deep-sea oxygen isotopic data (Zachos et al., 2001; Coxall et al., 2005) and stratigraphic records of sea-level change at mid- to low-latitude continental margins (e.g., Kominz and Pekar, 2001; Miller et al., 2005) provide estimates for the size of the early Oligocene ice sheet that vary greatly, from as small as ~50% of the volume of the present-day East Antarctic Ice Sheet (EAIS; e.g., Zachos et al., 2001) to as large as twice that volume (Coxall et al., 2005).

Stratigraphic and palynological data acquired close to Antarctica (e.g., Cooper et al., 1991; Raine and Askin, 2001) suggest a heavily glaciated continent at Oligocene-early Miocene glacial maxima and gradual cooling between successive glacial minima, from cool temperate conditions in the early Oligocene to tundra-like conditions in the latest Oligocene to early Miocene. They are, however, puzzlingly inconsistent with $\delta^{18}O$ evidence (an abrupt decrease in $\delta^{18}O$ values) for global warming and a significant decrease in EAIS volume during the late Oligocene, persisting into the early Miocene (Zachos et al. 2001). The latter data have been taken to indicate a decoupling during that interval between climate change and atmospheric pCO_2 estimates based on $\delta^{13}C$ data from alkenones (Pagani et al., 2005), with pCO_2 values gradually decreasing during the Eocene to Miocene, and reaching near modern levels by the latest Oligocene.

Calibrations of isotopic data against stratigraphically constrained sea-level changes allow apparent conflicts between proxies to be resolved, with all proxies brought into alignment with alkenone $\delta^{13}C$ evidence for a secular decrease in atmospheric pCO_2 for the entire interval between 34 and 16 Ma (Pagani et al., 2005).

Calibration of isotopic records to glacioeustasy

Ice-volume and their associated sea-level changes were determined by calibrating detrended amplitudes of apparent sea level (ASL, defined as eustasy plus water loading effects) based on 2-D flexural backstripping of the stratigraphy recovered from boreholes (Kominz and Pekar, 2001; Pekar et al., 2002) drilled as part of the New Jersey Coastal Plain Drilling Program (e.g., Miller et al., 1998) to $\delta^{18}O$ amplitudes for Oi-events identified by Miller et al. (1991) and Pekar and Miller (1996) in deep-sea records. These calibrations range from as low as 0.12‰/10 m ASL for the Weddell Sea Site 690 to 0.35‰/10 m for Equatorial Atlantic Ocean Site 929, with good to excellent correlation (see Pekar et al., 2002; Pekar et al., 2006 for discussion). Differences among the calibrations are attributable to variability in deep-sea temperatures among the sites between glacial maxima and minima at the million-year time scale. Oligocene $\delta^{18}O$ values of 3‰ or greater in deep-sea records are consistent with an EAIS of modern size and with bottom water temperatures $\leq$ 2.0˚ C (see Pekar et al., 2006 for discussion).

Discussion: Synthesis of Antarctic cryospheric history between 34-16 Ma

Calibrated isotopic records

Calibrated isotopic records indicate that ice volume increased from near zero during the late Eocene to 25% larger than the present-day EAIS (equivalent to an ASL lowering of 75 m) by the earliest Oligocene (33.5 Ma; Fig. 1). Oligocene ice volume was greatest at ~29 and 25 Ma, being some 30% larger than the present-day EAIS (equivalent to 80 m of ASL lowering). The EAIS was generally smaller between 33 and 29 Ma (with the exception of isotope event Oi1b [31.7-31.4 Ma]) and between 26 and 23.2 Ma, with ice volume during glacial maxima close to 80% of the present-day EAIS. Our calibrated records also show that ice volume rarely decreased below 50% of the present-day EAIS during Oligocene glacial minima. Ice volume during most of the early Miocene (23-16 Ma) was comparable to that of the Oligocene, ranging from 50 to 130% of the present-day EAIS (Pekar and DeConto, 2006).

Evidence from Antarctica

Stratigraphic evidence from boreholes and marine seismic reflection data indicate that the EAIS extended across the shelf at glacial maxima since the earliest Oligocene. Seismic data indicate ice grounding on the shelf, in some cases approaching the shelf edge, as early as the early Oligocene near Prydz Bay (Cooper et al., 1991; Hambrey et al., 1991), and no later than the late Oligocene in the western and eastern parts of the Ross Sea (Bartek et al., 1997; Sorlien et al., in press; Fig. 1). Hiatuses recognized at ~34 Ma, ~29-25 Ma, and ~23 Ma in the Cape Roberts boreholes coincide with times of maximum ice volume based on calibrated isotopic records (Fig. 1).

Palynological and leaf data from boreholes located in the western Ross Sea indicate a gradual cooling during the Oligocene, ranging from cool temperate during the latest Eocene to cold temperate during the early Oligocene to near-tundra-like conditions associated with glacial minima in the late Oligocene (e.g., Raine and Askin, 2001). Near-tundra-like conditions continued into the early Miocene. These records are thought to be from sediments preferentially representing interglacial intervals. The match between the timing of hiatuses in Antarctic cores and times of maximum ice volume suggests that during glacial times, these locations would have been either subaerially exposed or covered by ice.

Resolving the apparent decoupling of isotopic records with other climate records and pCO_2 estimates

A comparison of these records to atmospheric pCO_2 estimates (Pagani et al., 2005) suggests a strong if nonlinear link between climate and pCO_2 levels during the Oligocene and early Miocene. A sharp increase in ice volume at 33.5 Ma (early Oligocene) was followed by a decrease to about 80% of the present-day EAIS at glacial maxima. Ice volume at glacial maxima then gradually increased to between 100 and 130% of present-day levels between "mid" Oligocene and early Miocene (28 and 17 Ma), with volumes generally not decreasing below 50% of modern levels at glacial minima. During this same interval, pCO_2 generally decreased, reaching near-modern levels by the late Oligocene, and then continued at these levels throughout the early Miocene (Pagani et al., 2005).

Calibration of deep-sea $\delta^{18}O$ records against independently measured apparent sea-level changes resolves the hypothesized conflict between oceanographic and Antarctic climate records and evidence for a decrease in pCO_2 during late Oligocene to early Miocene time (see Zachos et al., 2001; Pagani et al., 2005). The apparently abrupt decrease in $\delta^{18}O$ values in the late Oligocene is an artifact of the way in which the record was spliced together at ~25 Ma, with $\delta^{18}O$ records from mainly southern sites below and Atlantic sites above (Fig. 1) and interpreted at face value (Pekar et al., 2002; Pekar et al., 2006). Our results indicate that there was no abrupt warming, nor any significant deglaciation in Antarctica. Therefore, the paleoclimate record is not decoupled from pCO_2 estimates.

An explanation for the apparent decoupling between non-calibrated isotopic composite records and other climate proxies, as well as with pCO_2 estimates, is that a warmer (presumably more saline) bottom water mass increased spatially in the deep sea in a number of ocean basins during the late Oligocene and early Miocene (Pekar et al., 2006; Pekar and DeConto, 2006). This is supported by high $\delta^{18}O$ values from Weddell Sea ODP Sites 689 and 690 between the "mid" Oligocene and the top of the available record at 24.5 Ma compared with coeval lower latitude isotopic records from the Atlantic and Indian Ocean basins.

Summary

The apparent decoupling between deep-sea $\delta^{18}O$ records and pCO_2 estimates and stratigraphic and vegetation data from Antarctica during the late Oligocene and early Miocene is resolved by calibrating the isotopic records, resulting in a consistent view of cryospheric and climate change in Antarctica between 34 and 16 Ma. It is not necessary to infer that any decoupling between climate change and CO_2 levels occurred, either in the late Oligocene or early Miocene.

Acknowledgments

This research was supported in part by National Science Foundation grant (OCE 99-11121 to N. Christie-Blick and S.F. Pekar). We wish to thank the co-editor, W. Wise, for his suggestions.



Figure 1. Modified from Pekar and Christie-Blick, 2007. (A) Apparent sea-level (ASL) estimates are derived by applying $\delta^{18}O$ to ASL calibrations to $\delta^{18}O$ records (Pekar et al., 2002; Pekar et al., 2006). The upper x-axis is the percent of the present-day EAIS (equivalent to ~60 m ASL). The lower x-axis is ASL change, with zero representing present-day sea level. Good agreement exists between ice-volume estimates from each of the calibrated records, in spite of different calibrations being used for each record. Blue bars represent times when ice volume was ≥ than present-day EAIS based on calibrated isotopic records. (B) Brown bars represent preserved sediment and red wavy lines, hiatuses identified in cores from the Cape Roberts Project (based on Florindo et al., 2005). Note the excellent agreement between ice volume ≥ present-day EAIS and the timings of the hiatuses. (C) Biological data (Askin, 2001) from Cape Roberts Project that show evidence for long-term climate cooling and full glacial conditions in Antarctica. (D) First occurrence of grounded ice based on core and seismic data around Antarctica (Cooper et al., 1991; Hambrey et al., 1991; Bartek et al., 1997; Sorlien et al., 2007). (E) pCO_2 estimates from Pagani et al. (2005) show decreasing values during the Oligocene, reaching pre-industrial levels by the latest Oligocene and continuing into the early Miocene. The range shown is due to the uncertainty involved in using carbon isotopic composition of sedimentary alkenones. The dashed line represents pre-industrial values (280 ppm), while the shaded box represent values that are predicted for this century (Watson et al., 2001). (F) Deep-sea $\delta^{18}O$ composite modified from Zachos et al. (2001), and including Sites 803 (Barrera et al., 1993), 1090 (Billups et al., 2002), and 1218 (Lear et al., 2004). The abrupt $\delta^{18}O$ decrease at circa 24.5 Ma is due to a change in the source of data from high latitude to low latitude sites, with Southern Ocean sites below and mainly western equatorial Atlantic Site 929 above (Pekar et al. 2002; Pekar et al., 2006).

References

Askin, C.B., 2001. Glacial influence from clast features in Oligocene and Miocene strata cored in CRP-2A and CRP-3, Victoria Land Basin, Antarctica. Terra Antartica 7, 493-501.

Barrera, E., Baldauf, J., Lohmann, K.C., 1993. Strontium isotope and benthic foraminifer stable isotopic results from Oligocene sediments at Site 803, In: Berger, W.H., Kroenke, L.W., Mayer, L.A., (Ed.), Proceedings of the Ocean Drilling Program, Scientific Results 130, Ocean Drilling Program, College Station, pp. 269-279.

Bartek, L.R., Andersen, J.L.R., Oneacre, T.A., 1997. Substrate control on distribution of subglacial and glaciomarine seismic facies based on stochastic models of glacial seismic facies deposition on the Ross Sea continental margin, Antarctica. Marine Geology 143, 223-262.

Billups, K., Channell, J.E.T., Zachos, J., 2002. Late Oligocene to early Miocene geochronology and paleoceanography from the subantarctic South Atlantic. Paleoceanography 17, 10.1029/2000PA000568.

Cooper, A.K., Barrett, P.J., Hinz, K., Traube, V., Leitchenkov, G., Stagg, H.M.J., 1991. Cenozoic prograding sequences of the Antarctic continental margin: A record of glacio-eustatic and tectonic events. Marine Geology 102, 175-213.

Coxall, H.K., Wilson, P.A., Palike, H., Lear, C.H., Backman, J., 2005. Rapid stepwise onset of Antarctic glaciation and deeper calcite compensation in the Pacific Ocean. Nature 433, 53-57.

Florindo, F., Wilson, G.S., Roberts, A.P., Sagnotti, L., Verosub, K.L., 2005. Magnetostratigraphic chronology of a late Eocene to early Miocene glacimarine succession from the Victoria basin, Ross Sea, Antarctica. Global and Planetary Change 45, 207-236.

Hambrey, M.J., Ehrmann, W.U., Larsen, B., 1991. Cenozoic glacial record of the Prydz Bay continental shelf, East Antarctica. In Barron, J., et al. (Eds.), Proceedings of the Ocean Drilling Program, Scientific Results 119, Ocean Drilling Program, College Station, Texas, pp. 77-132.

Kominz, M.A., Pekar, S.F., 2001. Oligocene eustasy from two-dimensional sequence stratigraphic backstripping. Geological Society of America Bulletin 113, 291-304.

Lear, C.H., Rosenthal, Y., Coxall, H.K., Wilson, P.A., 2004. Late Eocene to early Miocene ice sheet dynamics and the global carbon cycle. Paleoceanography 19, PA4015, doi:10.1029/2004PA001039.

Miller, K.G., Mountain, G.S., Browning, J.V., Kominz, M., Sugarman, P.J., Christie-Blick, N., Katz, M.E., and Wright, J.D., 1998. Cenozoic global sea level, sequences, and the New Jersey Transect: Results from coastal plains and continental slope drilling. Reviews of Geophysics 36, 569-601.

Miller, K.G., Wright, J.D., Fairbanks, R.G., 1991. Unlocking the ice house: Oligocene-Miocene oxygen isotopes, eustasy, and margin erosion. Journal of Geophysical Research 96, 6,829-6,848.

Miller, K.G., Kominz, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F., 2005. The Phanerozoic record of global sea-level change. Science 312, 1293-1297.

Pagani, M., Zachos, J.C., Freeman, K.H., Tipple, B., Bohaty, S., 2005. Marked decline in atmospheric carbon dioxide concentrations during the Paleogene. Science 309, 600-603.

Pekar, S. F., DeConto, R.M., 2006. High-resolution ice-volume estimates for the early Miocene: Evidence for a dynamic ice sheet in Antarctica. Palaeogeography, Palaeoclimatology, Palaeoecology 231, 101-109.

Pekar, S., Miller, K.G., 1996. New Jersey Oligocene "Icehouse" sequences (ODP Leg 150X) correlated with global $\delta^{18}O$ and Exxon eustatic records. Geology 24, 567-570.

Pekar, S.F., Christie-Blick, N., 2007, Resolving apparent conflicts between oceanographic and Antarctic climate records and evidence for a decrease in pCO_2 during the Oligocene through early Miocene (34-16 Ma): Palaeogeography, Palaeoclimatology, Palaeoecology, in press.

Pekar, S.F., Christie-Blick, N., Kominz, M.A., Miller, K.G., 2002. Calibrating eustasy to oxygen isotopes for the early icehouse world of the Oligocene. Geology 30, 903-906.

Pekar, S.F., Harwood, D.M., DeConto, R.M., 2006. Resolving a late Oligocene conundrum: deep-sea warming versus Antarctic glaciation. Palaeogeography, Palaeoclimatology, Palaeoecology 231, 29-40.

Raine, J.I., Askin, R.A., 2001. Oligocene and Early Miocene terrestrial palynology of the Cape Roberts Drillhole CRP-2/2A, Victoria Land Basin, Antarctica. Terra Antartica 7, 389-400.

Sorlien, C. C., B. P. Luyendyk, D. Wilson, R. Decesari, L. Bartek, J. Diebold, 2007, Oligocene development of the West Antarctic Ice Sheet recorded in eastern Ross Sea strata, Geolog, in press.

Watson, R.T. and the Core Writing Team (Eds.), 2001. Climate Change 2001: Synthesis Report, 2001. Intergovernmental Panel on Climate Change, Geneva, Switzerland. pp 184.

Zachos, J.C., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms and aberrations in global climate 65 Ma to present. Science 292, 686-693.