

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

2002 Bird Strike Committee-USA/Canada, 4th  
Annual Meeting, Sacramento, CA

Bird Strike Committee Proceedings

---

October 2002

# Increasing Air Safety at Eglin Air Force Base through Vulture Roost Dispersal

John S. Humphrey

*USDA, Wildlife Services, National Wildlife Research Center*

Follow this and additional works at: <http://digitalcommons.unl.edu/birdstrike2002>



Part of the [Environmental Health and Protection Commons](#)

---

Humphrey, John S., "Increasing Air Safety at Eglin Air Force Base through Vulture Roost Dispersal" (2002). *2002 Bird Strike Committee-USA/Canada, 4th Annual Meeting, Sacramento, CA*. 17.

<http://digitalcommons.unl.edu/birdstrike2002/17>

This Article is brought to you for free and open access by the Bird Strike Committee Proceedings at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2002 Bird Strike Committee-USA/Canada, 4th Annual Meeting, Sacramento, CA by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

### **Increasing Air Safety at Eglin Air Force Base through Vulture Roost Dispersal**

*John S. Humphrey, USDA, Wildlife Services, National Wildlife Research Center, 2820 East University Avenue, Gainesville, FL 32641 USA*

Forested wetlands provide attractive roost sites for black vultures (*Coragyps atratus*) and turkey vultures (*Cathartes aura*). Vultures entering and departing roosts, however, can create hazardous conditions for pilots arriving and departing nearby airports because vultures often use the same air column as aircraft. This was the case for Eglin Air Force Base and co-located Okaloosa Regional Airport in Niceville, Florida where 260 vultures roosted in nearby Turkey Creek Nature Trail. We evaluated the effectiveness of suspending a taxidermic vulture effigy in the roost, augmented by periodic use of a handheld laser management options to alleviate these aviation safety problems. The laser was used during first light and dusk on the 2<sup>nd</sup> and 3<sup>rd</sup> treatment day. Vulture numbers decreased 41% within 3 days and 100% 4 days after installation of the stimulus. Hanging a vulture effigy from a location within the roost creates an unfavorable roosting environment for vultures and offers a simple, effective means to manage problem roosts. The use of the handheld laser further decreases the desirability of the roost and quite possibly the time it takes to disperse the roost. In this study, dispersal of the vulture roost effectively resolved a potentially dangerous situation. In other cases however, dispersal of roosting birds may not decrease bird strike hazards.