

2-1-2007

HUBBLE SPACE TELESCOPE ULTRAVIOLET SPECTROSCOPY OF 14 LOW-REDSHIFT QUASARS

Rajib Ganguly

University of Wyoming, Laramie, WY, USA

Michael S. Brotherton

University of Wyoming, Laramie, WY, USA

Nahum Arav

University of Colorado, Boulder, arav@vt.edu

Sara R. Heap

NASA Goddard Space Flight Center, Greenbelt, MD, USA

Lutz Wisotzki

Astrophysikalisches Institut Potsdam, Potsdam, Germany

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsgaskell>



Part of the [Physics Commons](#)

Ganguly, Rajib; Brotherton, Michael S.; Arav, Nahum; Heap, Sara R.; Wisotzki, Lutz; Aldcroft, Thomas L.; Alloin, Danielle; Behar, Ehud; Canalizo, Gabriela; Crenshaw, D. Michael; De Kool, Martijn; Chambers, Kenneth; Cecil, Gerald; Chatzichristou, Eleni; Everett, John; Gabel, Jack R.; Gaskell, C. Martin; Galliano, Emmanuel; Green, Richard F.; Hall, Patrick B.; Hines, Dean C.; Junkkarinen, Vesa T.; Kaastra, Jelle S.; Kaiser, Mary Elizabeth; Kazanas, Demosthenes; Konigl, Arieh; Korista, Kirk T.; Kriss, Gerard A.; Laor, Ari; Leighly, Karen M.; Mathur, Smita; Ogle, Patrick; Proga, Daniel; Sabra, Bassem; Sivron, Ran; Snedden, Stephanie A.; Telfer, Randal; and Vestergaard, Marianne, "HUBBLE SPACE TELESCOPE ULTRAVIOLET SPECTROSCOPY OF 14 LOW-REDSHIFT QUASARS" (2007). *Martin Gaskell Publications*. 18.
<http://digitalcommons.unl.edu/physicsgaskell/18>

Authors

Rajib Ganguly, Michael S. Brotherton, Nahum Arav, Sara R. Heap, Lutz Wisotzki, Thomas L. Aldcroft, Danielle Alloin, Ehud Behar, Gabriela Canalizo, D. Michael Crenshaw, Martijn De Kool, Kenneth Chambers, Gerald Cecil, Eleni Chatzichristou, John Everett, Jack R. Gabel, C. Martin Gaskell, Emmanuel Galliano, Richard F. Green, Patrick B. Hall, Dean C. Hines, Vesa T. Junkkarinen, Jelle S. Kaastra, Mary Elizabeth Kaiser, Demosthenes Kazanas, Arie Konigl, Kirk T. Korista, Gerard A. Kriss, Ari Laor, Karen M. Leighly, Smita Mathur, Patrick Ogle, Daniel Proga, Bassem Sabra, Ran Sivron, Stephanie A. Snedden, Randal Telfer, and Marianne Vestergaard

HUBBLE SPACE TELESCOPE ULTRAVIOLET SPECTROSCOPY OF 14 LOW-REDSHIFT QUASARS¹

RAJIB GANGULY,² MICHAEL S. BROTHERTON,² NAHUM ARAV,³ SARA R. HEAP,⁴ LUTZ WISOTZKI,⁵ THOMAS L. ALDCROFT,⁶
DANIELLE ALLOIN,^{7,8} EHUD BEHAR,⁹ GABRIELA CANALIZO,¹⁰ D. MICHAEL CRENSHAW,¹¹ MARTIJN DE KOOL,¹²
KENNETH CHAMBERS,¹³ GERALD CECIL,¹⁴ ELENI CHATZICHRISTOU,¹⁵ JOHN EVERETT,^{16,17} JACK GABEL,³
C. MARTIN GASKELL,¹⁸ EMMANUEL GALLIANO,¹⁹ RICHARD F. GREEN,²⁰ PATRICK B. HALL,²¹
DEAN C. HINES,²² VESA T. JUNKKARINEN,²³ JELLE S. KAASTRA,²⁴ MARY ELIZABETH KAISER,²⁵
DEMOSTHENES KAZANAS,⁴ ARIEH KONIGL,²⁶ KIRK T. KORISTA,²⁷ GERARD A. KRISS,²⁸
ARI LAOR,⁹ KAREN M. LEIGHLY,²⁹ SMITA MATHUR,³⁰ PATRICK OGLE,³¹
DANIEL PROGA,³² BASSEM SABRA,³³ RAN SIVRON,³⁴
STEPHANIE SNEDDEN,³⁵ RANDAL TELFER,³⁶
AND MARIANNE VESTERGAARD³⁷

Received 2006 June 27; accepted 2006 October 4

ABSTRACT

We present low-resolution ultraviolet spectra of 14 low-redshift ($z_{\text{em}} \lesssim 0.8$) quasars observed with the *Hubble Space Telescope* STIS as part of a Snapshot project to understand the relationship between quasar outflows and luminosity. By design, all observations cover the C IV emission line. Ten of the quasars are from the Hamburg-ESO catalog, three are from the Palomar-Green catalog, and one is from the Parkes catalog. The sample contains a few interesting quasars, including two broad absorption line (BAL) quasars (HE 0143–3535 and HE 0436–2614), one quasar with a mini-BAL (HE 1105–0746), and one quasar with associated narrow absorption (HE 0409–5004). These BAL quasars are among the brightest known (although not the most luminous) since they lie at $z_{\text{em}} < 0.8$. We compare the properties of these BAL quasars to the $z_{\text{em}} < 0.5$ Palomar-Green and $z_{\text{em}} > 1.4$ Large Bright Quasar Survey samples. By design, our objects sample luminosities in between these two surveys, and our four absorbed objects are consistent with the $v \sim L^{0.62}$ relation derived by Laor & Brandt (2002). Another quasar, HE 0441–2826, contains extremely weak emission lines, and our spectrum is consistent with a simple power-law continuum. The quasar is radio-loud but has a steep spectral index and a lobe-dominated morphology, which argues against it being a blazar. The unusual spectrum of this quasar resembles the spectra of the quasars PG 1407+265, SDSS J1136+0242, and PKS 1004+13, for which several possible explanations have been entertained.

Key words: quasars: absorption lines — quasars: emission lines — surveys

Online material: color figures

¹ Based on observations made with the NASA/ESA *Hubble Space Telescope*, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

² Department of Physics and Astronomy, University of Wyoming, Laramie, WY, USA; ganguly@uwyo.edu.

³ Center for Astrophysics and Space Astronomy, University of Colorado, Boulder, CO, USA.

⁴ Laboratory of Astronomy and Solar Physics, NASA Goddard Space Flight Center, Greenbelt, MD, USA.

⁵ Astrophysikalisches Institut Potsdam, Potsdam, Germany.

⁶ Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA.

⁷ European Southern Observatory, Santiago, Chile.

⁸ UMR 7158, CEA-CNRS-Université Paris 7, DSM/DAPNIA/Service d’Astrophysique, CEA/Saclay, France.

⁹ Department of Physics, Technion, Haifa, Israel.

¹⁰ Department of Physics, University of California, Riverside, CA, USA.

¹¹ Department of Physics and Astronomy, Georgia State University, Atlanta, GA, USA.

¹² Research School of Astronomy and Astrophysics, Mount Stromlo Observatory, Weston Creek, ACT, Australia.

¹³ Institute for Astronomy, Honolulu, HI, USA.

¹⁴ Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC, USA.

¹⁵ Institute of Astronomy and Astrophysics, National Observatory of Athens, Athens, Greece.

¹⁶ Canadian Institute of Theoretical Astrophysics, University of Toronto, Toronto, ON, Canada.

¹⁷ Department of Astronomy, Department of Physics, and Center for Magnetic Self-Organization, University of Wisconsin, Madison, WI, USA.

¹⁸ Department of Physics and Astronomy, University of Nebraska, Lincoln, NE, USA.

¹⁹ Department of Physics, University of California, Davis, CA, USA.

²⁰ Large Binocular Telescope Observatory, Tucson, AZ, USA.

²¹ Department of Physics and Astronomy, York University, Toronto, ON, Canada.

²² Space Science Institute, Boulder, CO, USA.

²³ Center for Astrophysics and Space Sciences, University of California, San Diego, CA, USA.

²⁴ SRON Netherlands Institute for Space Research, Utrecht, Netherlands.

²⁵ Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD, USA.

²⁶ Department of Astronomy and Astrophysics and Enrico Fermi Institute, University of Chicago, Chicago, IL, USA.

²⁷ Department of Physics, Western Michigan University, Kalamazoo, MI, USA.

²⁸ Space Telescope Science Institute, Baltimore, MD, USA.

²⁹ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman, OK, USA.

³⁰ Department of Astronomy, Ohio State University, Columbus, OH, USA.

³¹ *Spitzer* Science Center, California Institute of Technology, Pasadena, CA, USA.

³² Department of Physics, University of Nevada, Las Vegas, NV, USA.

1. INTRODUCTION

Outflows from active galactic nuclei (AGNs) come in many observational classes. Seyfert galaxies show blueshifted UV and X-ray absorption lines hundreds of km s^{-1} wide (Crenshaw et al. 1999), while the UV troughs of quasar outflows can span tens of thousands of km s^{-1} as manifested in broad absorption line (BAL) quasars (Lynds 1967; Weymann et al. 1985; Turnshek et al. 1988). One of the current driving questions in the AGN field is what is the connection, if any, between the intrinsic luminosity of an AGN and the kinematic properties of the outflow (e.g., terminal velocity and velocity width of the observed trough).

Radiative acceleration, thought to be the principal driver of such outflows, predicts that the terminal velocity should scale as $v \sim L^n$, where $0.25 < n < 0.5$ (Arav et al. 1994). Qualitatively, such a progression is likely to exist given the observed fact that outflow in Seyfert galaxies terminate at $\sim 1000 \text{ km s}^{-1}$, while the BAL outflows extend out to $\sim 30,000 \text{ km s}^{-1}$. However, the quantitative trend is unclear given the lack of objects in between these populations. From an analysis of ~ 56 archived *Hubble Space Telescope* (*HST*) and *International Ultraviolet Explorer* (*IUE*) spectra of $z < 0.5$ quasars from the Palomar-Green survey, Laor & Brandt (2002) showed that such a trend may indeed exist. The soft X-ray-weak (defined as $\alpha_{\text{ox}} < -2.0$) quasars, which exhibit BALs in their UV spectra, show a relation of the form $v \sim L^{0.62 \pm 0.08}$.

This is a higher power-law index than predicted and implies that the radiation-pressure force multiplier has a luminosity dependence (Laor & Brandt 2002). The force multiplier (Castor et al. 1975) is a factor that expresses the sum total effect of all lines and edges in transferring momentum from the incident spectrum of photons to the gas. The luminosity dependence of the force multiplier arises from a variety of sources like the dust content and ionization state of the gas (e.g., Arav et al. 1994; Murray et al. 1995; Sulentic et al. 2006). Indeed, considerations of accretion-disk winds by Murray et al. (1995) in both the context of the broad emission lines and broad absorption lines require the presence of shielding gas that prevents X-rays from overionizing the outflowing gas. More recent considerations of the force multiplier show that it is also sensitive to black hole mass (Proga & Kallman 2004), which then favors a steeper index than predicted by Arav et al. (1994). The Laor & Brandt (2002) index is also consistent with the prediction from Proga et al. (1998) for AGNs with $L/L_{\text{Edd}} \geq M_{\text{max}}^{-1}$, where M_{max} is the maximum value of the force multiplier.

Another crucial issue is whether the fraction of objects that shows intrinsic absorption is luminosity dependent. Here we are facing a large gap between the established statistics for BAL quasars (see below) and the results from a biased sample of Seyfert galaxies that showed that 10 out of the available 17 *HST* UV spectra of Seyfert galaxies show intrinsic absorption (Crenshaw et al. 1999). Kriss (2002) found that a similar fraction of Seyfert galaxies (16/34) exhibit absorption in the $\text{O VI } \lambda\lambda 1031.926, 1037.617$ doublet. At low redshift, Ganguly et al. (2001) found $\sim 25\%$ of quasars from the *HST* Quasar Absorption Line Key Project appear to show absorption at $z_{\text{abs}} \sim z_{\text{em}}$ (i.e., ‘‘associated’’ absorption), comparable to the study of Laor & Brandt (2002) for the $z < 0.5$ Palomar-Green quasars. However, it must be noted that not all ab-

sorption found near quasars is necessarily intrinsic and, moreover, not necessarily part of an outflow. A follow-up study of low-redshift quasars with associated absorption showed that only $\sim 30\%$ were time variable (Wise et al. 2004). Furthermore, large statistical studies of absorption in quasars, like the *HST* Quasar Absorption Line Key Project, tend to be biased toward UV-bright targets in order to obtain adequate signal-to-noise ratio (S/N) spectra.

For high-redshift quasars ($z > 1.4$) intrinsic absorption (usually BALs) has been studied in large samples using ground-based telescopes. The Large Bright Quasar Survey (LBQS; Hewett et al. 1995) members at these redshifts have $M_V = -26$ to -28 and an intrinsic fraction of BAL quasars, in the redshift range $1.5 < z < 3.0$, of $22\% \pm 4\%$ (Hewett & Foltz 2003). The FIRST Bright Quasar Survey (White et al. 2000; Becker et al. 2001) members have $M_V = -26$ to -29 and a BAL incidence of 18% (Becker et al. 2000). Vestergaard (2003) found that about 25% of quasars show evidence for intrinsic, low-velocity C IV absorption from a careful consideration of possible contamination from intervening systems in a heterogeneous yet large sample of moderate redshift ($1.5 < z < 3.0$) quasars. In addition, Richards (2001) found that, in C IV absorption systems observed toward $z \sim 2.5$ FIRST quasars at large velocity separations, as many as 30% may be intrinsic to the background quasars. Such systems must clearly arise in a high-velocity outflow.

The relationship between the velocity of AGN outflows and their intrinsic luminosity, as well as the fraction of objects that show outflows, is poorly known due to two simple selection effects. The vast majority of AGN outflows are identified by detecting absorption features associated with the C IV $\lambda\lambda 1548.20, 1550.77$ doublet. From ground-based observations this line is only observed in objects with redshift $z > 1.3$ and, therefore, the population observed tends to be the higher luminosity quasars, of which $\sim 10\%$ are BAL quasars. In order to observe objects with $z < 1.3$ we must use space-based UV observatories. Only ~ 20 of the UV brightest AGNs that were observed with the *HST* (out of ~ 70) show evidence for outflows, compared with upward of 500 BAL quasars identified in ground-based observations. The statistics are especially poor for the luminosity range 10^{44} to $10^{45.5} \text{ ergs s}^{-1}$ (corresponding roughly to the absolute magnitude range $-21.9 \lesssim M_V \lesssim -25.6$).

To fill in this dearth of data, we proposed an *HST* Snapshot project to obtain low-dispersion STIS UV spectra of intermediate-luminosity AGNs. Unfortunately, the low efficiency of this program only yielded observations of 14 objects out of 200 approved targets, and we present the data here. Several of the objects are of individual interest based on their absorption properties. In the next section (§ 2) we present the spectra obtained and discuss our data reduction. In § 3 we present our results and compare them with those from other quasar samples. We briefly summarize our findings in § 4.

2. DATA

Our *HST* STIS (Woodgate et al. 1998; Kimble et al. 1998) observations were carried out using the G230L grating and the $52'' \times 0.5''$ slit, which provides spectra over the wavelength range $1570\text{--}3180 \text{ \AA}$ at a dispersion of $1.58 \text{ \AA pixel}^{-1}$ (and a 2 pixel per resolution element sampling rate). For these Snapshot observations, we used exposure times of 900 s with three exceptions. We used an exposure time of 720 s for the quasars HE 0354–5500, HE 0436–2614, and PG 1435–067. (These three quasars were part of the bright end of our initial list of 200 targets, defined as objects with $B < 16$, and therefore did not require a full 900 s exposure.)

³³ Faculty of Natural and Applied Sciences, Notre Dame University, Zouk Mosbeh, Lebanon.

³⁴ Department of Physics, Baker University, Baldwin City, KS, USA.

³⁵ Apache Point Observatory, Sunspot, NM, USA.

³⁶ Orbital Sciences Corporation, Greenbelt, MD, USA.

³⁷ Steward Observatory, The University of Arizona, Tucson, AZ, USA.




FIG. 1.—Spectra (flux vs. observed wavelength; *histogram*) for the subsample of quasars observed for our Snapshot program. The spectra are ordered by right ascension. In addition, we overlay the power-law fit (*smooth curve*) described in § 3.

We used the standard pipeline that provides fully reduced and calibrated spectra. According to the STIS Instrument Handbook (Kim-Quijano et al. 2003), data reduced using the pipeline have the following calibration uncertainties: 0.5–1.0 pixels (0.79–1.58 Å) in absolute wavelength calibration, and 4% (0.02 dex) in

absolute spectrophotometry. The S/N for our spectra were typically ~ 14 per pixel at 2000 and 3000 Å.

Figure 1 shows the spectra obtained for our program. The data are publicly available both at the Multimission Archive at Space Telescope (MAST)³⁸ and at the University of Wyoming AGN Research Group Web site.³⁹

3. RESULTS

3.1. Sample Characteristics

To characterize this sample and place it in context with larger samples of low-redshift quasars, we carried out power-law fits to all spectra with the goal of computing UV spectral slopes and luminosities. The fits were carried out using an arbitrary number of superposed Gaussian curves to mimic the contribution from emission lines. That is, we fit the following functional form to our spectra:

$$F_{\lambda} = F_{\lambda_o} \left[\left(\frac{\lambda}{\lambda_o} \right)^{\alpha} + \sum_{i=1}^m w_i \exp \left(-\frac{\lambda - \lambda_i}{\sigma_i} \right)^2 \right], \quad (1)$$

where F_{λ_o} is the normalizing flux at reference wavelength λ_o , α is the spectral index, m is the number of emission-line components each with a relative strength w_i , σ_i is the width, and the central wavelength is λ_i . The best fit was determined using the Numerical Recipes Marquardt-Levenburg χ^2 minimization routines (Press et al. 1992), and the optimal number of Gaussian components was determined using an F -test. In carrying out the fits, we omitted regions that were clearly, or potentially, affected by absorption (as subjectively determined by R. G.). The results of the power-law fits are listed in Table 1 and overlaid on the observed spectra in Figure 1. For uniformity, we used a common (observer's frame) reference wavelength, $\lambda_o = 1800$ Å for all fits. Table 1 lists the quasar name (col. [1]), quasar redshift (z , col. [2]), continuum flux density at 1800 Å (F_{λ_o} , col. [3]), continuum power-law index (α , col. [4], with the sign convention $F_{\lambda} \sim \lambda^{\alpha}$), the luminosity at rest-frame wavelength $\lambda_r = 3000$ Å (L_{λ_r} , col. [5]), and the C iv emission-line equivalent width for unabsorbed quasars (col. [6]). We note that in a few cases (e.g., PG 2233+134) for which the spectra cover wavelengths redward of the Al III + C III emission line, the continuum fit can be artificially elevated due to the presence of Fe II-III emission (Vestergaard & Wilkes 2001). The observer-frame flux density at observed wavelength λ_o was converted to rest-frame luminosity at rest wavelength λ_r via

$$\begin{aligned} \lambda_r L_{\lambda_r} &= 4\pi D_L(z)^2 (1+z) \lambda_r F_{\lambda_r} \\ &= 4\pi D_L(z)^2 (1+z) \lambda_r F'_{\lambda_r(1+z)} \\ &= 4\pi D_L(z)^2 (1+z) \lambda_r F'_{\lambda_o} \left[\lambda_r (1+z) / \lambda_o \right]^{\alpha}, \end{aligned} \quad (2)$$

where $D_L(z)$ is the luminosity distance to the quasar, and we have used the continuum portion of equation (1) in the second substitution. For clarity, we have used F' to indicate the flux in the observer frame (which is the flux derived using eq. [1]), and F to indicate the flux in the rest frame. The luminosity distance was computed using the listed quasar redshift and a $\Omega_M = 0.27$, $\Omega_{\Lambda} = 0.73$, $H_0 = 71$ km s⁻¹ Mpc⁻¹ cosmology. The mean luminosity of the sample is $\langle \lambda_r L_{\lambda_r}(3000 \text{ Å}) \rangle = (7.76 \pm 0.02) \times 10^{44}$ ergs s⁻¹, and has a range spanning $(0.3\text{--}31.4) \times 10^{45}$ ergs s⁻¹. Our fits imply a mean spectral index of $\langle \alpha_{UV} \rangle = -1.13 \pm 0.01$, with a standard deviation in the distribution of $\sigma_{\alpha} = 0.53$.

³⁸ Available at <http://archive.stsci.edu/index.html>; fits files only.

³⁹ Available at <http://physics.uwyo.edu/agn>; both fits and ASCII files.

TABLE 1
QUASAR PROPERTIES

QUASAR (1)	REDSHIFT (2)	POWER-LAW FIT			
		$F_{\lambda}(1800 \text{ \AA})$ ($10^{-15} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$) (3)	α_{UV} (4)	$\log \lambda L_{\lambda}$ (3000 \AA) (5)	$W_e(\text{C IV})$ (\AA) (6)
HE 0143–3535.....	0.446	1.51 ± 0.02	-0.51 ± 0.04	45.477 ± 0.006	...
PKS 0252–549.....	0.539	1.36 ± 0.02	-1.45 ± 0.04	45.260 ± 0.006	107 ± 5
HE 0354–5500.....	0.267	7.49 ± 0.05	-1.40 ± 0.02	45.103 ± 0.003	33 ± 2
HE 0409–5004.....	0.817	3.53 ± 0.03	-0.60 ± 0.02	46.493 ± 0.003	...
HE 0436–2614.....	0.690	3.02 ± 0.06	-0.65 ± 0.05	46.209 ± 0.009	...
HE 0441–2826.....	0.155	15.36 ± 0.04	-1.13 ± 0.01	45.207 ± 0.001	3.5 ± 0.9
HE 0502–2948.....	0.552	9.86 ± 0.05	-1.53 ± 0.01	46.107 ± 0.002	35 ± 1
HE 1006–1211.....	0.693	4.11 ± 0.03	-1.04 ± 0.02	46.172 ± 0.003	36 ± 2
HE 1101–0959.....	0.186	1.02 ± 0.02	-0.33 ± 0.07	44.439 ± 0.009	140 ± 7
HE 1105–0746.....	0.340	3.38 ± 0.03	-1.35 ± 0.03	45.236 ± 0.004	...
PG 1435–067.....	0.126	6.24 ± 0.15	-0.49 ± 0.06	44.794 ± 0.010	168 ± 8
PG 1552+085.....	0.119	2.92 ± 0.02	0.11 ± 0.02	44.573 ± 0.003	51 ± 4
PG 2233+134.....	0.325	6.14 ± 0.04	-1.33 ± 0.02	45.458 ± 0.003	7 ± 1
HE 2327–5522.....	0.494	2.72 ± 0.04	-1.14 ± 0.03	45.595 ± 0.006	119 ± 5

NOTES.—For the power-law fit, we adopt the convention $F_{\lambda} \sim \lambda^{\alpha}$. The rest-frame luminosity at 3000 \AA was computed assuming an $\Omega_M = 0.27$, $\Omega_{\Lambda} = 0.73$, $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ cosmology. The quoted 1σ error in both flux and luminosity reflects statistical fitting uncertainties only. The error associated with the uncertainty in absolute flux calibration, which dominates the error in both flux and luminosity, is 4%, or 0.02 dex. For quasars that do not show intrinsic/associated absorption, we list the C IV emission-line equivalent width [$W_e(\text{C IV})$].

In Figure 2, we show the region around the C IV emission line for all 14 quasars. Several quasars show absorption on top of the emission line, but most of these are identified with Galactic lines from Fe II and Mg II as indicated in the figure. Four quasars do appear to show intrinsic or associated C IV absorption, and we discuss these in the following sections.

3.2. Absorbed Quasars

Four of the 14 quasar spectra appear to show intrinsic/associated C IV absorption: HE 0143–3535, HE 0409–5004, HE 0436–2614, and HE 1105–0746. Figure 3 shows the spectra as luminosity versus rest wavelength, and ordered (*top to bottom*) by decreasing maximum ejection velocity of absorption. To characterize the absorption properties of these quasars, we carried out fits to the C IV emission line. We took the same approach as in § 3.1, and the fits are also shown (with the 1σ statistical uncertainty) in the figure. These fits allowed us to compute the following properties, which are listed in Table 2: “balmicity” index (col. [2]; Weymann et al. 1991), intrinsic absorption index (col. [3]; Hall et al. 2002), C IV absorption equivalent width (col. [4]), and maximum velocity of absorption (col. [5]).

We note that the fits use the minimum number of Gaussian components, and as a result there may be some systematic uncertainties. In particular, for the two BAL quasars, HE 0143–3535 and HE 0436–2614, if there is significant asymmetry in the intrinsic emission-line profiles (e.g., Wills et al. 1993; Brotherton et al. 1994; Richards et al. 2002 and references therein), it is possible that our fits do not fully recover the correct shape. Parameters such as the balmicity index that depend on the power-law fit only should not be affected by the systematics of trying to reproduce the emission-line profile.

3.3. Comparison to Other Samples

Figure 4 shows a comparison of the quasar luminosities from this sample to $z_{\text{em}} < 0.5$ quasars in the Bright Quasar Survey (BQS; Brandt et al. 2000; Boroson & Green 1992; Schmidt & Green 1983) and $z_{\text{em}} > 1.4$ broad absorption line quasars from the

LBQS (Gallagher et al. 2006; Hewett & Foltz 2003; Hewett et al. 1995). For the LBQS, we restrict the comparison of the luminosity distribution to the subsample of absorbed quasars from Gallagher et al. (2006) since this is the most interesting aspect of our initial survey. While the entire LBQS sample does extend down to $z_{\text{em}} = 0.2$, good ground-based spectroscopic observations of the C IV spectral region are available only for $z_{\text{em}} > 1.4$, where the UV doublet shifts into the optical band. The luminosities of our quasar sample lie in between these two surveys, although with some overlap with the low-redshift BQS.

Figure 5 shows a comparison of our quasar spectral indices to several other samples: Palomar-Green quasars from Neugebauer et al. (1987), LBQS BAL quasars from Gallagher et al. (2006), quasars used in the *FUSE* composite by Scott et al. (2004), and quasars from Shang et al. (2005). In general, there is agreement between the comparison samples (and in particular between the Palomar-Green, absorbed LBQS, and Shang et al. [2005] samples), with a peak near $\alpha \sim -1.8$. Scott et al. (2004) showed a correlation between the spectral index and luminosity, with more luminous objects having bluer spectra. This correlation nicely explains the difference in shape between the *FUSE* composite and the EUV portion of the *HST* composite spectrum from Telfer et al. (2002). The spectral indices derived for the quasars in our samples tend to be redder than those samples, with a peak near $\alpha \sim -1.4$ and an aforementioned mean of -1.14 .

One potential explanation for this is the effect of reddening. Galactic reddening is an unlikely explanation, as the largest color excess in our sample is $E(B - V) = 0.063$ toward PG 1435–067 and HE 1006–1211. Baskin & Laor (2005) noted that quasars (particularly from the BQS) appear redder (as measured by a two-point spectral index between 1549 and 4861 \AA) as more intrinsic C IV absorption is present, implying that intrinsic dust is present along sight lines that also produce intrinsic C IV absorption. While intrinsic reddening may be important, it is unlikely to be the source of reddening in our quasars, as our three reddest quasars (PG 1552+085, HE 1101–0959, and PG 1435–067) do not show intrinsic C IV absorption. In addition, our quasars are generally




FIG. 2.—Region around the C IV emission line for the 14 quasars in this sample. From top to bottom, the quasars are ordered in increasing redshift. Absorption features that appear in this wavelength range are marked, and the best-fit power law is also shown as a smooth curve.

redder than the LBQS BAL quasars from Gallagher et al. (2006), although this may arise from the luminosity effect described by Scott et al. (2004).

Since one of our initial goals was to test the radiative-driving hypothesis for quasar winds, we show in Figure 6 a plot of the maximum velocity of absorption against luminosity for our absorbed quasars, the BQS quasars from Laor & Brandt (2002), and the LBQS BAL quasars from Gallagher et al. (2006). From a consideration of soft X-ray weak quasars in the BQS sample, Laor & Brandt (2002) reported an apparent envelope to the maximum velocity of absorption as a function of luminosity (specifically M_V , but see also Gallagher et al. [2006] for a plot vs. 2500 Å luminosity). We reproduce the Laor & Brandt (2002) plot, revised for a concordance cosmology, with the four quasars from this work, also shown. The two BAL quasars from this sample, HE 0143–3535 and HE 0436–2614, both seem to lie close to the (cosmology-corrected) best-fit curve derived by Laor & Brandt (2002):

$$v_{\max} = 4100 \text{ km s}^{-1} \left[\frac{\nu L_{\nu}(2500 \text{ \AA})}{10^{44.67} \text{ ergs s}^{-1}} \right]^{0.62} \quad (3)$$

The other two absorbed quasars, like most of the absorbed BQS quasars, lie below this curve. There are several potential explanations for this. Since the measured velocities only measure the line-of-sight component and not the transverse (i.e., in the plane of the sky) component, the scatter below this curve is possibly explained by orientation effects. In addition, other driving mechanisms, such as magnetocentrifugal accretion disk winds (Everett 2005) or thermally driven winds (Krolik & Kriss 2001), may be important for absorbers at lower velocities. At very low velocities the absorbers may not even arise from an outflowing wind. They may arise from satellite galaxies around the quasar host galaxies, or even from the interstellar medium of the host galaxy (e.g., Ganguly et al. 2006; Hamann et al. 2001).

3.4. Notes on Individual Quasars

HE 0143–3535.—This is one of two BAL quasars in the sample. The BAL is detected clearly in both C IV and N V. It is marginally detected in H I Ly α in a relatively low S/N (≈ 6) region of the spectrum. Reproduction of the C IV emission line for this quasar is problematic (Fig. 3). Our best fit requires two components, a strong narrow component and a broad weaker component, as motivated by an inflection on the red wing that cannot be produced by a single Gaussian component. The components are offset by 1441 km s $^{-1}$, implying significant asymmetry. The uncertainty in the shape of the emission line, and consequently the uncertainty in the C IV absorption equivalent width, is significant. Regardless, if our assessment of the C IV absorption equivalent width is accurate, then this quasar may be significantly absorbed in the soft X-ray band. Furthermore, if the shape of the C IV emission line is accurate, then the absorption profile is consistent (within errors) of occulting the UV continuum only.

HE 0409–5004.—Three absorption doublets appear on the weak C IV emission line of this quasar, which is well fit by a single Gaussian. Two of those are the Mg II $\lambda\lambda 2796, 2803$ doublet at $z_{\text{abs}} = -0.00006$ (i.e., Galactic absorption) and $z_{\text{abs}} = 0.0143$. The third doublet, which appears blueward of the other two, is an associated narrow C IV absorber at $z_{\text{abs}} = 0.7865$ with a velocity offset of 4579 ± 165 km s $^{-1}$ from the emission redshift of the quasar. The absorption-line system is also detected in H I Ly α . The N V $\lambda\lambda 1238, 1242$ and O VI $\lambda\lambda 1031, 1038$ doublets are in our wavelength range, but neither is detected. We do detect absorption from low-ionization species, C II, S II–III, the Ly β line, and




FIG. 3.—Spectra [luminosity vs. relative velocity, with $\lambda = 1548.204 \text{ \AA} (1 + z_{\text{em}})$ defining the velocity zero point] of the C IV emission and absorption lines for the four AGNs that show intrinsic/associated absorption. From top to bottom, the AGN spectra are shown in order of decreasing maximum ejection velocity of absorption. The luminosity profiles are shown as a black histogram. The smooth curve around the C IV emission line is our assessment of the effective continuum level (i.e., power-law continuum plus emission lines), with the shaded region indicating the 1σ uncertainty. The horizontal bars indicate wavelength regions that were omitted from the fit due to the presence of absorption. [See the electronic edition of the Journal for a color version of this figure.]

possibly the Lyman limit. This implies that the gas is in a relatively low ionization state (compared to other associated narrow absorption line systems). Thus, it is possible that this system, while close in velocity, is not physically linked with the quasar central engine.

HE 0436–2614.—This quasar is the second in the sample that hosts a BAL. The BAL absorption in this quasar is very strong. The wavelength range of our spectrum gives us coverage down to the O VI $\lambda\lambda 1031, 1038$ doublet. We detect broad absorption from C IV, Si IV, N V, H I Ly α , P V, and O VI. The detection of P V, which has a low relative abundance, implies a combination of high column density in the flow and, perhaps, high metallicity (Hamann 1998). In the C IV BAL profile, there appears to be a curious transition in the apparent strength of the absorption (whether from coverage fraction or column density effects) around $v = -1.2 \times 10^4 \text{ km s}^{-1}$, where the emission line appears to terminate (Fig. 3). Redward of this velocity (i.e., closer to the emission line), the absorption appears to be completely saturated with full coverage of both the continuum and broad emission line regions. In the velocity range $-1.5 \times 10^4 \text{ km s}^{-1} \lesssim v \lesssim -1.2 \times 10^4 \text{ km s}^{-1}$, the absorption appears to have a flat bottom with a normalized

flux of ~ 0.31 (in units of the continuum plus emission-line flux). In the velocity range $-1.8 \times 10^4 \text{ km s}^{-1} \lesssim v \lesssim -1.5 \times 10^4 \text{ km s}^{-1}$, the profile is also flat-bottomed, but with a normalized flux of ~ 0.38 . Blueward of $v \sim -1.8 \times 10^4 \text{ km s}^{-1}$, the absorption tapers off (either due to coverage fraction change or column density change) to no apparent absorption at $\sim -2.4 \times 10^4 \text{ km s}^{-1}$.

HE 0441–2826.—The spectrum of this quasar is peculiar. The C IV emission line is very weak ($\sim 10\%$ of the continuum strength,

Quasar (1)	BI (km s^{-1}) (2)	AI (km s^{-1}) (3)	$W_a(\text{C IV})$ (\AA) (4)	v_{max} (km s^{-1}) (5)
HE 0143–3535.....	2829	3701	36 ± 4	13519 ± 207
HE 0409–5004.....	1.4 ± 0.3	5417 ± 165
HE 0436–2614.....	11592	12740	118 ± 3	24595 ± 177
HE 1105–0746.....	8 ± 1	5151 ± 224

NOTES.—For the BAL QSOs, we report balnicity (BI) and intrinsic absorption indices (AI) as defined by Weymann et al. (1991) and Hall et al. (2002). We also provide an estimate of the observed C IV absorption equivalent width (W_a). The quoted 1σ confidence error includes both statistical and effective-continuum fitting uncertainties.




FIG. 4.—Comparison of the luminosity distribution of quasars in our sample to that of the BQS and LBQS samples. Data for the BQS were taken from Laor & Brandt (2002). Data from the LBQS were taken from Gallagher et al. (2006). [See the electronic edition of the Journal for a color version of this figure.]




FIG. 5.—Distribution of UV spectral indices ($F_\lambda \sim \lambda^\alpha$) for our sample. The histogram is normalized to unit area. We compare this distribution to other samples, including Palomar-Green quasars from Neugebauer et al. (1987), LBQS BAL quasars from Gallagher et al. (2006), quasars used in the *FUSE* composite by Scott et al. (2004), and quasars from Shang et al. (2005). [See the electronic edition of the *Journal* for a color version of this figure.]

the lowest of the sample), and the spectrum is consistent with a simple power law with Galactic Fe II and Mg II absorption. (The redshift of the quasar is known from optical emission lines, so we are secure in the identification of this UV line as C IV.) In the rest frame of the quasar, the spectrum covers the wavelength range 1376–2726 Å. This range also covers the Si IV $\lambda\lambda$ 1393, 1402 doublet, the Al III $\lambda\lambda$ 1855, 1863 + C III] λ 1909 blend, and some of the Fe II UV multiplet, but these are not significantly detected. The quasar is detected in the NRAO-VLA Sky Survey (Condon et al. 1998) with a 1.4 GHz flux of 147.8 ± 4.5 mJy, and a radio core fraction of $\sim 0.24 \pm 0.01$. The quasar is also listed in the Parkes-MIT-NRAO survey catalog (PMN J0443–2820) with a 4.85 GHz flux of 56 ± 11 mJy. We conclude that the quasar has a radio loudness (Kellermann et al. 1994) of $\log R^* = 1.6 \pm 0.2$ and a radio spectral index of $\beta = -0.78 \pm 0.16$ ($F_\nu \sim \nu^\beta$). While weak/absent emission lines may be indicative of a blazar, the marginal radio-loudness, steep radio spectral index, and lobe-dominated morphology seem to rule this out. Alternatively, this quasar could be in the class of unusual emission-line objects that includes the radio-quiet PG 1407+265 ($z_{\text{em}} = 0.94$; McDowell et al. 1995), the radio-loud SDSS J1136+0242 ($z_{\text{em}} = 2.4917$; Hall et al. 2004), and possibly the intrinsic spectrum of the radio-loud BAL PKS 1004+13 ($z_{\text{em}} = 0.24$; Wills et al. 1999). These objects have very weak/absent and highly blueshifted high-ionization emission lines.

HE 1105–0746.—Our spectrum of this quasar covers the rest-frame wavelength range 1187–2350 Å, and the H I Ly α + N V, Si IV, C IV, He II, and Al III + C III] emission lines are clearly visible. The C IV emission line is well described by a single Gaussian curve, and part of the blue wing of the C IV emission line of this quasar appears to be absorbed. This absorption lies above the continuum and therefore the quasar has a zero “balnicity.” However, the width of the absorption is broader than the C IV doublet separation, so this is an example of a so-called mini-BAL. While the absorption does not dip below the continuum, the maximum depth of the absorption is equal to the strength of the continuum. That is, this mini-BAL is consistent with saturated absorption of the con-




FIG. 6.—Maximum velocity of absorption vs. 3000 Å luminosity plot from Laor & Brandt (2002, revised for a concordance cosmology). The solid curve represents their least-squares fit for the soft X-ray-weak quasars in the $z < 0.5$ BQS (see eq. [3], which has also been corrected for cosmology). The shaded region about the curve is the 1σ confidence uncertainty. We overplot the four absorbed quasars from this work. [See the electronic edition of the *Journal* for a color version of this figure.]

tinuum and no absorption of the C IV broad emission line. This has been observed before in BALs (e.g., Arav et al. 1999). Curiously, we do not detect mini-BAL absorption in N V or Si IV. The lack of Si IV absorption may be an ionization effect. The lack of N V absorption may be due to a combination of the noisiness of the spectrum in the wavelength region and dilution by the strong H I Ly α line if indeed the absorber does not occult the broad emission line region.

PG 1552+085.—Based on an *IUE* spectrum, Turnshek et al. (1997) claimed that this quasar was host to a BAL. However, a more recent analysis by Sulentic et al. (2006) of the *HST* STIS G230L spectrum presented here yield a balnicity index of ≈ 0 . The putative location of the C IV BAL lies in a low S/N region of the spectrum, and it is difficult to ascertain if the continuum blueward of the C IV broad emission line is absorbed.

4. SUMMARY

While our initial survey was designed to fill in the statistical gap between luminous quasars and Seyfert galaxies to further understand the relationship between outflows and luminosity, the low efficiency of the program did not make this feasible. Out of the approved list of 200 targets, only 14 objects were observed. The observations of 14 nearby bright AGNs resulted in the discovery of a few interesting objects that merit further study: two new, bright BAL quasars, one new mini-BAL quasar, and one quasar with unusually weak emission lines. Our observations are in agreement with the v_{max} -luminosity relation reported by Laor & Brandt (2002) and lie in between the BQS and LBQS samples.

Support for this work was provided by NASA through grant number HST-GO-09507 from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555.

REFERENCES

- Arav, N., Becker, R. H., Laurent-Muehleisen, S. A., Gregg, M. D., White, R. L., Brotherton, M. S., & de Kool, M. 1999, *ApJ*, 524, 566
- Arav, N., Li, Z.-Y., & Begelman, M. C. 1994, *ApJ*, 432, 62
- Baskin, A., & Laor, A. 2005, *MNRAS*, 356, 1029
- Becker, R. H., White, R. L., Gregg, M. D., Brotherton, M. S., Laurent-Muehleisen, S. A., & Arav, N. 2000, *ApJ*, 538, 72
- Becker, R. H., et al. 2001, *ApJS*, 135, 227
- Boroson, T. A., & Green, R. F. 1992, *ApJS*, 80, 109
- Brandt, W. N., Laor, A., & Wills, B. J. 2000, *ApJ*, 528, 637
- Brotherton, M. S., Wills, B. J., Steidel, C. C., & Sargent, W. L. W. 1994, *ApJ*, 423, 131
- Castor, J. I., Abbott, D. C., & Klein, R. I. 1975, *ApJ*, 195, 157
- Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B., & Broderick, J. J. 1998, *AJ*, 115, 1693
- Crenshaw, D. M., Kraemer, S. B., Boggess, A., Maran, S. P., Mushotzky, R. F., & Wu, C. 1999, *ApJ*, 516, 750
- Everett, J. E. 2005, *ApJ*, 631, 689
- Gallagher, S. C., Brandt, W. N., Chartas, G., Priddey, R., Garmire, G. P., & Sambruna, R. M. 2006, *ApJ*, 644, 709
- Ganguly, R., Bond, N. A., Charlton, J. C., Eracleous, M., Brandt, W. N., & Churchill, C. W. 2001, *ApJ*, 549, 133
- Ganguly, R., Sembach, K. R., Tripp, T. M., Savage, B. D., & Wakker, B. P. 2006, *ApJ*, 645, 868
- Hall, P. B., et al. 2002, *ApJS*, 141, 267
- . 2004, *AJ*, 127, 3146
- Hamann, F. 1998, *ApJ*, 500, 798
- Hamann, F. W., Barlow, T. A., Chaffee, F. C., Foltz, C. B., & Weymann, R. J. 2001, *ApJ*, 550, 142
- Hewett, P. C., & Foltz, C. B. 2003, *AJ*, 125, 1784
- Hewett, P. C., Foltz, C. B., & Chaffee, F. H. 1995, *AJ*, 109, 1498
- Kellermann, K. I., Sramek, R. A., Schmidt, M., Green, R. F., & Shaffer, D. B. 1994, *AJ*, 108, 1163
- Kim-Quijano, J., et al. 2003, *STIS Instrument Handbook*, Ver. 7.0 (Baltimore: STScI)
- Kimble, R. A., et al. 1998, *Proc. SPIE*, 3356, 188
- Kriss, G. A. 2002, in *ASP Conf. Ser. 255, Mass Outflow in Active Galactic Nuclei: New Perspectives*, ed. D. M. Crenshaw, S. B. Kraemer, & I. M. George (San Francisco: ASP), 69
- Krolik, J. H., & Kriss, G. A. 2001, *ApJ*, 561, 684
- Laor, A., & Brandt, W. N. 2002, *ApJ*, 569, 641
- Lynds, C. R. 1967, *ApJ*, 147, 396
- McDowell, J. C., Canizares, C., Elvis, M., Lawrence, A., Markoff, S., Mathur, S., & Wilkes, B. J. 1995, *ApJ*, 450, 585
- Murray, N., Chiang, J., Grossman, S. A., & Voit, G. M. 1995, *ApJ*, 451, 498
- Neugebauer, G., Green, R. F., Matthews, K., Schmidt, M., Soifer, B. T., & Bennett, J. 1987, *ApJS*, 63, 615
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical Recipes in C: The Art of Scientific Computing* (2nd ed.; Cambridge: Cambridge Univ. Press)
- Proga, D., & Kallman, T. R. 2004, *ApJ*, 616, 688
- Proga, D., Stone, J. M., & Drew, J. E. 1998, *MNRAS*, 295, 595
- Richards, G. T. 2001, *ApJS*, 133, 53
- Richards, G. T., Vanden Berk, D. E., Reichard, T. A., Hall, P. B., Schneider, D. P., SubbaRao, M., Thakar, A. R., & York, D. G. 2002, *AJ*, 124, 1
- Schmidt, M., & Green, R. F. 1983, *ApJ*, 269, 352
- Scott, J. E., Kriss, G. A., Brotherton, M., Green, R. F., Hutchings, J., Shull, J. M., & Zheng, W. 2004, *ApJ*, 615, 135
- Shang, Z., et al. 2005, *ApJ*, 619, 41
- Sulentic, J. W., Dultzin-Hacyan, D., Marziani, P., Bongardo, C., Braitto, V., Calvani, M., & Zamanov, R. 2006, *Rev. Mex. AA*, 42, 23
- Telfer, R. C., Zheng, W., Kriss, G. A., & Davidsen, A. F. 2002, *ApJ*, 565, 773
- Turnshek, D. A., Grillmair, C. J., Foltz, C. B., & Weymann, R. J. 1988, *ApJ*, 325, 651
- Turnshek, D. A., Monier, E. M., Sirola, C. J., & Espey, B. R. 1997, *ApJ*, 476, 40
- Vestergaard, M. 2003, *ApJ*, 599, 116
- Vestergaard, M., & Wilkes, B. J. 2001, *ApJS*, 134, 1
- Weymann, R. J., Morris, S. L., Foltz, C. B., & Hewett, P. C. 1991, *ApJ*, 373, 23
- Weymann, R. J., Turnshek, D. A., & Christiansen, W. A. 1985, in *Astrophysics of Active Galaxies and Quasi-Stellar Objects*, ed. J. S. Miller (Mill Valley: University Science Books), 333
- White, R. L., et al. 2000, *ApJS*, 126, 133
- Wills, B. J., Brandt, W. N., & Laor, A. 1999, *ApJ*, 520, L91
- Wills, B. J., Brotherton, M. S., Fang, D., Steidel, C. C., & Sargent, W. L. W. 1993, *ApJ*, 415, 563
- Wise, J. H., Eracleous, M., Charlton, J. C., & Ganguly, R. 2004, *ApJ*, 613, 129
- Woodgate, B. E., et al. 1998, *PASP*, 110, 1183