2011

3-Alkoxy-1,2-Dioxolanes: Synthesis and Evaluation as Potential Antimalarial Agents

Charles E. Schiaffo

University of Nebraska - Lincoln

Matthias Rottman

Swiss Tropical and Public Health Institute

Sergio Wittlin

Swiss Tropical and Public Health Institute

Patrick Dussault

University of Nebraska-Lincoln, pdussault1@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/chemistrydussault>

Schiaffo, Charles E.; Rottman, Matthias; Wittlin, Sergio; and Dussault, Patrick, "3-Alkoxy-1,2-Dioxolanes: Synthesis and Evaluation as Potential Antimalarial Agents" (2011). *Patrick Dussault Publications*. 20.

<http://digitalcommons.unl.edu/chemistrydussault/20>

This Article is brought to you for free and open access by the Published Research - Department of Chemistry at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Patrick Dussault Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

ACS Med Chem Lett. 2011 Apr 14; 2(4): 316–319.

PMCID: PMC3109996

Published online 2011 Jan 26. doi: [10.1021/ml100308d](https://doi.org/10.1021/ml100308d)

NIHMSID: [NIHMS268346](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3109996/)

3-Alkoxy-1,2-Dioxolanes: Synthesis and Evaluation as Potential Antimalarial Agents

[Charles E. Schiaffo](#),[†] [Matthias Rottman](#),^{‡§} [Sergio Wittlin](#),^{‡§} and [Patrick H. Dussault](#)^{✉,†}

[†]Department of Chemistry, University of Nebraska—Lincoln, Lincoln, Nebraska 68588, United States

[‡]Swiss Tropical and Public Health Institute, Socinstrasse 57, CH-4002 Basel, Switzerland

[§]University of Basel, CH-4003 Basel, Switzerland

[✉]Corresponding author.

*Tel: Phone: 402-472-6951. E-mail: pdussault1@unl.edu.

Received 2010 Dec 23; Accepted 2011 Jan 17.

Copyright © 2011 American Chemical Society

This article has been [cited by](#) other articles in PMC.

Abstract

[Go to:](#)

$IC_{50} = 9.2 \text{ nM}$ against
P. falciparum

A number of 3-alkoxy-1,2-dioxolanes exhibit promising levels of antimalarial activity against *Plasmodium falciparum*. A new route to the 1,2-dioxolane core is reported based on tandem peroxidation/cyclization of enones.

Keywords: Malaria, peroxide, 3-alkoxy-1,2-dioxolane, dioxolane

Malaria is a global health epidemic affecting approximately 250 million people and causing over 800000 deaths annually.¹ The peroxide artemisinin, along with closely related artesunate and arteether, are a critical part of therapies against drug-resistant strains of *P. falciparum*, the most virulent form of the disease.² Strong antimalarial activity has also been observed with a number of artemisinin analogues.^{3–7} However, reports of delayed parasite clearance in patients receiving artemisinin combination therapy (ACT) have led to renewed interest in structurally distinct classes of peroxides.⁸ Promising levels of antimalarial activity have been observed from a number of relatively simple peroxide skeletons.^{9,10–105} Struck by the high antimalarial activity of 1,2,4-

trioxolanes (ozonides), one of which is currently in phase III trials,^{105,11} we became interested in nearly isosteric 3-alkoxy-1,2-dioxolanes. We now report the synthesis of a number of alkoxydioxolanes along with in vitro antimalarial data.

The antimalarial activity of artemisinin is believed to derive from Fe(II)-mediated fragmentation of the peroxide to generate intermediate alkoxy radicals, which undergo rapid β -fragmentation or 1,5-hydrogen abstraction to generate reactive carbon radicals.^{12,13} The ozonides discussed above have also been established to undergo Fe(II)-mediated cleavage to generate carbon radicals.¹⁴ 1,2-Dioxolanes, which are nearly isosteric with ozonides, would appear to offer a promising platform for development of new peroxide antimalarials.¹⁵ However, to date, few 1,2-dioxolanes have demonstrated significant antimalarial activity.¹⁶ One reason may be the lack of an α -oxygen, resulting in a reduced propensity for β -scission in the derived alkoxy radicals.¹⁷ We postulated that suitably substituted 3-alkoxy-1,2-dioxolanes might be excellent antimalarial candidates, providing a stable framework that would undergo activation by Fe(II) to furnish α -oxygenated alkoxy radicals predisposed toward β -scission (Scheme 1).

[Scheme 1](#)

Competing Modes of Alkoxydioxolane Activation

3-Alkoxy-1,2-dioxolanes are available from the corresponding 1,2-dioxolan-3-ols via acid- or base-promoted etherification.^{18,19} We initially pursued the 3,5,5-trimethyl-1,2-dioxolan-3-ol (**1**) and 5-butyl-3,5-dimethyl-1,2-alkoxydioxolan-3-ol (**3**) core structures (Scheme 2). A number of methods have been reported for the synthesis of 1,2-dioxolanols, including base-mediated addition of hydrogen peroxide to α,β -unsaturated ketones,^{20,21} radical oxygenation of cyclopropanols,²² thermolysis of aza-hydroperoxides,²³ or addition of singlet oxygen to methylallyl aldehydes.²⁴ However, none of these methods have been generally applied. Peroxide **1** was readily prepared through base-mediated addition of hydrogen peroxide to mesityl oxide.²¹ However, as feared, application of similar conditions to the more hydrophobic 4-methyl-3-octen-2-one (**2**) mainly furnished the product of nucleophilic epoxidation.¹⁸ Fortunately, cobalt-mediated oxygenation of **2**,²⁵ originally envisaged as a route to an intermediate 3-trialkylsilylperoxy alkanone, directly afforded the desired 1,2-dioxolan-3-ol **3**. Acid-catalyzed transesterification to form the alkoxydioxolanes could be conducted on the 1,2-dioxolanol-3-ols (not shown)¹⁸ but proceeded more cleanly on the corresponding trimethylsilyl ethers (**1a** and **3a**; Table 1). Three of the alkoxydioxolanes (**5**, **7**, and **8**) demonstrated significant in vitro activity against *P. falciparum* with dioxolane **8** displaying an IC₅₀ less than 100 nM.^{26,101} On the basis of these results, we became interested in surveying influences of the following structural elements: a spirocyclic constraint at C₅/C₅, the size of the C₃ alkoxide, and the nature of the putative radical leaving group at C₃.

[Scheme 2](#)

Synthesis of Dioxolanol Core Structures

Table 1
Initial Alkoxydioxolane Synthesis and Bioassay

Entry	R ₁	R ₂	Yield (%)	IC ₅₀ (nM)
1a	H	H	20	>1000
2a	H	H	20	>1000
3a	H	H	20	>1000
4a	H	H	20	>1000
5a	H	H	20	>1000
6a	H	H	20	>1000
7a	H	H	20	>1000
8a	H	H	20	>1000

[Table 1](#)

Initial Alkoxydioxolane Synthesis and Bioassay

The synthesis of 1,2-dioxolanes incorporating a spirocyclic constraint is illustrated in Scheme 3. Prins reaction of methylenecyclohexane with acetaldehyde or phenylacetaldehyde furnished homoallyl alcohols **9** and **10**,²⁷ which

underwent Swern oxidation to furnish β,γ -unsaturated ketones **11** and **12**. Co-mediated dioxygenation (O_2 , Et_3SiH) by a modification of Isayama's protocol directly furnished 1,2-dioxolan-3-ols **13** and **14**.^{28,100} Although the dioxolanols could be carried on directly, purification was more easily conducted on the derived trimethylsilyl ethers, **13a** and **14a**. Dioxolanol **17**, which incorporates a better radical leaving group (benzyl) at C_3 , was prepared via a benzyl/methallyl ketone, as illustrated in Scheme 4.

[Scheme 3](#)

Synthesis of 5,5'-spiro-1,2-Dioxolan-3-ols

[Scheme 4](#)

Synthesis of a 3-Benzyl-1,2-dioxolan-3-ol

We were unsuccessful in applying a similar approach to a more hindered 1,2-dioxolan-3-ol (**20**) incorporating both a C_5/C_5' spirocycle and a branched side chain at C_3 (not shown). However, this target could be prepared via dioxygenation of a cyclopropanol.²² As illustrated in Scheme 5, conversion of dicyclohexylketone to the corresponding trimethylsilyl enol ether, followed by Simmons–Smith cyclopropanation, furnished trimethylsilyloxycyclopropane **18**, accompanied by varying amounts of **19**, presumably reflecting during work up. Desilylation with TBAF furnished a tetrasubstituted cyclopropanol (**19**), which could be converted to **20** by stirring in benzene under an atmosphere of oxygen.

[Scheme 5](#)

Dioxolanol via Oxidative Ring Expansion

Acid-catalyzed S_N1 transesterification of the dioxolanols or their trimethylsilyl ethers with primary alcohols furnished 3-alkoxydioxolanes **21–25** (Table 2). The alkoxydioxolanes proved to possess remarkable chemical stability, failing to react with either PPh_3 (≥ 24 h, room temperature) or $i\text{-Bu}_2AlH$ (≥ 4 h, room temperature).

dioxanol or dioxanol silyl ethers	ROH TsOH (0.1 equiv) solvent	product	Yield (%)
18	MeOH	21	80
19	MeOH	22	80
20	MeOH	23	80
21	MeOH	24	80
22	MeOH	25	80

[Table 2](#)

Additional Alkoxydioxolanes

The results of in vitro testing for compounds **21–25** (Table 2), when taken together with the results described in Table 1, suggest clear trends in terms of antimalarial structure–activity relationships. Activity is enhanced by a spirocyclohexyl constraint at C_5/C_5' and by the presence of steric bulk at C_3 . Dioxolanes **21** and **25**, which combine a spirocyclohexyl unit with either a bulky C_3 alkyl substituent or a large C_3 alkoxide displayed IC_{50} values below 10 nM.

Finally, we were interested in investigating the Fe(II) reactivity of the alkoxydioxolanes, and, in particular, the extent of cleavage of the derived alkoxy radicals (see Scheme 1). Alkoxydioxolane **25**, one of our most active of our initial leads, underwent reaction with $FeBr_2$ to furnish the corresponding 3-hydroxyester in 80% yield (reaction illustrated in Scheme 1), supporting the postulated formation and fragmentation of an alkoxy-substituted alkoxy radical.^{12,13} Similar results have been reported for a monocyclic alkoxydioxolane.²⁰

In conclusion, we have developed new and expanded routes to 1,2-dioxolan-3-ols and 3-alkoxy-1,2-dioxolanes. The latter are demonstrated to provide a highly promising platform for development of a new family of antimalarial peroxides.

Funding Statement

[Go to:](#)

National Institutes of Health, United States

Supporting Information Available

[Go to:](#)

Complete experimental procedures and characterization data; ^1H and ^{13}C NMR spectra for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes

[Go to:](#)

This work was supported by the Medicines for Malaria Venture and the Nebraska Research Initiative and conducted in facilities remodeled with support from the NIH (RR0116544-01). NMR spectra were acquired, in part, on spectrometers purchased with NSF support (MRI 0079750 and CHE 0091975). We thank Dr. Xuejun Liu for initial preparation of several dioxolanes, Tony Lloyd for technical assistance, and Prof. Jonathan Vennerstrom (University of Nebraska Medical Center) for helpful discussions.

Supplementary Material

[Go to:](#)

[ml100308d_si_001.pdf](#) (2.8M, pdf)

References

[Go to:](#)

1. World Health Organization Malaria Report 2008.
2. White N. J. Qinghaosu (Artemisinin): The Price of Success. *Science* 2008, 320, 330–334. [[PubMed](#)]
3. Moon D. K.; Singhal V.; Kumar N.; Shapiro T. A.; Posner G. H. Antimalarial preclinical drug development: A single oral dose of a 5-carbon-linked trioxane dimer plus mefloquine cures malaria-infected mice. *Drug Dev. Res.* 2010, 71, 76–81. [[PubMed](#)]
4. Walsh J. J.; Coughlan D.; Heneghan N.; Gaynor C.; Bell A. A novel artemisinin-quinine hybrid with potent antimalarial activity. *Bioorg. Med. Chem. Lett.* 2007, 17, 3599–3602. [[PubMed](#)]
5. Li Y.; Zhu Y.-M.; Jiang H.-J.; Pan J.-P.; Wu G.-S.; Wu J.-M.; Shi Y.-L.; Yang J.-D.; Wu B.-A. Synthesis and antimalarial activity of artemisinin derivatives containing an amino group. *J. Med. Chem.* 2000, 43, 1635–1640. [[PubMed](#)]
6. O'Neill P. M.; Pugh M.; Stachulski A. V.; Ward S. A.; Davies J.; Park B. K. Optimisation of the allylsilane approach to C-10 deoxo carba analogues of dihydroartemisinin: synthesis and in vitro antimalarial activity of new, metabolically stable C-10 analogues. *J. Chem. Soc., Perkin Trans. 1* 2001, 2682–2689.
7. Lin A. J.; Li L. Q.; Andersen S. L.; Klayman D. L. Antimalarial activity of new dihydroartemisinin derivatives. *J. Med. Chem.* 1992, 35, 1639–1642. [[PubMed](#)]
8. Noedl H.; Se Y.; Schaefer K.; Smith B. L.; Socheat D.; Fukuda M. M. Evidence of artemisinin-resistant malaria in western Cambodia. *N. Engl. J. Med.* 2008, 359, 2619–2620. [[PubMed](#)]
9. For an overview, see Tang Y.; Dong Y.; Vennerstrom J. L. Synthetic peroxides as antimalarials. *Med. Res. Rev.* 2004, 24, 425–448. [[PubMed](#)]
10. 1,2,4,5-Tetraoxanes: O'Neill P.; Amewu R.; Nixon G. L.; El Garah F. B.; Mungthin M.; Chadwick J.; Shone A. E.; Vivas L.; Lander H.; Barton V.; Muangnoicharoen S.; Bray P. G.; Davies J.; Park B. K.; Wittlin S.; Brun R.; Preschel M.; Zhang K.; Ward S. A. Identification of a 1,2,4,5-Tetraoxane Antimalarial Drug-Development Candidate (RKA 182) with Superior Properties to the Semisynthetic Artemisinins. *Angew. Chem., Int. Ed.* 2010, 49, 5693–5697. [[PubMed](#)]
11. Spiroperoxyketals: Ghorai P.; Dussault P. H.; Hu C. H. Synthesis of Spiro-bisperoxy ketals. *Org. Lett.* 2008, 10, 2401–2404. [[PubMed](#)]

12. 1,2,4-Trioxanes: Posner G. H.; Oh C. H.; Wang D.; Gerena L.; Milhous W. K.; Meshnick S. R.; Asawamadhasadka W. Mechanism-Based Design, Synthesis, and in vitro Antimalarial Testing of New 4-Methylated Trioxanes Structurally Related to Artemisinin. *J. Med. Chem.* 1994, 37, 1256–1258. [\[PubMed\]](#)
13. 1,2,4-Trioxolanes: Dong Y.; Wittlin S.; Sriraghavan K.; Chollet J.; Charman S. A.; Charman W. N.; Scheurer C.; Urwyler H.; Santo Tomas J.; Snyder C.; Creek D. J.; Morizzi J.; Koltun M.; Matile H.; Wang X.; Padmanilayam M.; Tang Y.; Dorn A.; Brun R.; Vennerstrom J. L. The Structure–Activity Relationship of the Antimalarial Ozonide Arterolane (OZ277). *J. Med. Chem.* 2010, 53, 481–91. [\[PubMed\]](#)
14. Fugi M. A.; Wittlin S.; Dong Y.; Vennerstrom J. L. Probing the antimalarial mechanism of artemisinin and OZ277 (arterolane) with nonperoxidic isosteres and nitroxyl radicals. *Antimicrob. Agents Chemother.* 2010, 54, 1042–1046. [\[PubMed\]](#)
15. Posner G. H.; O'Neill P. M. Knowledge of the Proposed Chemical Mechanism of Action and Cytochrome P450 Metabolism of Antimalarial Trioxanes Like Artemisinin Allows Rational Design of New Antimalarial Peroxides. *Acc. Chem. Res.* 2004, 37, 397–404. [\[PubMed\]](#)
16. Several alternatives to Fe(II)-mediated formation of reactive radicals have been proposed as the basis for the biological activity of peroxide antimalarials. For an excellent overview of this discussion, see O'Neill P. M.; Barton V. E.; Ward S. A. The Molecular Mechanism of Action of Artemisinin—The Debate Continues. *Molecules* 2010, 15, 1705–1721. [\[PubMed\]](#)
17. Tang Y.; Dong Y.; Wang X.; Sriraghavan K.; Wood J. K.; Vennerstrom J. L. Dispiro-1,2,4-trioxane Analogues of a Prototype Dispiro-1,2,4-trioxolane. *J. Org. Chem.* 2005, 70, 5103–5110. [\[PubMed\]](#)
18. Dai P.; Trullinger T. K.; Liu X.; Dussault P. H. Asymmetric Synthesis of 1,2-Dioxolane-3-acetic Acids. *J. Org. Chem.* 2006, 71, 2283–2292. [\[PubMed\]](#)
19. Martyn D. C.; Beletsky G.; Cortese J. F.; Tyndall E.; Liu H.; Fitzgerald M. M.; O'Shea T. J.; Liang B.; Clardy J. Synthesis and in vitro DMPK profiling of a 1,2-dioxolane-based library with activity against *Plasmodium falciparum*. *Bioorg. Med. Chem. Lett.* 2009, 19, 5657–5660. [\[PubMed\]](#)
20. Erhardt S.; Macgregor S. A.; McCullough K. J.; Savill K.; Taylor B. J. Model Studies of β -Scission Ring-Opening Reactions of Cyclohexyloxy Radicals. *Org. Lett.* 2007, 9, 5569–5572. [\[PubMed\]](#)
21. Dussault P. H.; Liu X. Lewis Acid-Mediated Displacements of Alkoxydioxolanes. *Org. Lett.* 1999, 1, 1391–1393. [\[PubMed\]](#)
22. Baumstark A. L.; Vasquez P. C.; Chen Y.-X. Synthesis and Thermolysis of Ketal Derivatives of 3-Hydroxy-1,2-dioxolanes. *J. Org. Chem.* 1994, 59, 6692–6696.
23. Alkoxydioxolanes have also been prepared via conjugate addition of hydroperoxyacetals to α,β -unsaturated esters: Liu H.-H.; Jin H.-X.; Wu Y.-K. Synthesis and cleavage studies of a 1,2-dioxolane-type peroxide. *Chin. J. Chem.* 2004, 22, 1029–1033.
24. Rieche A.; Schmitz E.; Grundemann E. Alkyl peroxides. XXIV. The structure of mesityl oxide peroxide. *Chem. Ber.* 1960, 93, 2443–8.
25. Gibson D. H.; DePuy C. H. Free radical reactions of cyclopropanol. The formation of 1,2-dioxolane derivatives. *Tetrahedron Lett.* 1969, 10, 2203–2206.
26. Baumstark A. L.; Vasquez P. C. Synthesis of pentasubstituted 3-hydroxy-1,2-dioxolanes. *J. Org. Chem.* 1992, 57, 393–395.
27. Adam W.; Catalani L. H.; Griesbeck A. Diastereoselective ene reaction in the photooxygenation of the silyl cyanohydrins of α,β -unsaturated aldehydes. *J. Org. Chem.* 1986, 51, 5494–5496.
28. Isayama S.; Mukaiyama T. Novel method for the preparation of triethylsilyl peroxides from olefins by the reaction with molecular oxygen and triethylsilane catalyzed by bis(1,3-diketono)cobalt(II). *Chem. Lett.* 1989, 573–6.
29. Reported in vitro antimalarial activity is based upon an average of two independent assays using the [^3H]-hypoxanthine incorporation method with the *P. falciparum* strain NF54 obtained from MR4: Huber W.; Koella J. C. A comparison of three methods of estimating EC50 in studies of drug resistance of

- malaria parasites. *Acta Trop.* 1993, 55, 257–261. [[PubMed](#)]
30. Desjardins R. E.; Canfield C. J.; Haynes J. D.; Chulay J. D. Quantitative assessment of antimalarial activity in vitro by a semiautomated microdilution technique. *Antimicrob. Agents Chemother.* 1979, 16, 710–718. [[PubMed](#)]
31. Snider B. B.; Rodini D. J.; Kirk T. C.; Cordova R. Dimethylaluminum chloride catalyzed ene reactions of aldehydes. *J. Am. Chem. Soc.* 1982, 104, 555–563.
32. Isayama S. An efficient method for the direct peroxygenation of various olefinic compounds with molecular oxygen and triethylsilane catalyzed by a cobalt(II) complex. *Bull. Chem. Soc. Jpn.* 1990, 63, 1305–10.
33. t-BuOOH activation of Co(II) catalysts for hydroazidations: Waser J.; Gaspar B.; Nambu H.; Carreira E. M. Hydrazines and Azides via the Metal-Catalyzed Hydrohydrazination and Hydroazidation of Olefins. *J. Am. Chem. Soc.* 2006, 128, 11693–11712. [[PubMed](#)]

Articles from ACS Medicinal Chemistry Letters are provided here courtesy of **American Chemical Society**