University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Faculty Publications: Department of Entomology

Entomology, Department of

3-3-2003

Genetic Sexing Research on Screwworms

Steven Skoda **USDA-ARS**

Dennis Berkebile USDA-ARS, DENNIS.BERKEBILE@ars.usda.gov

Margaret Allen **USDA-ARS**

Follow this and additional works at: http://digitalcommons.unl.edu/entomologyfacpub



Part of the Entomology Commons

Skoda, Steven; Berkebile, Dennis; and Allen, Margaret, "Genetic Sexing Research on Screwworms" (2003). Faculty Publications: Department of Entomology. Paper 19.

http://digitalcommons.unl.edu/entomologyfacpub/19

This Article is brought to you for free and open access by the Entomology, Department of at Digital Commons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



http://www.ars.usda.gov/research/publications/publications.htm?seq no 115=146950

Title: Genetic Sexing Research on Screwworms **Authors**

Skoda, Steven

■ Berkebile, Dennis

Allen, Margaret

Submitted to: APHIS-ARS-Mexico US Commission for the Eradication of Screwworm Meeting

Publication Type: Other

Publication Acceptance Date: March 3, 2003

Publication Date: March 3, 2003

Citation: SKODA, S.R., BERKEBILE, D.R., ALLEN, M.L. GENETIC SEXING RESEARCH ON SCREWWORMS. 6th Annual Meeting of the APHIS-ARS-Mexico US Commission for the Eradication of

Screwworm Meeting. 2003. Invited Oral Presentation

Interpretive Summary: Few changes have been made to the basic concepts for using Mendelian, or 'Classical', genetic techniques to develop genetic sexing strains of insects since they were first developed in the 1960's. They have been attempted for several species of insect pests with few successes. Scientists at the USDA-ARS Midwest Livestock Insects Research Unit have been developing techniques applicable to 'Classical' genetic approaches and, recently, have begun exploring the applicability of transgenic techniques for developing genetic sexing strains of screwworms. Linkage analysis has shown that genes responsible for several mutant lines are not near each other. Cytogenetic techniques are being rejuvenated for use in confirming results from both 'classical' and transgenic approaches. Screwworms have been genetically transformed to express the Green Fluorescent Protein and the resultant transgenic strains appear to be as fit as 'normal' strains. Further work is needed to determine screwworm specific lethality factors, sex determination factors must be defined, sex-specific lethality factors must be discovered, and an inducible/repressible mechanism must be developed before a reliable males-only, genetic sexing strain of screwworms can be developed for use in the mass rearing facility of the Program for the Eradication of Screwworms. Future research will emphasize developing transgenic techniques for screwworms through developing new collaborations to maximize resources.

Technical Abstract: Few changes have been made to the basic concepts for using Mendelian, or 'Classical', genetic techniques to develop genetic sexing strains of insects since they were first developed in the 1960's. They have been attempted for several species of insect pests with few successes. Scientists at the USDA-ARS Midwest Livestock Insects Research Unit have been developing techniques applicable to 'Classical' genetic approaches and, recently, have begun exploring the applicability of transgenic techniques for developing genetic sexing strains of screwworms. Linkage analysis has shown that genes responsible for several mutant lines are not near each other. Cytogenetic techniques are being rejuvenated for use in confirming results from both 'classical' and transgenic approaches. Screwworms have been genetically transformed to express the Green Fluorescent Protein and the resultant transgenic strains appear to be as fit as 'normal' strains. Further work is needed to determine screwworm specific lethality factors, sex determination factors must be defined, sex-specific lethality factors must be discovered, and an inducible/repressible mechanism must be developed before a reliable males-only, genetic sexing strain of screwworms can be developed for use in the mass rearing facility of the Program for the Eradication of Screwworms. Future research will emphasize developing transgenic techniques for screwworms through developing new collaborations to maximize resources.