

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

2000 Bird Strike Committee-USA/Canada, 2nd
Annual Meeting, Minneapolis, MN

Bird Strike Committee Proceedings

August 2000

ANALYSIS OF THE USE OF RADIO-CONTROLLED MODELS IN BIRD DISPERSAL

Nicholas B. Carter

Border Collie Rescue, Inc., 886 State Road 26, Melrose FL

Follow this and additional works at: <http://digitalcommons.unl.edu/birdstrike2000>



Part of the [Environmental Health and Protection Commons](#)

Carter, Nicholas B., "ANALYSIS OF THE USE OF RADIO-CONTROLLED MODELS IN BIRD DISPERSAL" (2000). *2000 Bird Strike Committee-USA/Canada, 2nd Annual Meeting, Minneapolis, MN*. 25.

<http://digitalcommons.unl.edu/birdstrike2000/25>

This Article is brought to you for free and open access by the Bird Strike Committee Proceedings at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2000 Bird Strike Committee-USA/Canada, 2nd Annual Meeting, Minneapolis, MN by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

ANALYSIS OF THE USE OF RADIO-CONTROLLED MODELS IN BIRD DISPERSAL

*Nicholas B. Carter, Border Collie Rescue, Inc., 886 State Road 26, Melrose FL 32666-3137USA
(352-473-0100; fax 352-473-6459; bcr@bcrescue.org)*

Border Collie Rescue (BCR), in cooperation with the Dover Air Force Base, conducted a series of field trials to test the efficacy of utilizing radio-controlled models (“RCs”) (aircraft and boats) in dispersing several species of birds from troublesome locations around the Dover AFB environment. Results of these trials suggest that RCs can be used to effectively harass seagulls, raptors, ducks, and geese from difficult areas, though there are a number of limitations that must be taken into account when determining the value of instituting a harassment program utilizing RCs. Though use of Border Collies in its wildlife management program had eliminated the majority of bird hazards at Dover AFB, a few troublesome areas and species remained, particularly those birds that were located in areas inaccessible to the dog (e.g. large quarries with heavy machinery, soaring raptors). Radio-controlled aircraft were highly effective in dispersing large flocks of loafing seagulls, raptors soaring in thermals and passing through the airfield environment, as well as flocks of ducks and teal feeding and resting on open bodies of water. They were also partially effective at discouraging seagulls and geese from utilizing the airspace over the base and seagulls feeding on earthworms on runways and taxiways, though several environmental and operational variables determined success. Radio-controlled aircraft were entirely ineffective in dispersing Canada geese from open water. Coordinated effort with a radio-controlled boat, however, proved to be an effective means of eliminating the Canada goose presence from large bodies of water.