

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

2006 Bird Strike Committee USA/Canada, 8th
Annual Meeting, St. Louis, MO

Bird Strike Committee Proceedings

August 2006

DNA IDENTIFICATION OF BIRDSTRIKE REMAINS – PROGRESS REPORT

Carla J. Dove

Smithsonian Institution, Washington, DC

Marcy Heacker

Smithsonian Institution, Washington, DC

Lee Weigt

Smithsonian Institution, Washington, DC

Follow this and additional works at: <http://digitalcommons.unl.edu/birdstrike2006>



Part of the [Environmental Health and Protection Commons](#)

Dove, Carla J.; Heacker, Marcy; and Weigt, Lee, "DNA IDENTIFICATION OF BIRDSTRIKE REMAINS – PROGRESS REPORT" (2006). *2006 Bird Strike Committee USA/Canada, 8th Annual Meeting, St. Louis, MO*. 25.
<http://digitalcommons.unl.edu/birdstrike2006/25>

This Article is brought to you for free and open access by the Bird Strike Committee Proceedings at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2006 Bird Strike Committee USA/Canada, 8th Annual Meeting, St. Louis, MO by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

(24) DNA IDENTIFICATION OF BIRDSTRIKE REMAINS – PROGRESS REPORT

Carla J. Dove, Marcy Heacker, and Lee Weigt, Smithsonian Institution, P.O. Box 37012, Washington, DC 20013-7012 USA

The FAA (William J. Hughes Technical Center), U.S. Air Force, and the Smithsonian Institution signed a five year Interagency Agreement in 2003 to develop a multi-level DNA-based identification method for bird strike remains. The tasks to be achieved include: establishment of a database of mitochondrial DNA sequences (12s, 16s, CytB and COI) for 300 species of birds; testing and refining DNA-based techniques for forensic samples; and development of user-friendly methods of collecting ‘snarge’. A separate collaboration between the Smithsonian and the University of Guelph (Canada) began in 2004 to sequence one of these target genes (COI, ‘barcode’) for all birds of the United States and Canada. This effort is now 93% complete and sequences are available for some 642 species of birds (462 provided by this project). In 2005, 83 bird strike cases (FAA, USAF) were submitted to the DNA lab for identification testing. A variety of material was tested for contamination, sample quantity and quality. The DNA method provided more specific identifications than morphological analyses alone in 66% of these cases. Twenty-one cases lacked viable DNA, but 5 of the non-viable cases contained sufficient feather material for microscopic identifications. Preliminary results show that molecular identifications are not reliable in some species of waterfowl and gulls (species known to hybridize widely) because of significant overlap in mtDNA sequences. Because of limitations in both methods, we recommend using a combination of molecular and morphological techniques for the best bird strike identification system. Testing continues on the use of a ‘DNA’ card to fix the samples in the field and a rigorous testing protocol is planned for DNA bird strike identifications during the 2006 Fall migration.