

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Human–Wildlife Interactions

Wildlife Damage Management, Internet Center for

4-1-2009

House cats as predators in the Australian environment: impacts and management

Christopher R. Dickman

University of Sydney, New South Wales, Australia, cdickman@bio.usyd.edu.au

Follow this and additional works at: <http://digitalcommons.unl.edu/hwi>



Part of the [Environmental Health and Protection Commons](#)

Dickman, Christopher R., "House cats as predators in the Australian environment: impacts and management" (2009). *Human–Wildlife Interactions*. Paper 27.

<http://digitalcommons.unl.edu/hwi/27>

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Human–Wildlife Interactions by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

House cats as predators in the Australian environment: impacts and management

CHRISTOPHER R. DICKMAN, Institute of Wildlife Research, School of Biological Sciences, University of Sydney, New South Wales 2006, Australia cdickman@bio.usyd.edu.au

Abstract: This paper provides an overview of the predatory activities of the house cat (*Felis catus*) in Australia, focusing principally on the interactions of domestic and stray cats with native species of prey. Like their free-living, or feral, counterparts, domestic cats take a broad range of prey, with small mammals, birds, and human-derived foods forming the bulk of the diet. Domestic and stray cats have contributed to declines of suburban populations of eastern barred bandicoots (*Perameles gunnii*) and superb lyrebirds (*Menura novaehollandiae*) in Victoria, Australia. The effects of cats on prey communities remain speculative. In Sydney, artificial nests placed in trees in forest remnants suffered less predation where cat activity was high rather than where it was low, indicating that cats beneficially reduced damage by introduced rats and other nest predators. However, high cat activity was associated with reduced bird diversity. Legislation to encourage responsible cat ownership has been passed in Australia; it should have positive outcomes for both wildlife conservation and cat welfare.

Key words: Australia, *Felis catus*, house cat, human–wildlife conflicts

OVER THE LAST 20 YEARS, there has been a surge of interest in the introduced house cat (*Felis catus*) in Australia and, in particular, the impact of the cat on native Australian fauna. Some studies have suggested that cats kill millions of native vertebrates each year (Paton 1990, 1991; Trueman 1991), and that certain species are represented disproportionately in the kill (Seebeck et al. 1991, Dowling et al. 1994). Predation from cats appears to have been a major contributor to declining populations of some threatened native species, such as the eastern barred bandicoot (*Perameles gunnii*) in Victoria and the rufous hare-wallaby (*Lagorchestes hirsutus*) in the Northern Territory (Dufty 1994, Gibson et al. 1994). Losses of such species have led to attempts to extirpate cats from local (<10,000 ha) or even regional (>10,000 ha) areas (Algar et al. 2002, Short et al. 2002, Short and Turner 2005), and many attempts have been made to design cat-specific traps, lures, or toxins (Algar and Burrows 2004, Wark 2004).

Other studies have argued, conversely, that the kill-rates of cats are lower than often is believed, especially in urban environments (Reark 1994), and that introduced vertebrates usually form the major part of their diet (Barratt 1997). Cats often are perceived to be beneficial as controllers of vermin on farms and rural properties (Ward 1994), and possibly to have positive effects on some native species by suppressing populations of introduced *Rattus*

spp. (Tidemann et al. 1994). In Australia, where cats were introduced in large numbers in the eighteenth and nineteenth centuries (Abbott 2002), negative impacts have been highlighted most often. The strong public affection for cats (e.g., Murray and Penridge 1997, Grayson et al. 2002) and limited empirical evidence of their actual impacts have hampered attempts to manage them effectively.

Studies of cat impact often have drawn a distinction between 2 kinds of cats. On the one hand, domestic cats have been viewed as pet or house cats that live in close connection with a household where all their ecological requirements are intentionally provided by humans (Moodie 1995). Such cats do not rely on hunting for food, but they may still impact on native fauna by their predatory activities. On the other hand, feral cats are free-living; they have minimal or no reliance on humans, and survive and reproduce in self-perpetuating populations



House cat. (Photo courtesy P. German)

(Moodie 1995). Individual cats may sometimes move between these 2 extremes, occupying the category of stray if there is partial dependency on humans for the provision of resource requirements.

In the present paper, I discuss the predatory activities of cats in Australia and review the interactions of cats with populations and communities of native prey species. I address 5 questions:

1. What do cats eat?
2. Are cats specialist or generalist predators?
3. What are the effects of cat predation on prey populations?
4. What are the effects of cat predation on prey communities?
5. What management protocols can be implemented to mitigate the impacts of cats?

I focus primarily on predation by domestic and stray cats because the impacts of feral cats on native fauna have been much studied and reviewed (e.g., Dickman 1996a, b; Risbey et al. 1999, 2000) and are currently the subject of a national threat abatement plan in Australia (Environment Australia 1999). In addition, there is emerging evidence that owned and stray cats often disperse into the natural environment and help to sustain feral populations (Denny et al. 2002, Hutchings 2003), and so contribute more to impacts on native fauna than has been realized hitherto. There has been limited documentation of the effects of domestic and stray cats on native fauna (e.g., Potter 1991, Siepen and Owens 1993, Paxton 1994). I review this information here, but also present new information from my own studies where possible.

What do cats eat in Australia?

While cats take a broad range of prey, small mammals and birds often feature most prominently in their diet. In a summary of 22 studies of the diet of feral cats from 20 localities in mainland Australia, Dickman (1996a) found that introduced rabbits (*Oryctolagus cuniculus*) and house mice (*Mus musculus*) are major dietary items in semi-arid and arid habitats, whereas marsupials predominate

in temperate forest habitats. In both forest and suburban habitats, the common ringtail possum (*Pseudocheirus peregrinus*) is depredated frequently. On islands of the Australia-Pacific region, birds often predominate in the diet of cats (Fitzgerald 1990, Dickman 1996a), although invertebrates also are a prominent part of their diet (Fitzgerald and Veitch 1985, Hayde 1992). Studies published since 1996 support the view that cats take a broad range of prey (Barratt 1997, Meek 1998, Murphy et al. 2004), including carrion, under certain conditions (Paltridge et al. 1997, Molsher et al. 1999).

Studies of cat diet in suburban and temperate forest habitats of the Sydney Basin, New South Wales, exemplify the range of prey taken by domestic and stray cats (Table 1). In the most suburban situation, at Cooper Park, Sydney, native vertebrates (small scincid lizards, birds, and the common brushtail possum [*Trichosurus vulpecula*]) together formed as little as 8–17% of the diet of cats by volume. Most of the diet was derived from human-provided sources of food. In contrast, at suburban North Head, Sydney, and in temperate forest at Olney and Kurungai on the city's northern fringe, mammals predominated in the diets, with 37 to 60% of the total volume being derived from native species (Table 1). Dietary differences among habitats probably reflect differences in prey availability. The common brushtail possum is the only terrestrial native mammal occurring in Cooper Park and surrounding areas, whereas native mammals predominate at the other sites (Matthews et al. 1999). Rabbits are likewise abundant at North Head, where they form 21% of the diet, by volume, of cats, but scarce at the other sites, where they form only 1 to 6% (Table 1). These results support previous findings (Fitzgerald 1988) that cats prey mostly on small vertebrates, especially mammals and birds. These results also are consistent with studies elsewhere in Australia (Wallis et al. 1996) and Europe (Goldschmidt-Rothschild and Lüps 1976, Borkenhagen 1979) that demonstrate large differences in the representation of prey species between suburban and less disturbed natural habitats.

Are cats specialist or generalist predators?

The predominance of small mammals and

Table 1. Diets of house cats (*Felis catus*) in suburban and temperate forest habitats of the Sydney Basin, New South Wales, shown as percentage volume of occurrence of food categories.

Food category	Suburban habitat		Forest habitat	
	Cooper Park (<i>n</i> ¹ = 37)	North Head (<i>n</i> = 24)	Olney State Forest (<i>n</i> = 12)	Kuring-gai National Park (<i>n</i> = 28)
Rabbit	1.3	21.5	1.3	6.3
House mouse		1.3	1.0	1.9
Other rodent ²	14.1	22.6	15.8	18.8
Bat				0.7
Marsupial	4.7	37.2	32.8	40.8
Bird	8.8	8.1	27.6	12.4
Reptile	3.6	3.9	4.2	6.0
Invertebrate	5.5	3.7	3.0	2.7
Scavenge	5.0		7.6	1.9
Other ³	57.0	1.7	6.6	8.5

¹Sample sizes, *n*, represent pooled results from analyses of feces at all sites and stomach contents at all sites except Cooper Park.

²In the suburban habitats, all "other rodents" ingested were introduced black rats (*Rattus rattus*), whereas in the forest habitats all "other rodents" taken were probably native bush rats (*R. fuscipes*).

³The category "other" represents cat fur, plant material, nonorganic items, unidentified materials, and foods likely to be of human origin, such as fish, bread, or commercial pet food. Full methodological details are given in Dickman (1996a).

birds in the diet of cats has led to the frequent assumption that these taxa are preyed upon selectively. However, dietary selectivity is more reliably indicated if a predator can be shown to take certain prey over others that are also available. Such selectivity may be demonstrated at the level of the individual predator or that of the population. In a particularly instructive study of population-level selectivity, Childs (1986) showed that cats in urban Baltimore, Maryland, USA, took small brown rats (*Rattus norvegicus*) weighing <200 g, while most (91%) of the rats available in the population weighed >200 g. In other studies, Barratt (1997) showed that cats in suburban Canberra, Australia, preferred house sparrows (*Passer domesticus*) and blackbirds (*Turdus merula*) to 9 other common species of birds, but avoided starlings (*Sturnus vulgaris*), possibly because starlings nest and roost high in trees and feed on the ground in flocks that can more readily detect approaching predators. Molsher et al. (1999) demonstrated that rabbits were being depredated selectively at Lake Burrendong, near Wellington, New South Wales, even after a 90% decline in rabbits' abundance due to rabbit calicivirus disease. At the individual level, cats have sometimes been

shown to selectively take certain prey species and to adopt specialized hunting methods to obtain them. Often these are species of small mammals or birds (Bradshaw 1992), but large animals such as the rufous hare-wallaby (*Lagorchestes hirsutus*; Gibson et al. 1994) and unusual prey, such as bats (Churcher and Lawton 1989) and grasshoppers (Hochstrasser 1970), also have been targeted. Observations of hunting by domestic and stray cats near human settlement at 2 locations have revealed clear specializations among individuals. On Rottneest Island, Western Australia, 3 of 5 cats appeared to be accomplished mousers, catching every house mouse (*Mus musculus*) that they hunted, but achieving only a 50% catch rate for either birds or lizards. These cats adopted a sit-and-wait strategy (Turner and Meister 1988), pouncing on mice from behind cover under conditions of semi-light or darkness. In the same population, by contrast, 1 cat achieved high rates of capture success on lizards and another cat on birds. Both foraged diurnally, the first pursuing lizard prey actively in sparse coastal heath vegetation, the second pouncing on birds from dense cover. Similar selectivity was observed by 1 cat at North Head that specialized on rabbits.

What are the effects of cat predation on prey populations?

The population-level impacts of feral cats on native fauna have been much discussed in Australia (Dickman 1996a, b) and elsewhere (King 1984, Fitzgerald 1988, 1990). Surprisingly, the impacts of domestic and stray cats have remained poorly studied in Australia, with only 2 well-documented studies.

The first study concerned the eastern barred bandicoot (*Perameles gunnii*), a small, rabbit-size marsupial, at the town of Hamilton, Victoria. Formerly widespread in southwestern Victoria and southeastern South Australia, this bandicoot had become restricted to the Hamilton area by the 1970s. During 1982 to 1983, the population comprised about 1,750 animals (Moon 1984), but had fallen to just 150 to 300 animals by 1989 (Lacy and Clark 1990). The precipitous population decline appeared to be driven by a high rate of mortality, especially of juveniles, with >42% of juvenile deaths being caused by cat predation (Dufty 1994).

The second case study concerned the superb lyrebird (*Menura novaehollandiae*) in Sherbrooke Forest, Victoria. The lyrebird population was about 130 animals in the 1960s, but had fallen to only 60 by 1988 (Bradley and Bradley 1990). Predation by cats, foxes (*Vulpes vulpes*) and domestic dogs (*Canis familiaris*) was identified as the major cause of the decline, with cats probably accounting for disproportionate mortality of young birds (Larkin 1989; H. Bradley, personal communication).

What are the effects of cat predation on prey communities?

Given the relatively small amount of information on the effects of cats on individual native species, it is not surprising that understanding of cat impacts on prey communities is meager. There is considerable speculation that cats may have indirect but deleterious effects on plant communities by reducing the abundance of avian pollinators, or by depleting rat kangaroos (*Bettongia* spp. and *Potorous* spp.) and other vectors of mycorrhizal fungi (Dickman 1996a). In contrast, circumstantial evidence suggests that cats may facilitate denser populations of forest birds by suppressing the numbers of predatory rats (Fitzgerald and Karl 1979, Ebenhard 1988, Tidemann et al. 1994).

One observational study on birds in Sydney illustrates the difficulty of disentangling the positive and negative effects of cats at the community level. In the first part of this study, in which I participated, Matthews et al. (1999) investigated the intensity of predation on artificial bird eggs and nests placed in trees in 24 forest patches throughout the Sydney metropolitan region. The predation rate, calculated as the percentage of nests attacked by predators, ranged from 45 to 100%. Avian predators were detected in all patches; black rats (*Rattus rattus*) attacked nests in 10 areas, and common ringtail possums and brown antechinus (*Antechinus stuartii*) damaged eggs in 2 areas each. In the second part of the study, I walked along foot tracks through each remnant and counted the numbers of cat feces encountered to obtain a rough index of cat activity. A plot of cat activity, expressed as feces-per-km of track, against nest predation rate revealed a strong negative correlation (Figure 1). As cats did not damage any nests in the 24 sites, the reduced levels of nest predation associated with high cat activity presumably reflect suppressive effects of cats on nest predators. Indeed, examination of the contents of the collected cat feces indicated that all of the nest predators noted above themselves fell victim to cats (Dickman, unpublished data), supporting the presumption that suppression is likely to have occurred. On their own, these findings indicate that domestic and stray cats may benefit tree-nesting birds in remnant forest patches in Sydney by reducing rates of nest predation.

In the final part of this study, the richness of all native species of birds was sampled in the same 24 forest patches by scoring species observed along foot tracks and plotting them against cat activity (Figure 2). The strong negative relationship suggests that cats reduce the total numbers of bird species that occur in forest remnants. Species that were absent from sites with high cat activity, but present elsewhere, included wrens (*Malurus cyaneus*), thornbills (*Acanthiza chrysorrhoa*, *A. pusilla*), wagtails (*Rhipidura leucophrys*), and other vulnerable small species that feed or nest close to the ground. These observations support the notion that cats depleted the avian community, presumably by direct predation. Confirmation awaits appropriate experimental studies.




Figure 1. Relationship between house cat (*Felis catus*) activity and nest predation rate in 24 remnant patches of forest in Sydney, New South Wales, Australia. Cat activity is expressed as an index based on numbers of cat feces found per km of foot tracks within patches. Nest predation rate is expressed as the percentage of artificial nests depredated per patch. Based on Matthews et al. (1999) and Dickman (unpublished data).




Figure 2. Relationship between house cat (*Felis catus*) activity and bird species richness in 24 remnant patches of forest in Sydney, New South Wales. Cat activity is expressed as an index based on numbers of cat feces found per km of foot tracks within patches. Bird species richness is expressed as the number of bird species heard or observed along foot tracks within patches pooled over 2 to 4 searches per patch.

What management protocols can be implemented to mitigate the impacts of cats?

Taken together, the findings of the above studies provide some evidence that domestic and stray cats impact negatively on some native species. If these cats help to establish

and maintain feral cat populations in less disturbed habitats outside conurbations (Denny et al. 2002, Say et al. 2003), their impacts may be more subtle, but also more pervasive, than realized. Research on this issue is continuing (Denny 2005). In built-up urban and suburban environments, it is likely that any direct impact of cats on native fauna will be secondary to the more dramatic effects of loss and modification of native vegetation by the suburbs themselves. However, in less disturbed areas adjoining reserves, national parks, or in remnants of native vegetation adjoining new residential developments, predation by domestic and stray cats on native species may be quite damaging (Barratt 1997).

In response to community perceptions about marauding cats, city councils and governments in all Australian states and territories have debated or passed by-laws to encourage responsible cat ownership (Department of Local Government 1994, Seebeck and Clunie 1998). Many municipalities also provide information packs to increase the awareness of owners about cat-wildlife interactions. By-laws vary greatly from council to council, but most include provisions for registration of pet cats, incentives for

sterilization, nighttime curfews, and stipulations for a maximum number of cats per property; some also allow for removal of unowned cats from parks and other areas of sensitive habitat.

Community surveys generally indicate strong support for legislation that promotes informed cat ownership, but weaker support for

proposals that restrict ownership or create cat-free zones (e.g., Grayson et al. 2002). Effective provisions should have twofold benefits. First, they should reduce the depletion of native wildlife in settled areas. Secondly, they should improve cat welfare by reducing the numbers of dumped, unwanted cats, and by reuniting lost pets with their owners. Despite the broad community support for education and control of cats, and the plethora of by-laws that has been passed in recent years, there has been no evaluation to date of the effectiveness of any existing programs. Such a review should be carried out as a matter of priority.

Acknowledgments

I thank C. McKechnie for assistance with much of the work described here, the Australian Research Council for funding, M. Crowther for comments on the manuscript, and T. English and S. Roy for encouragement to present this contribution.

Literature cited

- Abbott, I. 2002. Origin and spread of the cat, *Felis catus*, on mainland Australia, with a discussion of the magnitude of its early impact on native fauna. *Wildlife Research* 29:51–74.
- Algar, D., A. A. Burbidge, and G. J. Angus. 2002. Cat eradication on the Montebello Islands. Pages 14–18 in C. R. Veitch and M. N. Clout, editors. *Turning the tide: the eradication of invasive species*. Invasive Species Specialist Group of the World Conservation Union (IUCN), Auckland, New Zealand.
- Algar, D., and N. D. Burrows. 2004. Feral cat control research: Western Shield Review (February 2003). *Conservation Science Western Australia* 5:131–163.
- Barratt, D. G. 1997. Predation by house cats, *Felis catus*, in Canberra, Australia. Prey composition and preference. *Wildlife Research* 24:263–277.
- Borkenhagen, P. 1979. Zur Nahrungsökologie streunender Hauskatzen (*Felis sylvestris f. catus* Linné, 1758) aus dem Stradtbereich Kiel. *Zeitschrift für Säugetierkunde* 44:375–383.
- Bradley, H., and I. Bradley. 1990. *The lyrebird*. Wildlife Watch, Melbourne, Australia.
- Bradshaw, J. W. S. 1992. *The behaviour of the domestic cat*. C·A·B International, Wallingford, Oxon, UK.
- Childs, J. E. 1986. Size-dependent predation on rats (*Rattus norvegicus*) by house cats (*Felis catus*) in an urban setting. *Journal of Mammalogy* 67:196–199.
- Churcher, P. B., and J. H. Lawton. 1987. Predation by domestic cats in an English village. *Journal of Zoology* 212:439–455.
- Churcher, P. B., and J. H. Lawton. 1989. Beware of well-fed felines. *Natural History* 7:40–47.
- Denny, E. A. 2005. Ecology of free-living cats exploiting waste disposal sites—diet, morphometrics, population dynamics and population genetics. Dissertation, University of Sydney, Sydney, Australia.
- Denny, E., P. Yakovlevich, M. D. B. Eldridge, and C. R. Dickman. 2002. Social and genetic analysis of a population of free-living cats (*Felis catus*) exploiting a resource-rich habitat. *Wildlife Research* 29:405–413.
- Department of Local Government. 1994. *Proposals for the development of cat control legislation in Western Australia*. Department of Local Government, Perth, Australia.
- Dickman, C. R. 1996a. Overview of the impacts of feral cats on Australian native fauna. Australian Nature Conservation Agency, Canberra, Australia.
- Dickman, C. R. 1996b. Impact of exotic generalist predators on the native fauna of Australia. *Wildlife Biology* 2:185–195.
- Dowling, B., J. H. Seebeck, and K. W. Lowe. 1994. Cats and wildlife: results of a survey of wildlife admitted to shelters and animal welfare agencies in Victoria. Technical Report Series No. 134. Arthur Rylah Institute for Environmental Research, Department of Conservation and Natural Resources, Melbourne, Australia.
- Duffy, A. C. 1994. Population demography of the eastern barred bandicoot (*Perameles gunnii*) at Hamilton, Victoria. *Wildlife Research* 21:445–457.
- Ebenhard, T. 1988. Introduced birds and mammals and their ecological effects. *Viltrevy* 13:1–107.
- Environment Australia. 1999. *Threat abatement plan for predation by feral cats*. Environment Australia, Canberra, Australia.
- Fitzgerald, B. M. 1988. Diet of domestic cats and their impact on prey populations. Pages 123–147 in D. C. Turner and P. Bateson, editors. *The domestic cat: the biology of its behaviour*. Cambridge University Press, Cambridge, UK.
- Fitzgerald, B. M. 1990. Family Felidae. Pages

- 330–348 in C. M. King, editor. The handbook of New Zealand mammals. Oxford University Press, Auckland, New Zealand.
- Fitzgerald, B. M., and B. J. Karl. 1979. Foods of feral house cats (*Felis catus*) in forest of the Orongorongo Valley, Wellington. New Zealand Journal of Zoology 6:107–126.
- Fitzgerald, B. M., and C. R. Veitch. 1985. The cats of Herekopare Island, New Zealand: their history, ecology and effects on birdlife. New Zealand Journal of Zoology 12:319–330.
- Gibson, D. F., G. Lundie-Jenkins, D. G. Langford, J. R. Cole, D. E. Clarke, and K. A. Johnson. 1994. Predation by feral cats, *Felis catus*, on the rufous hare-wallaby, *Lagorchestes hirsutus*, in the Tanami Desert. Australian Mammalogy 17:103–107.
- Goldschmidt-Rothschild, B., and P. Lüps. 1976. Investigations concerning the nutritional ecology of domestic cats (*Felis catus*) which have “reverted to the wild state” in the canton of Berne (Switzerland). Revue Suisse Zoologie 83:723–735.
- Grayson, J., M. Calver, and I. Styles. 2002. Attitudes of suburban Western Australians to proposed cat control legislation. Australian Veterinary Journal 80:536–543.
- Hayde, K. A. 1992. Ecology of the feral cat *Felis catus* on Great Dog Island. Thesis, University of Tasmania, Hobart, Australia.
- Hochstrasser, G. 1970. Hauskatze frisst heuschrecken zur Sättigung. Säugetierkunde Mitteilungen 18:278.
- Hutchings, S. 2003. The diet of feral house cats (*Felis catus*) at a regional rubbish tip, Victoria. Wildlife Research 30:103–110.
- King, C. M. 1984. Immigrant killers. Oxford University Press, Oxford, New Zealand.
- Lacy, R. C., and T. W. Clark. 1990. Population viability assessment of the eastern barred bandicoot in Victoria. Pages 131–145 in T. W. Clark and J. H. Seebeck, editors. Management and conservation of small populations. Chicago Zoological Society, Brookfield, USA.
- Larkin, J. 1989. The song of Sherbrooke. Common Ground, June–July:21–25.
- Matthews, A., C. R. Dickman, and R. E. Major. 1999. The influence of fragment size and edge on nest predation in urban bushland. Ecography 22:349–356.
- Meek, P. D. 1998. Food items brought home by domestic cats *Felis catus* living in Booderee National Park, Jervis Bay. Proceedings of the Linnean Society of New South Wales 120:43–47.
- Molsher, R., A. Newsome, and C. R. Dickman. 1999. Feeding ecology and population dynamics of the feral cat (*Felis catus*) in relation to the availability of prey in central-eastern New South Wales. Wildlife Research 26:593–607.
- Moodie, E. 1995. The potential for biological control of feral cats in Australia. Australian Nature Conservation Agency, Canberra, Australia.
- Moon, B. R. 1984. Current distribution of the eastern barred bandicoot, *Perameles gunnii*, in Victoria. Technical Report Series 5. Arthur Rylah Institute for Environmental Research Department of Conservation, Forests and Lands, Melbourne, Australia.
- Murphy, E. C., R. J. Keedwell, K. P. Brown, and I. Westbrooke. 2004. Diet of mammalian predators in braided river beds in the central South Island, New Zealand. Wildlife Research 31:631–638.
- Murray, R. W., and H. E. Penridge, editors. 1997. Dogs and cats in the urban environment: a handbook of municipal pet management. Chiron Media, Brisbane, Australia.
- Paltridge, R., D. Gibson, and G. Edwards. 1997. Diet of the feral cat (*Felis catus*) in central Australia. Wildlife Research 24:67–76.
- Paton, D. C. 1990. Domestic cats and wildlife. Bird Observer 696:34–35.
- Paton, D. C. 1991. Loss of wildlife to domestic cats. Pages 64–69 in C. Potter, editor. The impact of cats on native wildlife. Australian National Parks and Wildlife Service, Canberra, Australia.
- Paxton, D. W., editor. 1994. Urban animal management. Australian Veterinary Association, Canberra, Australia.
- Potter, C., editor. 1991. The impact of cats on native wildlife. Australian National Parks and Wildlife Service, Canberra, Australia.
- Reark Research. 1994. The metropolitan domestic cat: a survey of the population characteristics and hunting behaviour of the domestic cat in Australia. Petcare Information and Advisory Service, Melbourne, Australia.
- Risbey, D. A., M. C. Calver, and J. Short. 1999. The impact of cats and foxes on the small vertebrate fauna of Heirisson Prong, Western Australia. Exploring potential impact using diet analysis. Wildlife Research 26:621–630.

- Risbey, D. A., M. C. Calver, J. Short, J. S. Bradley, and I. W. Wright. 2000. The impact of cats and foxes on the small vertebrate fauna of Heirison Prong, Western Australia. A field experiment. *Wildlife Research* 27:223–235.
- Say, L., F. Bonhomme, E. Desmarais, and D. Pontier. 2003. Microspatial genetic heterogeneity and gene flow in stray cats (*Felis catus*): a comparison of coat colour and microsatellite loci. *Molecular Ecology* 12:1669–1674.
- Seebeck, J., and P. Clunie. 1998. Predation of native wildlife by the cat (*Felis catus*). Action Statement 80. Department of Natural Resources and Environment, Melbourne, Australia.
- Seebeck, J. H., L. Greenwood, and D. Ward. 1991. Cats in Victoria. Pages 18–29 in C. Potter, editor. *The impact of cats on native wildlife*. Australian National Parks and Wildlife Service, Canberra, Australia.
- Short, J., and B. Turner. 2005. Control of feral cats for nature conservation. Population dynamics and morphological attributes of feral cats at Shark Bay, Western Australia. *Wildlife Research* 32:489–501.
- Short, J., B. Turner, and D. A. Risbey. 2002. Control of feral cats for nature conservation. Trapping. *Wildlife Research* 29:475–487.
- Siepen, G., and C. Owens, editors. 1993. *Cat management workshop proceedings*. Australian Nature Conservation Agency, Canberra and Queensland Department of Environment and Heritage, Brisbane, Australia.
- Tidemann, C. R., H. D. Yorkston, and A. J. Rus-sack. 1994. The diet of cats, *Felis catus*, on Christmas Island, Indian Ocean. *Wildlife Research* 21:279–286.
- Trueman, P. 1991. The impact of domestic and semi-domestic cats on the wildlife of southern Tasmania. Thesis, University of Tasmania, Hobart, Australia.
- Turner, D. C., and O. Meister. 1988. Hunting behaviour of the domestic cat. Pages 111–121 in D. C. Turner and P. Bateson, editors. *The domestic cat: the biology of its behaviour*. Cambridge University Press, Cambridge, UK.
- Wallis, R. L., H. Brunner, and J. H. Seebeck. 1996. Diet of red foxes and cats: their impact on fauna living in parks near Melbourne. *Victorian Naturalist* 113:300–305.
- Ward, S. 1994. Aspects of the ecology of semi-dependent farm cats, *Felis catus*. Thesis, Deakin University, Geelong, Australia.
- Wark, C. 2004. Auditory and visual lure systems: are they effective in attracting arid-zone feral cats? Thesis, University of Sydney, Sydney, Australia.



CHRISTOPHER R. DICKMAN is a professor of ecology at the School of Biological Sciences and director of the Institute of Wildlife Research, both at the University of Sydney, Australia. He is a fellow of the Royal Zoological Society of New South Wales. In his research, he investigates factors that influence the distribution and abundance of terrestrial vertebrates. He is curious to know the causes of the intriguing patterns of vertebrate distributions in the Australian fauna and is concerned that many species have declined or become extinct with the advent of European settlement. For the last 20 years, he has studied the exceptionally rich communities of small mammals and lizards of arid Australia. This research contributes to theoretical debate about the importance of biotic and physical processes in shaping population and species dynamics and to practical conservation gains. He was chair of the NSW Government Scientific Committee for seven years, sits on several scientific advisory bodies, such as the Science Advisory Committee for Earthwatch, and is a governor for WWF-Australia. He has served on editorial advisory boards for several journals, and is currently an associate editor of the *Journal of Applied Ecology* and the *Open Ecology Journal*. (Photo courtesy A. Greenville)