

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Transactions of the Nebraska Academy of Sciences
and Affiliated Societies

Nebraska Academy of Sciences

1-1-2001

THE TERRESTRIAL ISOPODS OF NEBRASKA (CRUSTACEA: ISOPODA)

William F. Rapp

Follow this and additional works at: <http://digitalcommons.unl.edu/tnas>



Part of the [Life Sciences Commons](#)

Rapp, William F, "THE TERRESTRIAL ISOPODS OF NEBRASKA (CRUSTACEA: ISOPODA)" (2001). *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*. Paper 28.

<http://digitalcommons.unl.edu/tnas/28>

This Article is brought to you for free and open access by the Nebraska Academy of Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Transactions of the Nebraska Academy of Sciences and Affiliated Societies by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

THE TERRESTRIAL ISOPODS OF NEBRASKA (CRUSTACEA: ISOPODA)

William F. Rapp

87 South Main Street
Pittsford, New York 14534

ABSTRACT

Nine species in three families of terrestrial isopods were collected by the author in Nebraska between 1948 and 1998: Oniscidae (*Oniscus asellus*), Cylisticidae (*Cylisticus convexus*), and Porcellionidae (*Armadillidium nasatum*, *A. vulgare*, *Metoponorthus pruinus*, *Porcellio laevis*, *P. scaber*, *P. spinicornis*, *Trachelipus rathkei*). Only *Armadillidium vulgare* was reported before 1948. All are native to Europe and Asia, and no species is native to the state.

† † †

Although terrestrial isopods are found in most parts of Nebraska, there are very few published records. If one consults Richardson (1905) or Van Name (1936), the two standard works on North American isopods, one will find no Nebraska records. This paper is based on specimens collected by the author between 1948 and 1998.

All species of terrestrial isopods found in Nebraska are of Old World origin and represent recent introductions. When European people migrated to North America they brought with them plants, soil and terrestrial isopods. As man moved west to Nebraska, he brought terrestrial isopods with him.

Terrestrial isopods are found in many different ecological niches. In general, they are distributed according to the moisture content of their habitats, which can vary from wet to dry determined by rainfall, and evaporation. They cannot survive in sunlight and are nocturnal and cryptozonic and are mainly active at night.

Isopods are found in areas where the soil is alkaline. Soils lacking in calcium can not support a population of isopods. Isopods break down organic vegetation, especially leaf litter, and they play an important role in the formation of humus. One species, *Armadillidium vulgare* Latr., under certain conditions is of economic importance, especially in greenhouses (Hatch 1947, Swenk 1929).

Isopods may be collected in many ways. Hand-picking from under wood, under loose bark on dead trees, under stones and many types of debris, as well as using pitfall traps, will often yield many specimens. Extraction of soil and humus by use of Berlese/Tullgren funnels will often yield isopods.

All specimens are deposited at the Illinois Natural History Survey, Champaign, Illinois.

DISTRIBUTION

The counties cited are shown in Fig. 1.

Family ONISCIDAE

Oniscus asellus Linnaeus

This species appears to be limited to extreme southeastern Nebraska. Brownsville, **Nemaha County**, 15 Nov. 1983 (under rocks along old railroad track south of village). Salem, **Richardson County**, 11 May 1975 (under boards around old mill).

Family CYLISTICIDAE

Cylisticus convexus (De Geer)—Fig. 2

Widely distributed especially south of the Platte River. Vandel (1962) states that it is common in Europe and Asia Minor. It has been transported to many parts of the world by man.

Family PORCELLIONIDAE

Trachelipus rathkei (Brandt)—Fig. 3

This is the most common and widely distributed isopod in Nebraska. It is commonly found in the eastern half of the state as a deciduous woodland species. Studies by the author (Rapp 1988) in an oak (*Quercus* sp.) woodland in southern Seward County, in which a series of one-square-meter samples of humus were extracted with a Berlese/Tullgren funnel, found a mean density of 6.7 per square meter. This translates to a theoretical population of 66,000 per hectare




Figure 1. Map of the counties of Nebraska.




Figure 2. Distribution map for *Cylisticus convenus*.




Figure 5. Distribution map for *Armadillidium vulgare*.




Figure 3. Distribution map for *Trachelipus rathkei*.




Figure 4. Distribution map for *Metoponorthus pruinosus*.

It is common in wooded areas along streams, especially in the eastern half of the state, where it is common in gardens in cities and villages. It is not found in the Sand Hills area. Vandel (1962) states that it is distributed in Central and Eastern Europe. However it is not found in the Mediterranean region or the Iberian Peninsula. It is found as far north as southwestern Finland (Lehtinen 1962).

***Metoponorthus pruinosus* (Brandt)—Fig. 4**

Some workers have placed this species in the genus *Porcellionides* Miers 1878, however Vandel (1962) has shown that it belongs in the genus *Metoponorthus* Budde-Lind 1870.

The distribution of this species is difficult to explain. It appears to be limited to cities and villages. The record from Cherry County is based on one specimen collected in a trash pile at a camp ground on Big Alkali Lake, August 14, 1980. Vandel (1962) considers that this species originated in the Mediterranean region and has been spread by man to many parts of the world.

***Porcellio laevis* Latreille**

This species is a southern form and apparently has not been able to adapt to Nebraska. It has been taken at the following stations: Chadron, **Dawes County**, May 18, 1981 (railroad yard); Maxwell, **Lincoln County**, August 21, 1974 (railroad right-of-way); Peru, **Nemaha County**, June 21, 1981 (around an old barn). Vandel (1962) considers this species to have originated in the Mediterranean region. It has become widespread in the warmer parts of the world.

***Porcellio scaber* Latreille**

To date there are only four records for this species. Keystone, **Keith County**, July 18, 1978 (under wood by railroad); Brownville, **Nemaha County**, June 16, 1976 (under stones around abandon building); Danbury, **Red Willow County**, July 7, 1981 (under wood by railroad); Valparaiso, **Saunders County**, 17 August 1976 (under boards by grain elevator).

East of the Mississippi River this is a common species around cities and villages. It appears to be a very successful colonized. It is very common in Europe and England.

***Porcellio spinicornis* Say**

This species is known from only two locations in Nebraska: **Jefferson County**, about 6 miles south of Fairbury, May 2, 1978 (under boards at abandoned farm); Beemer, **Cuming County**, October 18, 1973 (under boards along railroad). This is a European species which has been found in Eastern United States and Canada; It does not appear to be well established in North America. Vandel (1962) considers this species to have originated in western and northern Europe

***Armadillidium nasatum* Budde-Lund**

There is only one record: Auburn, **Nemaha County**, October 4, 1973. Taken under boards in an old building by the Missouri Pacific Railroad.

This species is usually found in greenhouses and caves. According to some workers, it can not overwinter in the north. It is common in greenhouses in eastern United States. According to Schultz (1961) it was first recorded in North America in 1902. Harding and Sutton (1985) report that it lives in the open in France and Spain.

***Armadillidium vulgare* (Latreille)—Fig. 5**

This species is widespread in the state, but appears to be more abundant in the southeast area. In most

areas it has been taken along railway right-of-ways, usually under boards and other debris. It is a common species east of the Mississippi River. In Europe is common and in southern England it is often a household pest. Man has spread it to many parts of the world.

IDENTIFICATION

Unfortunately, there are no modern keys to North American terrestrial isopods. Richardson (1905) and Van Name (1936, 1940, 1942) are the most detailed. Vandel (1960, 1962) is very detailed and covers most of the North American species. Muchmore (1990) has published an excellent key to North American genera.

LITERATURE CITED

- Hatch, M. H. 1947. The Chelifera and Isopoda of Washington and adjacent regions. *University of Washington Publications in Biology* 10 (5): 155–274.
- Harding, P. T., & S. L. Sutton. 1985. *Woodlice in Britain and Ireland: Distribution and Habitat*. Biological Records Centre, Institute of Terrestrial Ecology, Monks Wood Experimental Station, Abboys Ripton, Huntington, England.
- Muchmore, W. B. 1990. Terrestrial isopoda. In: D. L. Dindal, *Soil Biology Guide*. New York, Wiley: 805–817.
- Lehtinen, P. T. 1962. The isopod fauna of SW – Hame. *Lounais Hameen Luonto* 13: 38–40.
- Rapp, W. F. 1988. *Trachelipus rathkei* in North America. *ISOPODA* 2: 15–19.
- Richardson, H. 1905. *Monograph on the isopods of North America*. *USNH Bulletin* 54: 1–727.
- Schultz, George A. 1961. Distribution and establishment of a land isopod in North America. *Systematic Zoology* 10: 193–196.
- Swenk, M. N. 1929. *Armadillidium vulgare* in Nebraska. *Insect Pest Survey Bulletin* 9: 80.
- Vandel, A., 1960. Isopodes terrestres (Première partie). *Faune de France* 64: 1–416.
- . 1962. Isopodes terrestres (Deuxième partie). *Faune de France* 66: 417–931
- Van Name, W. G. 1936. The American land and freshwater isopod Crustacea. *Bulletin of the American Museum of Natural History* 71: 1–555.
- . 1940. Supplement to American isopod Crustacea. *Bulletin of the American Museum of Natural History* 77: 109–142.
- . 1942. A second supplement to the American land and freshwater isopod Crustacea. *Bulletin of the American Museum of Natural History* 80: 299–329.