

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

2008 Bird Strike Committee USA/Canada, 10th  
Annual Meeting, Orlando, Florida

Bird Strike Committee Proceedings

---

8-2008

## A Decade of United States Air Force Bat Strikes

Suzanne Peurach

*USGS Patuxent Wildlife Research Center, Smithsonian Institution, National Museum of Natural History*

Carla Dove

*USGS Patuxent Wildlife Research Center, Smithsonian Institution, National Museum of Natural History*

Laura Stepko

*US Air Force Safety Center, BASH Team, 9700 G Ave SE, Bldg 24499, Suite 266B, Kirtland AFB, NM 87117 USA*

Follow this and additional works at: <http://digitalcommons.unl.edu/birdstrike2008>



Part of the [Environmental Health and Protection Commons](#)

---

Peurach, Suzanne; Dove, Carla; and Stepko, Laura, "A Decade of United States Air Force Bat Strikes" (2008). *2008 Bird Strike Committee USA/Canada, 10th Annual Meeting, Orlando, Florida*. 35.

<http://digitalcommons.unl.edu/birdstrike2008/35>

This Article is brought to you for free and open access by the Bird Strike Committee Proceedings at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2008 Bird Strike Committee USA/Canada, 10th Annual Meeting, Orlando, Florida by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# A Decade of United States Air Force Bat Strikes

**Suzanne C. Peurach and Carla J. Dove**

USGS Patuxent Wildlife Research Center, Smithsonian Institution,  
National Museum of Natural History, PO Box 37012, MRC 111,  
Washington, DC 20013 USA

**Captain Laura Stepko**

US Air Force Safety Center, BASH Team, 9700 G Ave SE, Bldg 24499, Suite  
266B, Kirtland AFB, NM 87117 USA

From 1997 to 2007, 821 bat strikes were reported to the United States Air Force (USAF). Many samples were identified by macroscopic and/or microscopic comparisons with bat specimens housed in the Smithsonian's National Museum of Natural History, and in recent years by using molecular techniques. We received bat strike reports from 20 countries and 40 of the United States during this time frame. Forty-six percent of the strikes were identified to order; 5% were identified to family or genus; and 49% were identified to the species level. Fifty-five of the 101 bat strikes submitted for DNA analysis since 2006 resulted in species-level identifications. Twenty-five species of bat have been recorded striking USAF planes world-wide. The most commonly identified bat involved in USAF strike impacts is the Brazilian free-tailed bat (*Tadarida brasiliensis*; n=173), followed by the red bat (*Lasiurus borealis*; n=83). Bat strikes peaked during the spring and fall with more than 57% taking place between August and October; 82% of the strikes that recorded a mishap time occurred between 2100 and 0900; and more than 12% of the bat strikes were reported at or above 1000 feet. Although less than 1% of the bat strike reports indicated damage to USAF aircraft, cumulative damages for 1997-2007 totaled over \$825,000.00 and more than half of this sum is attributed to five bat strikes. Only five of the ten most damaging bat strikes were identified to the species level because we did not receive samples, or the evidence received was insufficient for identification. Improving the identification of bat species involved in aircraft strikes will increase our knowledge of bat movement, life history, and behavior, and help improve flying safety by predicting times, areas, and altitudes of increased bat movement and activity.

*Abstract of paper presented at Bird Strike Committee USA/Canada Meeting, Lake Mary and Sanford, Florida, August 18-21, 2008.*