

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural  
Research Service, Lincoln, Nebraska

---

8-1-2004

## Activation of the *Oryza sativa* non-symbiotic haemoglobin-2 promoter by the cytokinin-regulated transcription factor, ARR1

Emily Ross

*University of Nebraska-Lincoln*

Julie M. Stone

*University of Nebraska-Lincoln*, [jstone2@unl.edu](mailto:jstone2@unl.edu)

Christian Elowsky

*University of Nebraska-Lincoln*, [celowsky@unl.edu](mailto:celowsky@unl.edu)

Raul Arredondo-Peter

*Universidad Autónoma del Estado de Morelos, Avenida Universidad 1001, Colonia Chamilpa, 62210 Cuernavaca, Morelos, México*, [ra@uaem.mx](mailto:ra@uaem.mx)

Robert V. Klucas

*University of Nebraska-Lincoln*

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>



Part of the [Agricultural Science Commons](#)

---

Ross, Emily; Stone, Julie M.; Elowsky, Christian; Arredondo-Peter, Raul; Klucas, Robert V.; and Sarath, Gautam, "Activation of the *Oryza sativa* non-symbiotic haemoglobin-2 promoter by the cytokinin-regulated transcription factor, ARR1" (2004). *Publications from USDA-ARS / UNL Faculty*. 37.  
<https://digitalcommons.unl.edu/usdaarsfacpub/37>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Emily Ross, Julie M. Stone, Christian Elowsky, Raul Arredondo-Peter, Robert V. Klucas, and Gautam Sarath



RESEARCH PAPER

# Activation of the *Oryza sativa* non-symbiotic haemoglobin-2 promoter by the cytokinin-regulated transcription factor, ARR1

Emily J. H. Ross<sup>1,\*</sup>, Julie M. Stone<sup>1</sup>, Christian G. Elowsky<sup>2</sup>, Raul Arredondo-Peter<sup>3</sup>, Robert V. Klucas<sup>1,†</sup> and Gautam Sarath<sup>4</sup>

<sup>1</sup> Department of Biochemistry and Plant Science Initiative, University of Nebraska-Lincoln, Lincoln, Nebraska 68588, USA

<sup>2</sup> Johns Hopkins University, School of Medicine, Division of Gastroenterology, 720 Rutland Ave., Baltimore, MD 21205, USA

<sup>3</sup> Laboratorio de Biofísica y Biología Molecular, Facultad de Ciencias, Universidad Autónoma del Estado de Morelos, Avenida Universidad 1001, Colonia Chamilpa, 62210 Cuernavaca, Morelos, México

<sup>4</sup> USDA-ARS, Wheat, Sorghum and Forage Unit, University of Nebraska-Lincoln, Lincoln, Nebraska, USA

Received 19 November 2003; Accepted 21 June 2004

## Abstract

Using *in silico* methods, several putative phytohormone-responsive *cis*-elements in the *Oryza sativa* non-symbiotic haemoglobin (*NSHB*) 1-4 and *Arabidopsis thaliana* *NSHB*1-2 promoters have been identified. An *OsNSHB2* promoter::GUS reporter gene fusion shows tissue-specific expression in *A. thaliana*. GUS expression was observed in roots, the vasculature of young leaves, in flowers, and in the pedicel/stem junction. In transient assays, activity of the *OsNSHB2* promoter was significantly up-regulated in the presence of the cytokinin, 6-benzylaminopurine (BA). Deletion analyses indicated that the full-length promoter was required for maximal *trans*-activation in the presence of cytokinin. Mutation of the single cytokinin-regulated ARR1-binding element abolished promoter activation in response to cytokinin. Constitutive expression of ARR1 under the control of the 35S cauliflower mosaic virus promoter enhanced wild-type *OsNSHB2* promoter activity, but had no effect on the activity of the mutated promoter in the absence of cytokinin. However, overexpression of ARR1 in the presence of cytokinin resulted in super-activation of the wild-type promoter. The mutated promoter was only moderately activated in the presence of cytokinin and ARR1, indicating that the

*OsNSHB2* promoter can be regulated by the ARR1 protein, but requires other cytokinin-induced factors for optimal activation. This is the first report that identifies a *trans*-acting factor involved in the activation of a *NSHB* gene.

Key words: ARR proteins, cytokinins, gene regulation, non-symbiotic plant haemoglobin.

## Introduction

Cytokinins are hormones that have been implicated in numerous aspects of plant development, including but not limited to cell division, shoot initiation and growth, and leaf senescence (Mok and Mok, 2001). Cytokinins, as well as auxins, are able to reprogramme terminally differentiated leaf cells into stem cells and maintain indefinite shoot regeneration in tissue culture, suggesting that these hormones are master regulators of plant growth and development (Bolwell *et al.*, 1995; Hwang *et al.*, 2002; Mok and Mok, 2001).

Cytokinins are sensed by histidine protein kinase receptors (CRE1, AHK2, and AHK3) that transmit signals via histidine phosphotransfer proteins to nuclear response regulators (ARRs) to activate or repress transcription. The

\* Present address and to whom correspondence should be sent: Department of Entomology, Plant Industry 306, University of Nebraska-Lincoln, Lincoln, NE 68583, USA. Fax: +1 402 472 4687. E-mail: [eross2@unl.edu](mailto:eross2@unl.edu)

† Deceased.

members of the ARR family are classified into three subgroups, the pseudo-response regulators that lack the critical Asp acceptor site, type-A, and type-B. Type-A and type-B are subgrouped based on the architecture of their polypeptides and the cytokinin-inducible expression patterns of their transcripts (Imamura *et al.*, 1999; Kiba *et al.*, 1999). The type-A ARRs are cytokinin primary response genes that contain only a receiver domain and function as repressors of cytokinin signalling (Hwang and Sheen, 2001). The type-B ARRs contain an N-terminal receiver domain, which acts as a repressor until cytokinin-induced phosphorylation liberates it to its active conformation, and a C-terminal output domain containing a Myb-DNA binding domain (Hwang *et al.*, 2001; Sakai *et al.*, 2001). The type-B ARRs regulate the type-A ARRs, by binding to their promoters and activating transcription. Genes other than type-A ARRs that are regulated by type-B ARRs are not well understood. Recent expression profiling of cytokinin action in *A. thaliana* has revealed a subset of genes whose expression is altered by cytokinin. This work, while limited to examining the genes represented on the approximately 8300-element Affymetrix GeneChip, also identified a sequence motif enriched in the upstream regions of cytokinin-up-regulated genes (Rashotte *et al.*, 2003).

Haemoglobins (HBs) are multifunctional proteins found in a wide variety of organisms (Kundu *et al.*, 2003). Documented functions of HBs include storage and transport of oxygen (Bulow *et al.*, 1999; Giardina *et al.*, 1995), nitric oxide detoxification (Bulow *et al.*, 1999; Wittenberg *et al.*, 2002), and facilitating oxygen diffusion in symbiotic tissues (Jacobsen-Lyon *et al.*, 1995). While the physiological functions for several HBs have been identified, the function(s) of ubiquitous plant non-symbiotic haemoglobins (NSHBs) are largely unknown (Kundu *et al.*, 2003). Plants contain three known types of HBs, the symbiotic (SYMHBs), which facilitate oxygen diffusion in the nitrogen-fixing nodule, truncated HBs of unknown function, and NSHBs. The X-ray crystallographic structures of the NSHBs are strikingly similar to the SYMHBs (Hargrove *et al.*, 2000), however, these proteins differ in their biochemical properties. The NSHBs are divided into two groups; class 1 exhibit very tight oxygen binding (Arrendondo-Peter *et al.*, 1997; Hargrove *et al.*, 2000) and class 2 exhibit oxygen binding kinetics more similar to the SYMHBs (Kundu *et al.*, 2003; Trevaskis *et al.*, 1997). Therefore, the regulation of the *NSHB* genes is currently being pursued to further understand their potentially different physiological roles.

NSHBs are encoded by multigene families, which suggest multiple or essential functions. Four *NSHB* iso-genes are found in two clusters in the *Oryza sativa* genome: *OsNSHB1/3/4* and *OsNSHB2* (Lira-Ruan *et al.*, 2001), and *Arabidopsis thaliana* contains two *NSHBs* (Hunt *et al.*, 2002). Each gene has its own separate and distinct promoter, signifying potential differential expression of these

genes. Overexpression of *AtNSHB1* (class 1) can protect *A. thaliana* plants from hypoxia and enhance seedling growth, whereas *AtNSHB2* (class 2) expression is enhanced by addition of exogenous cytokinins (Hunt *et al.*, 2002). The molecular mechanisms behind these physiological observations have yet to be elucidated.

Tissue expression patterns dictated by a number of dicotyledonous *NSHB* upstream regulatory regions (promoters) have been analysed by transgenic expression of promoter::reporter gene fusions. The activities of *NSHB* promoters of *Parasponia andersonii* and *Trema tomentosa* were localized to the root meristem and vascular cylinder of transgenic tobacco (Bogusz *et al.*, 1990). The *Casuarina glauca NSHB* promoter was primarily active in *L. corniculatus* roots, including tip meristems, the vascular stele, and the parenchyma internal to the endodermis (Jacobsen-Lyon *et al.*, 1995). *AtNSHB1* and 2 promoters are active in distinct tissues under normal conditions and respond differentially to specific environmental stimuli, such as hypoxia (*AtNSHB1*) and cytokinin treatment (*AtNSHB2*) (Hunt *et al.*, 2001). These results underscore the necessity for a detailed understanding of specific promoter regulation *in planta*. Such studies are critical in deciphering the functions of individual NSHB proteins *in vivo*.

Monocot NSHBs have been immunologically localized to differentiating tissues, such as the vasculature and root cap cells, and germinating seeds (Arechaga-Ocampo *et al.*, 2001; Lira-Ruan *et al.*, 2001; Ross *et al.*, 2001). However, expression of individual monocot *NSHB* genes (class 1) is not known. Based on the expression of individual dicot *NSHBs*, it is likely that monocot *NSHBs* are also differentially expressed.

In this work, a sequence-based approach has been taken to identify putative *cis*-elements that regulate the expression of the *OsNSHB1-4* and *AtNSHB1* and 2 genes. All of the known *Os* and *At NSHB* promoters contain *cis*-regulatory elements implicated in hormone responses. Moreover, the cytokinin-activated ARR1-binding-*cis*-regulatory element was prevalent. Deletion and site-directed mutational analyses of a predicted ARR1-binding *cis*-element in the *OsNSHB2* promoter confirmed its functionality in cytokinin response. Co-expression studies of promoter::GUS fusions with the transcription factor *ARR1* (D'Agostino *et al.*, 2000) up-regulated the *OsNSHB2* promoter in the absence of cytokinin. Therefore, *OsNSHB2* can be added to the growing list of genes (other than type-A ARRs) regulated by the type-B ARRs (Rashotte *et al.*, 2003).

## Materials and methods

### Analysis of non-symbiotic haemoglobin promoters

Promoter sequences for *Oryza sativa NSHB1*, 3 and 4, and *NSHB2* (Accession nos. AF335504 and AF335503, respectively) and *Arabidopsis thaliana NSHB1* and 2 (Accession nos. U94998 and U94999 [PUBMED] and At2g16060 and At3g10520 [Arabidopsis database],

respectively) were analysed for known consensus *cis*-regulatory sequences utilizing the Signal Scan Search technology (Higo *et al.*, 1999), available at the PLACE website (<http://www.dna.affrc.go.jp/htdocs/PLACE/>). In addition, manual searches were performed for the sequence AGATT found in the promoter of ARR6 (Sakai *et al.*, 2001).

#### Cloning of the *O. sativa* NSHB2 5'-upstream regulatory sequence

*O. sativa* var. Jackson genomic DNA was extracted from 5 g of plant tissue using previously described methods (Dellaporta *et al.*, 1983). The predicted 960 bp *OsNSHB2* promoter region was amplified using 1 nM of both a forward primer engineered with a *Hind*III restriction site (underlined): 5'-AAGCTTATGGGTCGTTCTCGAAGGCT-3', and a reverse primer (Hb2Pro.RVS) engineered with an *Eco*RI restriction site (underlined): 5'-GAATTCTAAAGAGGTTGCTTC-CT-3'. PCR reactions were performed at an annealing temperature of 55 °C in a solution containing approximately 5 ng of *O. sativa* genomic DNA, 1 nM dNTPs, 1 mM MgCl<sub>2</sub>, 1× PCR buffer (Invitrogen) and 1 unit of *Taq* DNA polymerase (Invitrogen) in a final volume of 25 µl. The 'full-length' promoter was denoted *HB2p*.

Deletions from the 5' end of *HB2p* were generated by PCR. A truncated 652 bp promoter ( $\Delta 960/652p$ ;  $\Delta 960$  to 653) was amplified using a forward primer engineered with a *Hind*III site (5'-AAGCTTCCCAAGTAAACGAGGAAACGA-3') and the Hb2Pro.RVS primer. The 272 bp promoter ( $\Delta 960/272p$ ;  $\Delta 960$  to 273) was amplified using a *Hind*III-engineered forward primer (5'-AAGCTTGATATCCGCC-GTAGTCTCCA-3') and the Hb2Pro.RVS primer. The predicted ARR1-binding element (AGATT; (Sakai *et al.*, 2001) of the wild-type *HB2p* was mutated to a *Bam*HI site (GGATCC) using the megaprimer method (Sarkar and Sommer, 1990), resulting in the *hb2p* mutant promoter. The primer used to mutate this site was 5'-GCTGCATCCGGCACAGGATCCTTCGGCTCGAATGGA-3'. The resulting promoters *HB2p*,  $\Delta 960/652p$ ,  $\Delta 960/272p$ , and *hb2p* were cloned into PCR2.1-TOPO (Invitrogen), and the sequences were verified at the UNL-DNA Sequencing Facility.

#### Construction of NSHB2 promoter fusions to the GUS reporter gene

*OsNSHB2* promoter constructs were ligated into the 5'-*Hind*III and 3'-*Eco*RI sites of the plasmid pPTN134 (obtained from the UNL-Plant Transformation Facility, <http://www.biotech.unl.edu/>) immediately upstream of the *Escherichia coli*  $\beta$ -glucuronidase gene (*GUS*). The resulting plasmids, denoted *HB2p::GUS*,  $\Delta 960/652p::GUS$ ,  $\Delta 960/272p::GUS$ , and *hb2p::GUS*, were used for transient bombardment experiments. All plasmids were maintained in *E. coli* XL1-Blue cells (Stratagene).

#### Plant transformation of *A. thaliana* with the *HB2p::GUS* construct

The *HB2p::GUS* construct was ligated into the binary vector pZP212 (Hajdukiewicz *et al.*, 1994) to generate plasmid pZP2P. pZP2P was introduced into *Agrobacterium tumefaciens* strain C58C1 (pMP90) (Koncz and Schell, 1986) using triparental mating (Ditta *et al.*, 1980) with the pRK2013 plasmid in helper *E. coli* HB101 (Horsch *et al.*, 1985). pZP2P was transformed into *A. thaliana* (Col-0) using the floral dip method (Bechtold *et al.*, 1993). Transformed plants were selected on MS plates supplemented with kanamycin containing 100 mg l<sup>-1</sup> cefotaxime in 4% (w/v) top agar. The T<sub>2</sub> and T<sub>3</sub> populations of *A. thaliana* were harvested and analysed for GUS activity by incubating plant organs in the GUS substrate 5-bromo-4-chloro-3-indoyl- $\beta$ -glucuronic acid for 3 h at 37 °C and washing three times with 70% (v/v) ethanol. Light microscopy was performed on a Nikon SMZ800 microscope.

#### Detection of native Hb2 transcripts in *O. sativa* and GUS transcripts in stably transformed *A. thaliana* plants

*O. sativa* var. Jackson and *A. thaliana* plants harbouring the *Hb2p::GUS* construct were sprayed in triplicate with approximately 5–10 ml of either dH<sub>2</sub>O+0.05% Tween-20 or 3 mg l<sup>-1</sup> 6-benzylaminopurine+0.05% Tween-20 at 0 h and 1 h. Approximately 200 mg of tissue was harvested at 0, 1, and 2 h from each plant. Total RNA was isolated using Trizol (Invitrogen). A cDNA mix was obtained using reverse transcriptase (Ambion) and random hexamer primers from a starting concentration of 1 µg RNA. Samples were standardized using an 18S RNA primer pair (Ambion). Native *Hb2* transcripts in *O. sativa* were analysed using the specific oligonucleotide primers F: 5'-CCATGGCTCTCGTG-GAGGGAACAACGGCGTGTGGGGGGA-3' and R: 5'-CAATTCTCACTCAGCAGGCTTCA-3' to generate a 477 bp product. GUS transcripts in *A. thaliana* were analysed using the oligonucleotide primers F: 5'-CCACGCCGAAACACCTGGGTGGACG-3' and R: 5'-GAGAGGTTAAGCCGACAGCTGC-3' to generate a 546 bp product. PCR reactions were 95 °C 2 min, 35 cycles of 95 °C 30 s, 60 °C 30 s, 72 °C 2 min, and 72 °C for 5 min. Transcripts were analysed on a 1% agarose gel.

#### Cloning of the *A. thaliana* ARR1 cDNA

The 1863 bp coding sequence of the *ARR1* gene was amplified from a reverse-transcribed cDNA mixture from various tissues of *A. thaliana* (Col-0) using 1 nM of a forward primer engineered with a *Nco*I restriction site (underlined): 5'-CCATGGCATTG-TCTCTGCTCCGGAAGAACAACATGG-3' and a reverse primer engineered with a *Bam*HI site (underlined): 5'-GGATCCTCAAC-CGGAATGTTATCGATGGAGTATGCG-3'. PCR reactions were performed for 40 cycles at an annealing temperature of 58 °C in a solution containing approximately 5 ng of cDNA, 3% (v/v) DMSO, 1 nM dNTPs, 1 mM MgCl<sub>2</sub>, 1× PCR buffer, and 1 unit of *Taq* Precision Plus polymerase (Stratagene) in a final volume of 25 µl. The amplified product was ligated into PCR2.1 TOPO (Invitrogen) and verified by DNA sequencing. The amplified *ARR1* gene was then ligated into the *Nco*I and *Bam*HI sites of pPTN134 to produce the 35S::*ARR1* construct.

#### Transient transformation of NSHB2 promoter::GUS constructs


The 35S::*GUS* (positive control), 35S alone (negative control), *HB2p::GUS*,  $\Delta 960/652p::GUS$ ,  $\Delta 960/272p::GUS$ , or *hb2p::GUS* were introduced into sterilized tobacco (*Nicotiana tabacum* cv. Xanthii) leaf discs using microprojectile bombardment with the PDS-1000 He Biolistic Particle Delivery System (BioRad) at 1100 psi. Leaf discs had been previously incubated for 48 h on either MS agar (Murashige and Skoog, 1962) or MS agar containing 3 mg l<sup>-1</sup> 6-benzylaminopurine (BAP) (Sigma) under an 18/6 h light/dark period at 24 °C. Bombardments were performed in triplicate with 1.25 µg plasmid DNA per treatment coated on tungsten particles. The leaf discs were incubated under similar conditions as above and harvested 48 h following bombardment. Extracts of the discs were analysed using a quantitative GUS assay performed with 1 mM methylumbelliferyl glucuronide (MUG) as substrate (Jefferson, 1987). GUS activity was expressed as nmol of 4-methylumbelliferone (4-MU) released mg protein<sup>-1</sup> min<sup>-1</sup>. Protein concentrations were determined using the BCA protein assay (Pierce) using bovine serum albumin as a standard.

Statistical analysis was conducted using the PROC-GLM program of SAS 8e. Analysis of variance was performed for each treatment. Means were separated at the 5% probability level with the LSD test when a significant *F* ratio occurred ( $P \geq 0.05$ ). Significant differences are indicated by different letters above the bars.

## Results

### Detection of consensus cis-element regulatory sequences in the upstream promoter regions of the *O. sativa* and *A. thaliana* NSHB genes

The sequences of the *O. sativa* and *A. thaliana* NSHB promoters were analysed for potential consensus sequences using the PLACE database (<http://www.dna.affrc.go.jp/htdocs/PLACE/>), a collection of known plant regulatory promoter elements reported in the literature (Higo *et al.*, 1999). Because all known regulatory sequences are not found in the PLACE database, a pattern search was used to identify the cytokinin-responsive ARR1-binding element (AGATT; Sakai *et al.*, 2000). The *Os* and *At* NSHB promoters (approximately 1 kbp upstream of the translation start sites) contain a number of putative *cis*-elements known to confer regulation by plant hormones (Fig. 1). In particular, the cytokinin-responsive ARR1-binding element (Oka *et al.*, 2002; Sakai *et al.*, 2000) was found in most NSHB promoters analysed, and the consensus site determined by expression profiling (GATCTT; Rashotte *et al.*, 2003) was found in the *OsNSHB4* promoter. Each NSHB promoter also contains at least one W-box (TTTGACT) (Eulgem *et al.*, 2000), involved in response to plant defence signalling (Eulgem *et al.*, 1999; Yu *et al.*, 2001). A GCC-box (AGCCGCC) was only detected in *OsNSHB2*. This element is involved in an ethylene response to pathogens (Ohme-Takagi *et al.*, 2000) and abiotic stress (Fujimoto *et al.*, 2000). Each of the *OsNSHB* promoters, but not those from *A. thaliana* contain at least one ACG box (AACGTT), which bind members of the abscisic acid-responsive bZip family of transcriptional activators (Hobo *et al.*, 1999; Izawa *et al.*,



**Fig. 1.** The *O. sativa* and *A. thaliana* non-symbiotic haemoglobin genes possess a number of predicted *cis*-regulatory elements in their promoter regions. Shown are the predicted *cis*-regulatory regions upstream of the *OsNSHB1-4* and *AtNSHB1* and 2 coding sequences. Computational analyses of *O. sativa* and *A. thaliana* sequences were performed using analysis tools from the PLACE website (<http://www.dna.affrc.go.jp/htdocs/PLACE/>) and manual searches for *cis*-regulatory elements not present in the database. Black spot, ARR1 binding element; black square, TATA box; black cross, ACG-box; plus sign, GCC-box; black diamond, W-box; black inverted triangle, GATCTT.

1994; Toyofuku *et al.*, 1998). The presence of these candidate promoter elements suggests that the *O. sativa* and *A. thaliana* NSHB genes may be hormonally regulated.

### Activity of the *O. sativa* NSHB2 gene promoter in transgenic *A. thaliana*


The activity of a monocot (*O. sativa*) promoter in a dicot (*A. thaliana*) has been evaluated to address the conservation of NSHB promoter regulation *in planta*. The activities of the *AtNSHB1* and 2 promoters in transgenic *A. thaliana* tissues were generated using promoter::GUS fusions (Hunt *et al.*, 2001). Similarly, activity of the *OsNSHB2* gene promoter (*HB2p*) was analysed in transgenic *A. thaliana* using GUS as a reporter gene. The *OsNSHB2* promoter was chosen over *OsNSHB1*, 3, and 4 based on its unique location on the *O. sativa* genome (Lira-Ruan *et al.*, 2001), and because it contains a single putative ARR1-binding element (Fig. 1), permitting evaluation of this single *cis*-element without interference from the other identical *cis*-acting elements.

In the T<sub>2</sub> and T<sub>3</sub> generations of plants harbouring the *OsHB2p*::GUS construct, light microscopy showed strong GUS staining in root vasculature, root cap cells, and root hairs throughout plant development (Fig. 2A–C). GUS activity was observed in the cotyledon and leaf vasculature during early development (Fig. 2D, E), but is absent in cauline leaves (Fig. 2F). In young flowers, GUS expression in the short style was observed prior to the opening of the bud (Fig. 2G), and throughout flower development (Fig. 2G–I). In inflorescences with open flowers, GUS staining was evident in the vasculature of the sepals, petals, and filaments (Fig. 2G, H). In addition, elevated GUS activity was detected in pollen throughout pollen maturation and anther dehiscence (Fig. 2H). GUS activity was also evident in pedicel abscission zones in late flower development through silique formation (Fig. 2I).

Using confocal laser-scanning microscopy to visualize *HB2p*::GUS expression, GUS activity was again evident in the abscission zone of the pedicel (Fig. 2J). Staining was also observed in the nectaries (Fig. 2K), and in the short style throughout development of the flower and maturation of the silique (Fig. 2L). Finally, in areas of the plant wounded during excision, GUS staining was pronounced, indicating that the *HB2p*::GUS was activated at wound sites (Fig. 2M). Expression of the *OSNSHB2* promoter overlaps with the activities of the promoters of both *AtNSHB1* and 2 (Hunt *et al.*, 2001). These data suggest that monocot and dicot NSHB regulation is similar and that molecular components necessary for expression of a monocot promoter are conserved in a dicot.

### Activation of the *O. sativa* NSHB2 promoter by exogenous cytokinin application

Putative ARR1 binding elements are found in all of the NSHB promoters analysed (one is located greater than 1



**Fig. 2.** The *OsNSHB2* promoter is active in *A. thaliana*. Expression of the *OsNSHB2* promoter::*GUS* construct in *A. thaliana* (Col-0) T<sub>2</sub> and T<sub>3</sub> plants was analysed by staining with GUS substrate (X-Gluc) and visualized using a light microscope (A–I) and a confocal laser scanning microscope (J–M). (A) Young roots; junction of root and shoot; (B) young root tips; (C) cauline-leaf stage root tips; (D) cotyledons; (E) primary leaves; (F) cauline leaves; (G) young inflorescences; (H) an inflorescence at anthesis; (I) maturing siliques; (J) mature floral organ abscission zone; (K) nectaries and anther vasculature; (L) stigma and style; and (M) stem wound site. Arrows indicate areas of significant GUS staining.


kbp upstream of the *OsNSHB3* translational start site; data not shown). The activation of a type-A cytokinin-responsive gene *ARR6* in *A. thaliana* occurs through the binding of the transcription factor ARR1 to the ARR1-binding elements found in the *ARR6* promoter. Three ARR1-binding elements exist in the 200 bp region upstream from the *ARR6* translation start site (Sakai *et al.*, 2001), however, it is not known if all the ARR1-binding elements are required or bind to the ARR1 protein during cytokinin activation of the *ARR6* gene. Only one putative ARR1-binding element was identified in the 960 bp *OsNSHB2* promoter. The presence of only one ARR1 binding element in the *OsNSHB2* promoter allowed it to be directly tested whether a single element was adequate for gene activation by ARR1 binding to its cognate site on the *OsNSHB2* promoter.

To analyse the *OsNSHB2* promoter response to cytokinin, *HB2p*::*GUS* was used in transient transformation assays via particle bombardment. Sterile tobacco leaf discs were placed on MS agar (control) or MS agar supplemented with 6-benzylaminopurine (BA). Leaf discs were individually bombarded with (i) the negative control plasmid, containing

the CAMV 35S promoter, but no *GUS* gene; (ii) a positive control containing 35S::*GUS*; or (iii) a plasmid containing the *HB2p*::*GUS* construct. GUS activity was determined 48 h after bombardment. Figure 3 indicates that basal GUS activity for tobacco discs incubated on MS medium and bombarded with 35S, 35S::*GUS*, and *HB2p*::*GUS* was  $1.5 \pm 0.3$ ,  $11.4 \pm 1.1$ , and  $13.0 \pm 1.0$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>, respectively. The activity measured with 35S::*GUS* and *HB2p*::*GUS* were significantly different from 35S alone ( $F=25.5$ ;  $df=5, 17$ ;  $P<0.0001$ ). Inclusion of BA in the incubation medium increased GUS activity in leaf discs bombarded with the *HB2p*::*GUS* construct to  $37.5 \pm 10$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>, compared with  $1.0 \pm 0.6$  and  $11.4 \pm 3.6$  nmol mg<sup>-1</sup> protein min<sup>-1</sup> for discs bombarded with plasmids containing the 35S and 35S::*GUS* constructs, respectively. Therefore, BA application had a significant effect on the activity of the *HB2* promoter (3-fold) and no effect on the 35S promoter (Fig. 3). These results are consistent with activation of the *OsNSHB2* promoter through a cytokinin-regulatory element, like that of the putative ARR1-binding element (Fig. 1).

To identify the specific element of the *OsNSHB2* promoter that was mediating the cytokinin activation, deletions or mutations of *HB2p* fused to the *GUS* reporter gene were analysed using transient bombardment assays. Figure 4A shows the regions of the promoter that were either deleted or mutated. The first deletion ( $\Delta 960/652p$ ) removed the ACG box upstream at -689 bp. The second deletion ( $\Delta 960/272p$ ) removed all of the identified canonical *cis*-acting elements including the W-box (-448 bp), GCC-box (-436 bp) and ARR1-site (-290 bp). To determine if the ARR1-binding element was primarily responsible for the observed cytokinin activation, the ARR1-binding element was mutated to a *Bam*HI restriction site (AGATT→GGATCC) to generate the mutant promoter *hb2p*. This mutation was confirmed by DNA sequencing and restriction enzyme analysis (data not shown).


GUS activity driven by the different promoter constructs in tobacco leaf discs incubated on MS agar supplemented with BA is shown in Fig. 4B. Tissue bombardment with the negative control construct (35S) showed minimal GUS activity ( $1.0 \pm 0.6$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>). By contrast, bombardment with the positive control construct, *HB2p*::*GUS*, resulted in an approximately 30-fold increase in GUS activity ( $33.5 \pm 3.2$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>), consistent with results obtained for this construct in earlier experiments (Fig. 3). Mutation of the ARR1-binding site (*hb2p*) significantly diminished GUS activity in the presence of BA ( $7.3 \pm 3.2$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>), a 4.6-fold decrease in GUS activity compared with the activity observed in discs bombarded with the full length *HB2p*::*GUS* vector ( $F=20.16$ ;  $df=4, 14$ ;  $P<0.0001$ ). Similarly, deletion of the putative ACG-box alone ( $\Delta 960/652p::*GUS*) caused a ~2-fold decrease in GUS activity, which was significantly different from the GUS activity observed for all the other$



**Fig. 3.** The cytokinin, BA, activates the *OsNSHB2* promoter::GUS fusion in transient transformation assays. Tobacco discs incubated on either MS agar or MS agar+BA were bombarded with constructs containing 35S; 35S::GUS, or *HB2p*::GUS. GUS activity is expressed as nmol of 4-MU released  $\text{mg}^{-1}$  protein  $\text{min}^{-1}$ . Data from three independent experiments were pooled and analysed. Error bars indicate standard deviation. Means indicated by the same letter are not significantly different ( $P \geq 0.05$ , LSD-test). Main effect:  $F=25.5$ ;  $df=5, 17$ ;  $P < 0.0001$ .

promoter elements (Fig. 4B). As expected, removal of all the identified putative *cis*-acting elements ( $\Delta 960/272p$ ::GUS) resulted in GUS activity similar to the 35S negative control,  $2.8 \pm 2.2$  versus  $1.0 \pm 0.6$  nmol  $\text{mg}^{-1}$  protein  $\text{min}^{-1}$  (Fig. 4B). Analysis of *hb2p*::GUS,  $\Delta 960/652p$ ::GUS, and  $\Delta 960/272p$ ::GUS in the absence of BA resulted in GUS activity with no significant difference with the negative control lacking the GUS gene (Fig. 4B). The *HB2p*::GUS shows some activity in the absence of BA, similar to the results shown in Fig. 3, whereas the mutated and deleted versions have less. The significance levels for the negative control in the treatment lacking and containing BA were equivalent, as seen in Fig. 3. These data demonstrated a striking activation of the *HB2p*::GUS construct in the presence of cytokinins that was abolished by mutation or removal of the ARR1-binding site.

ARR1-responsive genes are induced fairly rapidly upon perception of the cytokinin signal (Hutchinson and Keiber, 2002; Oka *et al.*, 2002; Sakai *et al.*, 2001). To verify that a similar response was occurring in the *HB2p*::GUS transformed plants, plants were sprayed with water or with 3 mg  $\text{l}^{-1}$  BA in the presence of a wetting agent, Tween-20. All above-ground plant parts were harvested immediately following spraying, and 1 h and 2 h post-treatment. Cytokinin application enhanced GUS transcript levels within 1 h and were substantially elevated after 2 h, compared with the water controls (Fig. 4C). These data confirmed the early response of the *HB2p* promoter to exogenous cytokinins *in planta*. Accumulation of *HB2* transcripts was also detected in the native rice species using the same procedure (Fig. 4D).



**Fig. 4.** Deletion or mutation of the *OsNSHB2* promoter affects activation by cytokinin. (A) Deletion and mutational strategy for the *OsNSHB2* promoter region. Promoters are designated as wild-type (*HB2p*); deleted 308 bp ( $\Delta 960/652p$ ) and 688 bp  $\Delta 960/272p$ ; and full-length promoter with mutated ARR1-binding site (*hb2p*). Black spot, ARR1 binding element; black square, TATA box; black cross, ACG-box; plus sign, GCC-box; black diamond, W-box. (B) Cytokinin (BA) activation of derivations of the *OsNSHB2* promoter, analysed by transient bombardment assays. Tobacco discs were incubated on MS agar+BA and bombarded with either 35S,  $\Delta 960/272p$ ::GUS,  $\Delta 960/652p$ ::GUS, *hb2p*::GUS, or *HB2p*::GUS constructs. GUS activity is expressed as nmol of 4-MU released  $\text{mg}^{-1}$  protein  $\text{min}^{-1}$ . Data from three independent experiments were pooled and analysed. Error bars indicate standard deviation. Means indicated by the same letter are not significantly different ( $P \geq 0.05$ , LSD-test). Main effect:  $F=20.16$ ;  $df=4, 14$ ;  $P < 0.0001$ . (C) Activation of GUS in of *HB2p*::GUS carrying *A. thaliana* plants. Plants were sprayed with distilled water containing 0.05% Tween-20 (Water) or 3 mg  $\text{l}^{-1}$  BA containing 0.05% Tween-20 (BA). Total RNA was extracted at 0, 1, and 2 h after spraying and an RT reaction performed using GUS specific primers that would yield a 546 bp product. Loading was standardized using an 18S primer pair. (D) Activation of *HB2* by cytokinin in native *O. sativa*. Plants were treated in a similar manner as in


### Activation of *O. sativa* HB2p promoter by co-bombardment with 35S::ARR1

To investigate the role of the ARR1-binding element in the cytokinin-induced activation of the *OsNSHB2* promoter further, co-bombardment with the *AtARR1* gene was performed (Fig. 5). Tobacco leaf discs were co-bombarded with either *HB2p::GUS* or *hb2p::GUS* and 35S or 35S::ARR1 on MS agar and MS agar supplemented with BA. Levels of GUS activity in discs maintained on MS medium alone and co-bombarded with *HB2p::GUS*+35S, *hb2p::GUS*+35S, or *hb2p::GUS*+35S::ARR1 constructs were  $7.8 \pm 4.9$ ,  $6.2 \pm 3.8$ , and  $6.4 \pm 3.8$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>, respectively (Fig. 5). These values were statistically similar to those obtained from tobacco leaf discs maintained on MS medium supplemented with BA and co-bombarded with the *hb2p::GUS*+35S constructs ( $F=53.89$ ;  $df=7, 47$ ;  $P<0.0001$ ). Co-expression of ARR1 in leaf discs maintained on MS medium alone enhanced GUS activity driven only by the wild-type *HB2* promoter. A 3-fold increase was observed in leaf discs maintained on BA-supplemented MS medium and co-bombarded with the *HB2p::GUS*+35S constructs ( $15.2 \pm 2.9$  versus  $21.9 \pm 6.8$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>, respectively), indicating that there was a positive relationship between expression of the ARR1 gene and activity of *HB2p*. There was no effect of co-bombardment of ARR1 and the mutated *hb2p::GUS* constructs in the absence of BA (Fig. 5), suggesting that only the wild-type *HB2* promoter was able to respond to excess ARR1 in tobacco tissues.

In the presence of BA, co-bombardment of leaf discs with 35S::ARR1 with either the *HB2p::GUS* or *hb2p::GUS* constructs significantly enhanced GUS activity compared with the other treatments (Fig. 5). The highest level of GUS activity,  $60.1 \pm 10.8$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>, was observed in leaf discs incubated on MS agar supplemented with BA and co-bombarded with the wild-type, full-length promoter and 35S::ARR1. Unexpectedly, in the presence of BA, coexpression with ARR1 had a positive impact on GUS activity ( $34.0 \pm 7.9$  nmol mg<sup>-1</sup> protein min<sup>-1</sup>) driven by *hb2p*, with a mutated putative ARR1-binding element. In both instances there was an approximate 3-fold activation compared with the respective controls.

## Discussion

Plant non-symbiotic haemoglobins are encoded by multi-gene families, in species with significant genomic and/or EST sequences available. This observation raises the question of whether these proteins are essential (i.e. redundant) or have multiple physiological functions. Regard-



**Fig. 5.** Overexpression of the *A. thaliana* cytokinin response regulator, ARR1, activates the *NSHB2* promoter in the absence of exogenous cytokinin. Co-bombardment analysis of the wild-type and mutated *OsNSHB2* promoters in the absence and presence of *A. thaliana* ARR1 under the control of the cauliflower mosaic virus 35S promoter. Assays were conducted with *HB2p::GUS* or *hb2p::GUS* and 35S or 35S::ARR1. GUS activity is expressed as nmol of 4-MU released mg<sup>-1</sup> protein min<sup>-1</sup>. Data from six independent experiments were pooled and analysed. Error bars indicate standard deviation. Means indicated by the same letter are not significantly different ( $P \geq 0.05$ , LSD-test). Main effect:  $F=53.89$ ;  $df=7, 47$ ;  $P<0.0001$ .

less, a better understanding of the expression patterns of different isoforms is essential to elucidating their individual physiological functions *in planta*.

*In silico* analyses of *O. sativa* and *A. thaliana* *NSHB* promoters (1 kbp upstream of the translation start sites) has revealed a number of putative *cis*-elements, implicating that *NSHB* expression may be regulated in part through hormones and abiotic stresses (Fig. 1). These elements include putative binding sites for the Arabidopsis Response Regulators (ARRs), specifically the type-B ARR1. ARR1 has been shown to be involved in early responses to cytokinins (Oka *et al.*, 2002; Sakai *et al.*, 2001). There are three major classes of ARRs: (i) the pseudo-response regulators that lack the critical Asp acceptor site; (ii) the type-A ARRs which lack a DNA binding domain and function as repressors of cytokinin signalling; and (iii) the type-B ARRs, which contain Myb-like DNA binding domains and function as transcriptional activators of cytokinin-induced gene expression (Haberer and Kieber, 2002; Hwang *et al.*, 2002; Imamura *et al.*, 1999).

The presence of these elements is consistent with previous reports regarding the activity of the *NSHB* promoters

Fig. 4C. RT reaction performed using *HB2*-specific primers that would yield a 477 bp product. Loading was standardized using an 18S primer pair.

*in planta*. For instance, cytokinin was shown to activate some *NSHBs* (Hendriks *et al.*, 1998; Hunt *et al.*, 2001), and hypoxia induces expression from the *AtNSHB1* promoter (Hunt *et al.*, 2001; Sowa *et al.*, 1999). It is likely that a combination of these *cis*-elements and their cognate *trans*-acting factors are required for the activation/deactivation of these promoters *in planta*.

The organ and tissue patterns of GUS activity driven by the *OsNSHB2* promoter in *A. thaliana* (Fig. 2) were largely similar to those observed for plants transformed with *AtNSHB1* and *AtNSHB2* promoter::*GUS* constructs (Hunt *et al.*, 2001), suggesting conservation of signal transduction pathways in monocots and dicots. The *AtNSHB1* promoter is activated during germination, in the roots and root tips of mature plants and under hypoxic conditions. By contrast, the *AtNSHB2* promoter drives GUS activity in older plants, in floral tissues, and in response to cytokinins (Hunt *et al.*, 2001). In *A. thaliana*, the *OsNSHB2* promoter-driven *GUS* expression occurs in cell types undergoing cellular differentiation or sensing some aspect of the environment (Fig. 2). The promoter activity correlates well with the localization of *OsNSHB* proteins in several differentiating cell types (Ross *et al.*, 2001). *OsNSHB2* activity was also abundant at the sites of tissue excision where a wound response has been initiated (Bouquin *et al.*, 1997; Peck and Kende, 1998; Rushton, 2002), indicating that endogenous hormonal status regulates expression of the *OsNSHB2* gene.

Analysis of transient bombardment assays indicates that cytokinin significantly (3-fold) activates the *OsNSHB2* promoter (Fig. 3). These data are within a similar activation range of *ARR1* and *ARR2* to other target sequences (Sakai *et al.*, 2000). Partial deletion ( $\Delta 960/652p$ ) confers significantly less activation in the presence of cytokinin, deletion of the promoter to a minimal promoter ( $\Delta 960/272p$ ), and mutation of the putative *ARR1* binding site (*hb2p*) abolish cytokinin activation of the *OsNSHB2* promoter (Fig. 4). These data suggest that the *OsNSHB2* promoter is, in part, activated by a cytokinin-induced transcription factor or transcriptome, and that the putative *ARR1*-binding site is involved in this cytokinin-related transcriptional activation. Co-transformation of *ARR1* and the wild-type promoter resulted in enhanced activation in the absence of cytokinin, while a similar experiment with the mutant promoter resulted in the abolition of promoter activation. In the presence of cytokinin, activation of these promoters with *ARR1* was 3-fold higher (Fig. 5) in both cases. These data indicate that other cytokinin-induced *trans*-acting factors are involved in the activation of both promoters (Rashotte *et al.*, 2003).


Mutation of the reported *ARR1* binding site found in the promoters of *OsNSHB2* (this work) and *A. thaliana* *ARR6*, AGATT (Sakai *et al.*, 2001), to a *Bam*HI site, GGATCC, might not be expected to abolish DNA binding by *ARR1*, based on previous electrophoretic mobility shift assay

(EMSA) analyses (Sakai *et al.*, 2000), in which the authors concluded that the central 'GAT' was critical for binding. However, this mutation clearly abolished activation of the *hb2p* promoter by exogenous cytokinin application (Figs 4B, 5).

The EMSA results, based on mutations of the sequence 5'-GGATT-3', suggested that position one was irrelevant and specific mutations in positions two, three and four were critical for binding, leading to the conclusion that the central 'GAT' was critical for binding (Sakai *et al.*, 2000). However, the mutations analysed were not conservative, purine to pyrimidine and vice versa. Interestingly, non-conservative mutations in position 5 (from T to G and T to A; pyrimidine to purine) also abolished binding, while a conservative mutation in this position (T to C; pyrimidine to pyrimidine) actually enhanced binding. Therefore, the nature of the nucleotide at various positions seems critical. These observations raise doubts as to the importance of the central 'GAT' sequence. It is possible that conservative mutations of nucleotides two, three, or four would also have no effect on DNA binding. Therefore, the effect of the mutation cannot be accurately predicted from these experiments. However, it is clear that the mutation made in the *hb2* promoter resulted in the abolition of cytokinin responsiveness (Fig. 5).

A recent paper describing expression profiling of cytokinin action in *A. thaliana* identified a 5'-GATCTT-3' consensus sequence prevalent in cytokinin-up-regulated gene promoters (Rashotte *et al.*, 2003). Because the Affymetrix GeneChip used (8300 genes) did not contain *AtNSHB1* and 2, the potential up-regulation of these genes for cytokinin relative to other identified genes could not be ascertained.

A model for the activation of the *OsNSHB2* promoter by *ARR1* and cytokinin is shown in Fig. 6. It has previously been suggested that native *ARR1* is repressed on its receiver domain and that liberation of *ARR1* from the endogenous repressor could be the result of a cytokinin-dependent phosphotransfer (Hwang *et al.*, 2002, 2001; Oka *et al.*, 2002). Therefore, in the absence of exogenous cytokinin, overexpression of *ARR1* is sufficient to mimic cytokinin responses, thereby resulting in a moderate *trans*-activation of the wild-type promoter. The application of exogenous cytokinin induces other factors that amplify the cytokinin signal transduction pathway through a feed-forward mechanism in which cytokinin-dependent phosphorylation of *ARR1* results in deactivation of the repressor. Activation of the cytokinin-responsive signalling mechanisms, coupled to the overexpression of *ARR1*, results in significantly higher activation of the *OsNSHB2* promoter. This model correlates well to the model of *ARR1* activation proposed by others (Hwang *et al.*, 2002, 2001; Oka *et al.*, 2002). In addition, the data suggest that competent *ARR1* binding to the *ARR1 cis*-element is required for maximal activation of the *OsNSHB2* promoter as previously



**Fig. 6.** Model for the activation of the *OsNSHB2* promoter by *A. thaliana* ARR1. (A) In the absence of cytokinin (BA), overexpression of the *A. thaliana* ARR1 protein activates the *OsNSHB2* promoter; and (B) mutation of the putative ARR1 binding site abolishes this activation. (C) Overexpression of ARR1 has a dramatic effect on the activity of the wild-type promoter in the presence of cytokinin; and (D) a moderate (but significant) effect on the activity of the mutated *hb2p* promoter in the presence of cytokinin.

suggested (Sakai *et al.*, 2000). Further dissection of the mechanism of ARR1 gene activation will give a better understanding of the regulation of *NSHB* gene expression by cytokinins.

Figure 1 supports the cytokinin-regulation of *AtNSHB2*, which has already been reported by Hunt *et al.* (2001). However, an equivalent number of ARR1 binding sites were also detected in the promoter of *AtNSHB1*, which are not in close proximity (further than 300 bp) from the start site. This promoter, however, was reported not to be activated by cytokinins in the same study (Hunt *et al.*, 2001). Perhaps the close proximity of the ARR1 binding sites to the translation start sites, as noted for the *ARR6* promoter by Sakai *et al.* (2001), are responsible for the observed cytokinin-mediated activation of both the *OsNSHB2* (this work) and the *A. thaliana NSHB2* (Hunt *et al.*, 2001). If so, it might be hypothesized that the *OsNSHB1*, but not *OsNSHB3* or *OsNSHB4*, is also activated by cytokinin (Fig. 1). A test of this hypothesis remains to be performed.

*OsNSHB2* is considered a class 1 NSHB based on phylogenetic analysis and oxygen-binding kinetics. Class 1 NSHBs are reported as being active in germination and roots and root tips under hypoxia (Hunt *et al.*, 2001). Class 2 NSHBs are reported as proteins that bind oxygen in a more similar fashion to the symbiotic HBs, are expressed during later development in floral tissues, and are induced by cytokinins. The *OsNSHB2* promoter is interestingly

induced by cytokinins and is activated during all of development in tissues where *AtNSHB1* and 2 are activated. This suggests that, perhaps, the NSHB classification system requires more parameters and that functions of the two classes may overlap in some cases.

Cytokinins are involved in several differentiation pathways (Mok and Mok, 2001). Differentiation processes result in metabolic redirection and generally require high levels of energy and a high demand for oxygen for maintenance of cellular integrity during the differentiation process. Thus, NSHBs may function to maintain a specific cellular microenvironment that is part of a cellular protection mechanism during differentiation (Kundu *et al.*, 2003; Ross *et al.*, 2001). NSHBs could sense changes in the cellular energy status and may be involved in a signalling cascade that assists in the differentiation process. This understanding that *NSHBs* are activated in the presence of cytokinins by ARR1 through the ARR1-binding element can now be explored using the plethora of available *A. thaliana* mutants.

### Acknowledgements

The authors would like to thank Dr Tiffany Heng-Moss of UNL for aid in statistical analysis. This work was supported by in part by NIH Grant Number 1 P20 RR16469 from the BRIN Program of the National Center for Research Resources (GS) and by the University of Nebraska-Lincoln Agriculture Research Division. This is a contribution of the University of Nebraska Agricultural Research

Division, Lincoln, NE 68583. Journal Series No. 13844. We would like to dedicate this work to the late Dr Robert V Klucas.

## References

- Arechaga-Ocampo E, Saenz-Rivera J, Sarath G, Klucas RV, Arredondo-Peter R. 2001. Cloning and expression analysis of hemoglobin genes from maize (*Zea mays* ssp. *mays*) and teosinte (*Zea mays* ssp. *parviglumis*). *Biochimica et Biophysica Acta* **1522**, 1–8.
- Arredondo-Peter R, Hargrove MS, Sarath G, Moran JF, Lohrman J, Olson JS, Klucas RV. 1997. Rice hemoglobins. Gene cloning, analysis, and O<sub>2</sub>-binding kinetics of a recombinant protein synthesized in *Escherichia coli*. *Plant Physiology* **115**, 1259–1266.
- Bechtold N, Ellis J, Pelletier G. 1993. In planta *Agrobacterium* gene transfer by infiltration of adult *Arabidopsis thaliana* plants. *CR Academy Science Paris, Life Sciences* **316**, 1194–1199.
- Bogusz D, Llewellyn DJ, Craig S, Dennis ES, Appleby CA, Peacock WJ. 1990. Nonlegume hemoglobin genes retain organ-specific expression in heterologous transgenic plants. *The Plant Cell* **2**, 633–641.
- Bolwell GP, Butt VS, Davies DR, Zimmerlin A. 1995. The origin of the oxidative burst in plants. *Free Radical Research* **23**, 517–532.
- Bouquin T, Lasserre E, Pradier J, Pech JC, Balague C. 1997. Wound and ethylene induction of the ACC oxidase melon gene CM-ACO1 occurs via two direct and independent transduction pathways. *Plant Molecular Biology* **35**, 1029–1035.
- Bulow L, Holmberg N, Lilius G, Bailey JE. 1999. The metabolic effects of native and transgenic hemoglobins on plants. *Trends in Biotechnology* **17**, 21–24.
- D'Agostino IB, Deruere J, Kieber JJ. 2000. Characterization of the response of the *Arabidopsis* response regulator gene family to cytokinin. *Plant Physiology* **124**, 1706–1717.
- Dellaporta SL, Wood J, Hicks JB. 1983. A plant DNA mini-preparation: version II. *Plant Molecular Biology Reporter* **1**, 19–21.
- Ditta G, Stanfield S, Corbin D, Helinski DR. 1980. Broad-host-range DNA cloning system for gram-negative bacteria: Construction of a gene bank of *Rhizobium meliloti*. *Proceedings of the National Academy of Sciences, USA* **27**, 7347–7451.
- Eulgem T, Rushton PJ, Robatzek S, Somssich IE. 2000. The WRKY superfamily of plant transcription factors. *Trends in Plant Science* **5**, 199–206.
- Eulgem T, Rushton PJ, Schmelzer E, Hahlbrock K, Somssich IE. 1999. Early nuclear events in plant defence signalling: rapid gene activation by WRKY transcription factors. *EMBO Journal* **18**, 4689–4699.
- Fujimoto SY, Ohta M, Usui A, Shinshi H, Ohme-Takagi M. 2000. *Arabidopsis* ethylene-responsive element binding factors act as transcriptional activators or repressors of GCC box-mediated gene expression. *The Plant Cell* **12**, 393–404.
- Giardina B, Messana I, Scatena R, Castagnola M. 1995. The multiple functions of hemoglobin. *Critical Reviews in Biochemistry and Molecular Biology* **30**, 165–196.
- Haberer G, Kieber JJ. 2002. Cytokinins. New insights into a classic phytohormone. *Plant Physiology* **128**, 354–362.
- Hajdukiewicz P, Svab Z, Maliga P. 1994. The small, versatile pPZP family of *Agrobacterium* binary vectors for plant transformation. *Plant Molecular Biology* **25**, 989–994.
- Hargrove MS, Brucker EA, Stec B, Sarath G, Arredondo-Peter R, Klucas RV, Olson JS, Phillips Jr GN. 2000. Crystal structure of a nonsymbiotic plant hemoglobin. *Structure Fold Design* **8**, 1005–1014.
- Hendriks T, Scheer I, Quillet MC, Randoux B, Delbreil B, Vasseur J, Hilbert JL. 1998. A nonsymbiotic hemoglobin gene is expressed during somatic embryogenesis in *Cichorium*. *Biochimica et Biophysica Acta* **1443**, 193–197.
- Higo K, Ugawa Y, Iwamoto M, Korenaga T. 1999. Plant cis-acting regulatory DNA elements (PLACE) database: 1999. *Nucleic Acids Research* **27**, 297–300.
- Hobo T, Kowayama Y, Hattori T. 1999. A bZIP factor, TRAB1, interacts with VP1 and mediates abscisic acid-induced transcription. *Proceeding of the National Academy of Sciences, USA* **96**, 15348–15353.
- Horsch RB, Fry JE, Hoffmann NL, Eichholtz D, Rogers SG, Fraley RT. 1985. A simple and general method for transferring genes into plants. *Science* **227**, 1229–1231.
- Hunt PW, Klok EJ, Trevaskis B, Watts RA, Ellis MH, Peacock WJ, Dennis ES. 2002. Increased level of hemoglobin 1 enhances survival of hypoxic stress and promotes early growth in *Arabidopsis thaliana*. *Proceeding of the National Academy of Sciences, USA* **99**, 17197–17202.
- Hunt PW, Watts RA, Trevaskis B, Llewellyn DJ, Burnell J, Dennis ES, Peacock WJ. 2001. Expression and evolution of functionally distinct haemoglobin genes in plants. *Plant Molecular Biology* **47**, 677–692.
- Hutchinson C, Keiber J. 2002. Cytokinin signaling in *Arabidopsis*. *The Plant Cell* **14**, Supplement, S47–S59.
- Hwang I, Chen HC, Sheen J. 2002. Two-component signal transduction pathways in *Arabidopsis*. *Plant Physiology* **129**, 500–515.
- Hwang I, Sheen J. 2001. Two-component circuitry in *Arabidopsis* cytokinin signal transduction. *Nature* **413**, 383–389.
- Imamura A, Hanaki N, Nakamura A, Suzuki T, Taniguchi M, Kiba T, Ueguchi C, Sugiyama T, Mizuno T. 1999. Compilation and characterization of *Arabidopsis thaliana* response regulators implicated in His-Asp phosphorelay signal transduction. *Plant Cell Physiology* **40**, 733–742.
- Izawa T, Foster R, Nakajima M, Shimamoto K, Chua NH. 1994. The rice bZIP transcriptional activator RITA-1 is highly expressed during seed development. *The Plant Cell* **6**, 1277–1287.
- Jacobsen-Lyon K, Jensen EO, Jorgensen JE, Marcker KA, Peacock WJ, Dennis ES. 1995. Symbiotic and nonsymbiotic hemoglobin genes of *Casuarina glauca*. *The Plant Cell* **7**, 213–223.
- Jefferson RA. 1987. Assaying chimeric genes in plants: the GUS gene fusion system. *Plant Molecular Biology Reporter* **5**, 387–405.
- Kiba T, Taniguchi M, Imamura A, Ueguchi C, Mizuno T, Sugiyama T. 1999. Differential expression of genes for response regulators in response to cytokinins and nitrate in *Arabidopsis thaliana*. *Plant and Cell Physiology* **40**, 767–771.
- Koncz C, Schell J. 1986. The promoter of T<sub>L</sub>-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of *Agrobacterium* binary vector. *Molecular and General Genetics* **204**, 383–396.
- Kundu S, Premer SA, Hoy JA, Trent III JT, Hargrove MS. 2003. Direct measurement of equilibrium constants for high-affinity hemoglobins. *Biophysical Journal* **84**, 3931–3940.
- Lira-Ruan V, Sarath G, Klucas RV, Arredondo-Peter R. 2001. Synthesis of hemoglobins in rice (*Oryza sativa* var. Jackson) plants growing in normal and stress conditions. *Plant Science* **161**, 279–287.
- Mok DW, Mok MC. 2001. Cytokinin metabolism and action. *Annual Review of Plant Physiology and Plant Molecular Biology* **52**, 89–118.
- Murashige T, Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum* **15**, 473–497.

- Ohme-Takagi M, Suzuki K, Shinshi H.** 2000. Regulation of ethylene-induced transcription of defense genes. *Plant and Cell Physiology* **41**, 1187–1192.
- Oka A, Sakai H, Iwakoshi S.** 2002. His-Asp phosphorelay signal transduction in higher plants: receptors and response regulators for cytokinin signaling in *Arabidopsis thaliana*. *Genes and Genetic Systems* **77**, 383–391.
- Peck SC, Kende H.** 1998. Differential regulation of genes encoding 1-aminocyclopropane-1-carboxylate (ACC) synthase in etiolated pea seedlings: effects of indole-3-acetic acid, wounding, and ethylene. *Plant Molecular Biology* **38**, 977–982.
- Rashotte A, Carson S, To J, Kieber J.** 2003. Expression profiling of cytokinin action in *Arabidopsis*. *Plant Physiology* **132**, 1998–2011.
- Ross EJ, Shearman L, Mathiesen M, Zhou YJ, Arredondo-Peter R, Sarath G, Klucas RV.** 2001. Nonsymbiotic hemoglobins in rice are synthesized during germination and in differentiating cell types. *Protoplasma* **218**, 125–133.
- Rushton P.** 2002. Exciting prospects for plants with greater disease resistance. *Trends in Plant Science* **7**, 325.
- Sakai H, Aoyama T, Oka A.** 2000. *Arabidopsis* ARR1 and ARR2 response regulators operate as transcriptional activators. *The Plant Journal* **24**, 703–711.
- Sakai H, Honma T, Aoyama T, Sato S, Kato T, Tabata S, Oka A.** 2001. ARR1, a transcription factor for genes immediately responsive to cytokinins. *Science* **294**, 1519–1521.
- Sarkar G, Sommer SS.** 1990. The 'megaprimer' method of site-directed mutagenesis. *BioTechniques* **8**, 404–407.
- Sowa AW, Guy PA, Sowa S, Hill RD.** 1999. Nonsymbiotic haemoglobins in plants. *Acta Biochimica Poland* **46**, 431–445.
- Toyofuku K, Umemura T, Yamaguchi J.** 1998. Promoter elements required for sugar-repression of the RAmy3D gene for alpha-amylase in rice. *FEBS Letters* **428**, 275–280.
- Trevaskis B, Watts RA, Andersson CR, Llewellyn DJ, Hargrove MS, Olson JS, Dennis ES, Peacock WJ.** 1997. Two hemoglobin genes in *Arabidopsis thaliana*: the evolutionary origins of leghemoglobins. *Proceeding of the National Academy of Sciences, USA* **94**, 12230–12234.
- Wittenberg JB, Bolognesi M, Wittenberg BA, Guertin M.** 2002. Truncated hemoglobins: a new family of hemoglobins widely distributed in bacteria, unicellular eukaryotes, and plants. *Journal of Biological Chemistry* **277**, 871–874.
- Yu D, Chen C, Chen Z.** 2001. Evidence for an important role of WRKY DNA binding proteins in the regulation of NPR1 gene expression. *The Plant Cell* **13**, 1527–1540.