## ON A THEOREM OF HÖLDER

DONALD W. MILLER, University of Wisconsin and University of Nebraska

1. Introduction. A well-known result, due to Hölder [1], is the following: The symmetric group  $S_n$  has outer automorphisms if and only if n=6. The classical proof of the existence of a class of outer automorphisms of  $S_6$ , as formulated by Burnside [2], rests in part on the theory of primitive groups and entails extensive computation. In this note we offer a direct method for constructing such automorphisms.

The author is grateful to Professor R. H. Bruck for raising this problem and for subsequent helpful remarks.

2. Construction of an outer automorphism of  $S_6$ . Let  $S_6$  be defined on the set  $M = \{1, 2, 3, 4, 5, 6\}$ ; let I denote the identity of  $S_6$ . Call two elements of  $S_6$  disjoint if no element of M is displaced by both of them.

Define the mapping  $\psi$  by:  $(1\ 2)\psi = (1\ 2)(3\ 6)(4\ 5) = P_2$ ,  $(1\ 3)\psi = (1\ 3)(2\ 4)(5\ 6)$ =  $P_3$ ,  $(1\ 4)\psi = (1\ 4)(2\ 6)(3\ 5) = P_4$ ,  $(1\ 5)\psi = (1\ 5)(2\ 3)(4\ 6) = P_5$ ,  $(1\ 6)\psi = (1\ 6)(2\ 5)(3\ 4) = P_6$ . Write  $N = \{2, 3, 4, 5, 6\}$ ,  $\mathcal{O} = \{P_i | i \in \mathbb{N}\}$ . Note that the elements of  $\mathcal{O}$  include as factors the 15 distinct transpositions of  $S_6$ ; consequently  $\mathcal{O}$  is transitive on M. Moreover, for  $i, j, k \in M, i \neq j$ ,

$$P_i^2 = I, \quad kP_i \neq kP_j, \quad iP_j \neq i.$$

If i, j, k are distinct elements of N, then

$$iP_j = jP_k = kP_i$$

cannot hold. For, if so, write  $iP_j = q$  and  $N = \{i, j, k, q, r\}$ . Now  $q = fP_r$  for some f in M. Certainly f is not one of i, j, k, or q. But if f = r then  $q = rP_r = 1$ , contradicting  $i \neq j$ .

If  $P_i$ ,  $P_j$ ,  $P_k$  are distinct elements of  $\mathcal{O}$ , then

$$(2) (P_i P_k P_j) P_i = P_j (P_i P_k P_j).$$

It is sufficient to prove that  $P_k$  commutes with  $P_iP_jP_i$ , for then  $P_kP_iP_jP_i=P_iP_jP_iP_iP_i$ ,  $P_iP_kP_jP_i=P_jP_iP_kP_jP_i$ ,  $P_iP_kP_jP_i=P_jP_iP_kP_jP_j$ ,  $P_iP_kP_jP_i=P_jP_iP_kP_j$ . Now

$$Q = P_i P_j P_i = P_j P_i P_j = (1 \quad i P_j P_i)(i \quad j P_i)(j \quad i P_j).$$

Each of the three transpositions of Q is a factor of some  $P_k$ ,  $k \neq i$ , j. If Q should have two cycles in common with some  $P_t$  then  $Q = P_t$ . But in that case the dis-

played representation of Q would yield  $iP_j = jP_t$ ,  $iP_jP_i = t$  (so  $iP_j = tP_i$ ), whence  $tP_i = iP_j = jP_t$ , contradicting (1). (Thus we can write  $Q = (a \ b)(c \ d)(e \ f)$ ,  $P_k = (a \ b)(c \ f)(d \ e)$ . But then  $QP_k = (c \ e)(d \ f) = P_kQ$ .

If  $A_1, \dots, A_n, B, C$  are distinct elements of  $\mathcal{O}$ , then

(3) 
$$B(CA_1 \cdot \cdot \cdot A_n B) = (CA_1 \cdot \cdot \cdot A_n B)C.$$

If n=1, (3) follows from (2). Assume inductively that (3) holds for n; then  $B(CA_1 \cdot \cdot \cdot A_n A_{n+1} B)$ 

$$= B(CA_1 \cdot \cdot \cdot A_n B)(BA_{n+1} B) = (CA_1 \cdot \cdot \cdot A_n BC)(A_{n+1} BA_{n+1})$$

$$= (CA_1 \cdot \cdot \cdot A_n A_{n+1})(A_{n+1}BCA_{n+1})BA_{n+1} = (CA_1 \cdot \cdot \cdot A_n A_{n+1})(BCA_{n+1}B)BA_{n+1}$$

$$= (CA_1 \cdot \cdot \cdot A_n A_{n+1} B) C.$$

Further, if  $A_1, \dots, A_n, B, C$  are distinct elements of  $\mathcal{O}$ , then

(4) 
$$CB(A_1 \cdot \cdot \cdot A_n)B = B(A_1 \cdot \cdot \cdot A_n)BC.$$

For by (3),  $CBA_1 \cdot \cdot \cdot A_nB = BC(BCA_1 \cdot \cdot \cdot A_nB) = BC(CA_1 \cdot \cdot \cdot A_nBC) = B(A_1 \cdot \cdot \cdot A_nBC)$ .

Define the mapping  $\theta$  as follows. Let  $a_1, \dots, a_n$  be distinct elements of N and write  $(1 \ a_i)\psi = A_i$ . Then set

(5) 
$$I\theta = I, \qquad (1a_1 \cdot \cdot \cdot \cdot a_n)\theta = A_1 \cdot \cdot \cdot \cdot A_n, \\ (a_1a_2 \cdot \cdot \cdot \cdot a_n)\theta = A_n A_1 A_2 \cdot \cdot \cdot \cdot A_n, \qquad (QR)\theta = (Q\theta)(R\theta),$$

where Q, R are arbitrary disjoint cycles of  $S_6$ . By (3),

$$(a_1a_2\cdot\cdot\cdot a_n)\theta=A_1A_2\cdot\cdot\cdot A_nA_1.$$

Clearly  $\theta$  maps  $S_6$  into itself.

To show that  $\theta$  is single-valued it will be sufficient to establish that if  $Q = (a_1 \cdot \cdot \cdot a_m)$ ,  $R = (b_1 \cdot \cdot \cdot b_n)$  are arbitrary disjoint cycles in  $S_6$ , then

- (i)  $(QR)\theta = (RQ)\theta$ ;
- (ii)  $(a_1a_2 \cdot \cdot \cdot a_m)\theta = (a_2a_3 \cdot \cdot \cdot a_ma_1)\theta$ .

If Q displaces 1 then  $Q\theta$  is uniquely defined; if not, (ii) follows from (3). As to (i), suppose without loss of generality that R does not displace 1; then  $R\theta$  is of the form  $BA_1 \cdot \cdot \cdot A_n B$ , so by successive applications of (4),  $(QR)\theta = (Q\theta)(R\theta) = (R\theta)(Q\theta) = (RQ)\theta$ .

For arbitrary elements Q, R of  $S_6$ ,  $(QR)\theta = (Q\theta)(R\theta)$ . To prove this it is sufficient to consider the case where R is a transposition (since every element of  $S_6$  is a product of transpositions). If Q and R are disjoint the asserted relation is trivial. Hence we write Q as a product of disjoint cycles and let Q' denote the product of those factors of Q which are not disjoint from R. We need to show that  $(Q'R)\theta = (Q'\theta)(R\theta)$ .

Let 1, e, f,  $a_1$ ,  $\cdots$ ,  $a_m$ ,  $b_1$ ,  $\cdots$ ,  $b_n$  denote distinct elements of M.

- (i) If  $Q' = (1 \ a_1 \cdot \cdot \cdot \cdot a_m)$ ,  $R = (1 \ b_1)$ , then  $(Q'\theta)(R\theta) = A_1 \cdot \cdot \cdot A_m B_1 = (1 \ a_1 \cdot \cdot \cdot \cdot a_m \ b_1)\theta = (Q'R)\theta$ .
- (ii) If  $Q' = (e \ a_1 \cdot \cdots \cdot a_m)$ ,  $V = (e \ b_1 \cdot \cdots \cdot b_n)$ , then  $(Q'\theta)(V\theta) = (EA_1 \cdot \cdots \cdot A_m E) \cdot (EB_1 \cdot \cdots \cdot B_n E) = EA_1 \cdot \cdots \cdot A_m B_1 \cdot \cdots \cdot B_n E = (e \ a_1 \cdot \cdots \cdot a_m \ b_1 \cdot \cdots \cdot b_n)\theta = (Q'V)\theta$ .
- (iii) If  $Q' = (1 \ a_1 \cdot \cdots \cdot a_m \ e \ b_1 \cdot \cdots \cdot b_n)$ ,  $R = (1 \ e)$ , with  $m, n \ge 0$ , then  $(Q'\theta)(R\theta) = A_1 \cdot \cdots \cdot A_m(EB_1 \cdot \cdots \cdot B_n E) = A_1 \cdot \cdots \cdot A_mB_nEB_1 \cdot \cdots \cdot B_n = [(1 \ a_1 \cdot \cdots \cdot a_m) \cdot (e \ b_1 \cdot \cdots \cdot b_n)]\theta = (Q'R)\theta$ .
- (iv) If  $Q' = (1 \ a_1 \cdot \cdot \cdot a_m)(e \ b_1 \cdot \cdot \cdot b_n)$ ,  $R = (1 \ e)$ , then  $(Q'\theta)(R\theta) = A_1 \cdot \cdot \cdot A_m E B_1 \cdot \cdot \cdot \cdot B_n E E = A_1 \cdot \cdot \cdot \cdot A_m E B_1 \cdot \cdot \cdot \cdot B_n = (1 \ a_1 \cdot \cdot \cdot \cdot a_m \ e \ b_1 \cdot \cdot \cdot \cdot b_n)\theta = (Q'R)\theta$ .
- (v) If  $Q' = (e \ a_1 \cdot \cdots \cdot a_m f \ b_1 \cdot \cdots \cdot b_n)$ ,  $R = (e \ f)$ , with  $m, n \ge 0$ , then by (4),  $(Q'\theta)(R\theta) = (EA_1 \cdot \cdots \cdot A_m FB_1 \cdot \cdots \cdot B_n E)(EFE) = (EA_1 \cdot \cdots \cdot A_m)(FB_1 \cdot \cdots \cdot B_n FE)$ =  $(EA_1 \cdot \cdots \cdot A_m)(EFB_1 \cdot \cdots \cdot B_n F) = [(e \ a_1 \cdot \cdots \cdot a_m)(f \ b_1 \cdot \cdots \cdot b_n)]\theta = (Q'R)\theta$ .
- (vi) If  $Q' = (e \ a_1 \cdots a_m)(f \ b_1 \cdots b_n) = Q_1' Q_2'$ ,  $R = (e \ f)$ , then, by (ii),  $(Q'\theta)(R\theta) = (Q_1'\theta)(Q_2'\theta)(R\theta) = (Q_1'Q_2'R)\theta = (Q'R)\theta$ .

 $\theta$  is an automorphism of  $S_6$ . Indeed, the kernel, K, of  $\theta$  is a normal subgroup of  $S_6$ , so K is one of  $S_6$ ,  $A_6$ ,  $\{I\}$ , where  $A_6$  denotes the alternating group of degree 6. But  $[(3\ 6)(4\ 5)]\theta = (3\ 6)(4\ 5)$ , so  $K \neq S_6$ ,  $K \neq A_6$ . Therefore  $K = \{I\}$  so  $\theta$  is 1-1 and hence an automorphism.

Finally,  $\theta$  is outer since  $(1\ 3\ 5)\theta = (1\ 2\ 6)(3\ 5\ 4)$ , whereas if  $\theta$  were inner it would map every conjugate class of  $S_6$  onto itself. This completes the proof.

We observe in conclusion that *all* outer automorphisms of  $S_6$  are obtainable with the aid of the above construction. Indeed, as shown by Hölder [1], the automorphism group of  $S_6$  has order 1440=2(6!); thus the group,  $\mathfrak{I}$ , of inner automorphisms is of index 2 in the full automorphism group. Hence if  $\theta$  is any outer automorphism of  $S_6$  then the right coset  $\mathfrak{I}\theta$  includes all outer automorphisms of  $S_6$ .

## References

- 1. O. Hölder, Bildung zusammengesetzter Gruppen, Math. Ann., vol. 46, 1895, pp. 321-422.
- 2. W. Burnside, Theory of Groups of Finite Order, Cambridge, 1911.