

2011

## Stress relief to augment fertility: the pressure mounts

William H. Catherino

*Uniformed Services University of the Health Sciences*, [catheriw@mail.nih.gov](mailto:catheriw@mail.nih.gov)

Follow this and additional works at: <http://digitalcommons.unl.edu/usuhs>

 Part of the [Medicine and Health Sciences Commons](#)

---

Catherino, William H., "Stress relief to augment fertility: the pressure mounts" (2011). *Uniformed Services University of the Health Sciences*. 45.

<http://digitalcommons.unl.edu/usuhs/45>

This Article is brought to you for free and open access by the U.S. Department of Defense at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Uniformed Services University of the Health Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# Stress relief to augment fertility: the pressure mounts

William H. Catherino, M.D., Ph.D.

Program in Reproductive and Adult Endocrinology, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland; and Department of Obstetrics and Gynecology, Uniformed Services University of the Health Sciences, Bethesda, Maryland

Current studies have not conclusively demonstrated an objective and consistent marker of an aberrant stress response; an effect of such a stress response on reproductive outcome; or a benefit of counseling on reproductive outcome in such patients. (*Fertil Steril*® 2011;95:2462–3. ©2011 by American Society for Reproductive Medicine.)

**Key Words:** Infertility, stress response, alpha-amylase, psychological testing, psychological counseling

Dr. Buck Louis and colleagues provide intriguing data supporting the use of salivary  $\alpha$ -amylase as a marker for stress. This marker correlated with day-specific probability of conception, although not with fecundability (1). Although the function of salivary  $\alpha$ -amylase is to digest dietary starch, there is evidence that  $\alpha$ -amylase rapidly increases in the setting of stressful stimuli (2–4). To place the present article in context, several critical questions must be addressed.

## WHAT IS STRESS?

Stress is a physical, mental, or emotional response to events that causes bodily or mental tension. Extreme stress (e.g., anorexia nervosa) certainly can disrupt reproductive function, but the impact of less dramatic stress in humans is unclear. In addition, determining what is stressful is complex, in that individual responses to stressful stimuli can differ dramatically. The relevant issue is stress response, rather than the stress itself. Were all stressful stimuli deleterious to fertility, it is likely our species (as well as all other species) would have ceased to exist long ago. Assuming that an excessive or aberrant stress response is indeed deleterious and a method existed to remove this deleterious response, then a technique to accurately identify patients with such an aberrant response would be extremely helpful.

## HOW IS AN ABERRANT STRESS RESPONSE CURRENTLY MEASURED?

Personality tests are the most common method to identify traits that correlate to aberrant stress response with poor reproductive outcome (5–21). However, despite the wide array of testing, no consistent personality characteristic has been identified. Physiologic responses are thought to be less subjective than personality tests and have also been studied as stress response markers, with similar mixed results (6, 11, 16). Finally, various biochemical markers of acute (1, 11, 15, 17) and chronic (1, 22, 23) stress have been studied. Unfortunately, the correlation between these markers and poor reproductive outcome have been subtle or nonexistent. Either the current methods do not adequately measure aberrant stress response, or the impact of aberrant stress response on fertility is truly subtle or nonexistent.

Received May 24, 2011; accepted May 24, 2011.

W.H.C. has nothing to disclose.

Reprint requests: William H. Catherino, M.D., Ph.D., Uniformed Services University of the Health Sciences, Department of Obstetrics and Gynecology, Building A, Room 3078, 4301 Jones Bridge Road, Bethesda, MD 20814-4799 (E-mail: [catheriw@mail.nih.gov](mailto:catheriw@mail.nih.gov)).

In addition to identifying the ideal objective measure of the stress response, the timing of such a measure can confound findings. Clinicians exert significant effort to predict the outcome of a future infertility intervention, providing prognostic information that can correlate with both stress and outcome. The finding of elevated stress at the beginning of an infertility intervention may represent recognized expectation of poor outcome, and elevated stress toward the end of an IVF cycle may reflect a recognized poor progress, whereas elevated stress after the cycle may reflect disappointment of a failed cycle. To avoid these biases, stress analysis must be done before the collection of any prognostic data, and data must be age-matched.

## DOES STRESS NEGATIVELY IMPACT FERTILITY?

The studies on the association between an aberrant stress response and poor fertility outcomes are mixed, with studies supporting an association (1, 6, 8, 11, 12, 15, 17, 22, 24–26), studies refuting an association (13, 18, 27–29), and even a study suggesting a positive impact of stress on reproduction (14). In addition, there are studies demonstrating a negative impact of stress on male reproduction (19, 30, 31). It remains unclear whether the mixed literature represents difficulty in measurement, difficulty in determining the appropriate fertility endpoint, statistical aberrations resulting in one outcome or the other, or a subtle impact requiring larger sample sizes.

There is also ample anecdotal and documented evidence demonstrating that the diagnosis of infertility and the process of IVF are stressful events (10, 11, 20, 21, 32, 33). Few life experiences rival infertility assessment in its level of personal invasiveness, which certainly increases stress. Removing these confounders is difficult, which can make assessment of the independent influence of an aberrant stress response on fertility outcomes challenging. This is particularly valid given the wide variation in individual responses to the stress of infertility or IVF.

## DOES INTERVENTION REDUCE STRESS?

Assuming that an aberrant stress response is considered a negative influence on fertility intervention, is there a way to intervene such that the stress response itself can be mitigated? Fortunately, this issue does not seem to be controversial. There is evidence that counseling does decrease the stress response in an infertility population (16). Regardless of the impact on fertility, decreasing the stressful experience is of value. The existence of counseling services, regardless of whether the couple participates, is reassuring to patients (34, 35).

## DOES REDUCED STRESS IMPROVE OUTCOME?

Assuming that an aberrant stress response has a negative impact on fertility, and assuming that this aberrancy can be corrected with counseling, does counseling improve fertility rates? Unfortunately the data are limited and not encouraging; counseling does not seem to improve fertility outcomes (36), except perhaps in the extreme case involving induced hypogonadotropic hypogonadism (37). There is certainly opportunity to improve on the counseling intervention and study design to document benefit, but until there is clear and documented benefit, the emotional cost of assuming a negative impact of stress on reproductive outcomes is troublesome.

## IMPLICATIONS OF A "STRESS" DIAGNOSIS

Response to stressful stimuli is highly individualized, and assigning such a response as a causal factor for poor reproductive performance is controversial. Theoretically, an aberrant response can be "unlearned," providing a patient-controlled method for improving fertility. However, should fertility intervention fail, it is possible for both patient and clinical staff to infer that the blame for such failure was an inability of the patient to control their stress response. Such an interpretation adds additional guilt and frustration. Until it

is possible to convincingly demonstrate a negative impact of an aberrant stress response on fertility, until we can adequately measure this stress response, and until we can provide an effective intervention that improves reproductive outcome, it is prudent to avoid adding to the challenges our patients already undergo.

## CONCLUSIONS

Buck Louis and colleagues address one of the challenges in studying the question of aberrant stress response on fertility, that of an easily measurable marker. However, the lack of correlation with fecundability remains a concern. Given the controversy that exists in the literature, it is unlikely that any single study will address the optimal stress response marker, confirmation of an aberrant stress response on poor fertility outcomes, and confirmation that a specific intervention corrects the problem. Proponents who believe that a deleterious stress response does harm reproduction can argue that negative studies used the incorrect tool to identify the aberrant stress response or that the studies were underpowered, whereas opponents can argue that the positive findings to date are spurious and inconsistent. Ultimately, relevance will be based on confirmation of a causal relationship and successful intervention.

## REFERENCES

1. Buck Louis GM, Lum KJ, Sundaram R, Chen Z, Kim S, Lynch CD, et al. Stress reduces conception probabilities across the fertile window: evidence in support of relaxation. *Fertil Steril* 2011;95:2184–9.
2. Bosch JA, Brand HS, Ligtenberg TJ, Bermond B, Hoogstraten J, Nieuw Amerongen AV. Psychological stress as a determinant of protein levels and salivary-induced aggregation of *Streptococcus gordonii* in human whole saliva. *Psychosom Med* 1996;58:374–82.
3. Skosnik PD, Chatterton RT, Swisher T, Park S. Modulation of attentional inhibition by norepinephrine and cortisol after psychological stress. *Int J Psychophysiol* 2000;36:59–68.
4. Takai N, Yamaguchi M, Aragaki T, Eto K, Uchihashi K, Nishikawa Y. Effect of psychological stress on the salivary cortisol and amylase levels in healthy young adults. *Archives Oral Biol* 2004;49:963–8.
5. Kemeter P. Studies on psychosomatic implications of infertility—effects of emotional stress on fertilization and implantation in in-vitro fertilization. *Hum Reprod* 1988;3:341–52.
6. Facchinetti F, Matteo ML, Artini GP, Volpe A, Genazzani AR. An increased vulnerability to stress is associated with a poor outcome of in vitro fertilization-embryo transfer treatment. *Fertil Steril* 1997;67:309–14.
7. Lindheim SR, Sauer MV. Stress and infertility: "the chicken or the egg?". *Fertil Steril* 1997;68:384–5.
8. Sanders KA, Bruce NW. Psychosocial stress and treatment outcome following assisted reproductive technology. *Hum Reprod* 1999;14:1656–62.
9. Csemiczky G, Landgren BM, Collins A. The influence of stress and state anxiety on the outcome of IVF-treatment: psychological and endocrinological assessment of Swedish women entering IVF-treatment. *Acta Obstet Gynecol Scand* 2000;79:113–8.
10. Verhaak CM, Smeenk JM, Eugster A, van Minnen A, Kremer JA, Kraaijaat FW. Stress and marital satisfaction among women before and after their first cycle of in vitro fertilization and intracytoplasmic sperm injection. *Fertil Steril* 2001;76:525–31.
11. Gallinelli A, Roncaglia R, Matteo ML, Ciaccio I, Volpe A, Facchinetti F. Immunological changes and stress are associated with different implantation rates in patients undergoing in vitro fertilization-embryo transfer. *Fertil Steril* 2001;76:85–91.
12. Klonoff-Cohen H, Chu E, Natarajan L, Sieber W. A prospective study of stress among women undergoing in vitro fertilization or gamete intrafallopian transfer. *Fertil Steril* 2001;76:675–87.
13. Anderheim L, Holter H, Bergh C, Möller A. Does psychological stress affect the outcome of in vitro fertilization? *Hum Reprod* 2005;20:2969–75.
14. Cooper BC, Gerber JR, McGettrick AL, Johnson JV. Perceived infertility-related stress correlates with in vitro fertilization outcome. *Fertil Steril* 2007;88:714–7.
15. Li XH, Ma YG, Geng LH, Qin L, Hu H, Li SW. Baseline psychological stress and ovarian norepinephrine levels negatively affect the outcome of in vitro fertilization. *Gynecol Endocrinol* 2011;27:139–43.
16. Facchinetti F, Tarabusi M, Volpe A. Cognitive-behavioral treatment decreases cardiovascular and neuroendocrine reaction to stress in women waiting for assisted reproduction. *Psychoneuroendocrinology* 2004;29:162–73.
17. Smeenk JM, Verhaak CM, Vingerhoets AJ, Sweep CG, Merkus JM, Willemssen SJ, et al. Stress and outcome success in IVF: the role of self-reports and endocrine variables. *Hum Reprod* 2005;20:991–6.
18. Sheiner E, Sheiner EK, Potashnik G, Carel R, Shoham-Vardi I. The relationship between occupational psychological stress and female fertility. *Occup Med (Lond)* 2003;53:265–9.
19. Clarke RN, Klock SC, Geoghegan A, Travassos DE. Relationship between psychological stress and semen quality among in-vitro fertilization patients. *Hum Reprod* 1999;14:753–8.
20. Greenfeld DA. Does psychological support and counseling reduce the stress experienced by couples involved in assisted reproductive technology? *J Assist Reprod Genet* 1997;14:186–8.
21. Beutel M, Kupfer J, Kirchmeyer P, Kehde S, Köhn FM, Schroeder-Printzen I, et al. Treatment-related stresses and depression in couples undergoing assisted reproductive treatment by IVF or ICSI. *Andrologia* 1999;31:27–35.
22. Demyttenaere K, Nijs P, Evers-Kiebooms G, Koninckx PR. Coping and the ineffectiveness of coping influence the outcome of in vitro fertilization through stress responses. *Psychoneuroendocrinology* 1992;17:655–65.
23. Milad MP, Klock SC, Moses S, Chatterton R. Stress and anxiety do not result in pregnancy wastage. *Hum Reprod* 1998;13:2296–300.
24. Demyttenaere K, Nijs P, Evers-Kiebooms G, Koninckx PR. Personality characteristics, psychoendocrinological stress and outcome of IVF depend upon the etiology of infertility. *Gynecol Endocrinol* 1994;8:233–40.
25. Boivin J, Takefman JE. Stress level across stages of in vitro fertilization in subsequently pregnant and non-pregnant women. *Fertil Steril* 1995;64:802–10.
26. Barzilai-Pesach V, Sheiner EK, Sheiner E, Potashnik G, Shoham-Vardi I. The effect of women's occupational psychological stress on outcome of fertility treatments. *J Occup Environ Med* 2006;48:56–62.
27. Merari D, Feldberg D, Elizur A, Goldman J, Modan B. Psychological and hormonal changes in the course of in vitro fertilization. *J Assist Reprod Genet* 1992;9:161–9.
28. Harlow CR, Fahy UM, Talbot WM, Wardle PG, Hull MG. Stress and stress-related hormones during in-vitro fertilization treatment. *Hum Reprod* 1996;11:274–9.
29. Milad MP, Klock SC, Moses S, Chatterton R. Stress and anxiety do not result in pregnancy wastage. *Hum Reprod* 1998;13:2296–300.
30. Ragni G, Caccamo A. Negative effect of stress of in vitro fertilization program on quality of semen. *Acta Eur Fertil* 1992;23:21–3.
31. Sheiner EK, Sheiner E, Carel R, Potashnik G, Shoham-Vardi I. Potential association between male infertility and occupational psychological stress. *J Occup Environ Med* 2002;44:1093–9.
32. Boivin J, Takefman JE. Impact of the in-vitro fertilization process on emotional, physical and relational variables. *Hum Reprod* 1996;11:903–7.
33. Bringhentti F, Martinelli F, Ardent R, La Sala GB. Psychological adjustment of infertile women entering IVF treatment: differentiating aspects and influencing factors. *Acta Obstet Gynecol Scand* 1997;76:431–7.
34. Mazure CM, Greenfeld DA. Psychological studies of in vitro fertilization/embryo transfer participants. *J In Vitro Fert Embryo Transf* 1989;6:242–56.
35. Freeman EW, Boxer AS, Rickels K, Tureck R, Mastroianni L. Psychological evaluation and support in a program of in vitro fertilization and embryo transfer. *Fertil Steril* 1985;43:48–53.
36. Kupka MS, Dorn C, Richter O, Schmutzler A, van der Ven H, Kulczycki A. Stress relief after infertility treatment—spontaneous conception, adoption and psychological counselling. *Eur J Obstet Gynecol Reprod Biol* 2003;110:190–5.
37. Berga SL, Loucks TL. Use of cognitive behavior therapy for functional hypothalamic amenorrhea. *Ann N Y Acad Sci* 2006;1092:114–29.