# University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Nebraska Bird Review

Nebraska Ornithologists' Union

1-1-1999

## Roseate Spoonbill in Otoe County

Stephen J. Dinsmore Fort Collins, CO

Follow this and additional works at: http://digitalcommons.unl.edu/nebbirdrev

Part of the Ornithology Commons, Poultry or Avian Science Commons, and the Zoology Commons

Dinsmore, Stephen J., "Roseate Spoonbill in Otoe County" (1999). *Nebraska Bird Review.* Paper 46. http://digitalcommons.unl.edu/nebbirdrev/46

This Article is brought to you for free and open access by the Nebraska Ornithologists' Union at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Bird Review by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Vol. 67 No.

#### 80

#### Roseate Spoonbill in Otoe County

Stephen J. Dinsmore 612 1/2 West Magnolia St. Fort Collins, CO 80521

On 5 August 1997, W. Ross Silcock and I were birding the heron roost at th Waubonsie Wildlife Area in Fremont County, lowa. We were hoping to see th Roseate Spoonbill I found there on 4 August (see Dinsmore 1998), and though there might be a chance it would fly into Nebraska upon leaving the roost. At 5,3 a.m., the bird left the roost. It flew directly over us (in lowa) and headed to the north with a Great Egret. We continued to watch the bird as it gained altitude as was joined by a second Great Egret. The birds were now flying northwest. After couple minutes, all three birds turned suddenly to the west, and were soon over Nebraska. They then turned south and appeared to be following the bluffs along th Nebraska side of the Missouri River. At this time, they were barely visible through a 20x spotting scope. We watched them fly south over Nebraska City an eventually disappear to the south. When we last saw them, they were flying wes of the Otoe County Omaha Public Power District (OPPD) power plant stack, clearly over Nebraska. We lost sight of the birds at 6:15 a.m. The Roseate Spoonbill wa very distinctive because of the color and bill shape. It was slightly smaller an shorter-legged than a Great Egret, and flew with more rapid wingbeats. The b was long, laterally compressed, and had a spatulate shape. The head, neck, an body were white. The wings were pink, and in flight the primaries had dusky tip The tail was also pink. The legs were pinkish with darker knees. On the basis of the mostly white head and neck and the dusky wingtips, I aged the bird as an immatur (first-fall plumage). On 15 August 1997, I briefly observed the spoonbill in flight over the wetlands near the OPPD plant in Otoe County.

This sighting represents the fourth record of a Roseate Spoonbill in Nebraska. Other records are on 5 June 1932 in Buffalo County (Bray et al 1986), 2 on 20 August 1966 near Hastings, Adams County (Maunder 1966), and 1 on 16 September 1884 in Lancaster County (Bray et al 1986).

### Literature Cited

Bray, T. E., B. K. Padelford, and W. R. Silcock. 1986. The Birds of Nebraska: A Critically Evaluated List. Published by the Authors. Bellevue, Nebraska.

Dinsmore, S. J. 1998. Roseate Spoonbill in southwestern lowa. <u>Iowa Bird Life</u> 68: 136-137.

Maunder, V. 1966. Roseate Spoonbill. Nebraska Bird Review 34: 77.