

11-2006

Incorporation of Polyethylene Glycol into Self-Assembled Monolayers Enhances Substrate-Mediated Gene Delivery by Nonspecifically-Bound Complexes

Angela K. Pannier

University of Nebraska - Lincoln, apannier2@unl.edu

Lonnie D Shea

l-shea@northwestern.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/biosysengpres>



Part of the [Biological Engineering Commons](#)

Pannier, Angela K. and Shea, Lonnie D, "Incorporation of Polyethylene Glycol into Self-Assembled Monolayers Enhances Substrate-Mediated Gene Delivery by Nonspecifically- Bound Complexes" (2006). *Conference Presentations and White Papers: Biological Systems Engineering*. 46.

<http://digitalcommons.unl.edu/biosysengpres/46>

This Article is brought to you for free and open access by the Biological Systems Engineering at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Conference Presentations and White Papers: Biological Systems Engineering by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Podium presentation
American Institute of Chemical Engineers Annual Meeting
November 13, 2006, San Francisco, CA

Incorporation of Polyethylene Glycol into Self-Assembled Monolayers Enhances Substrate-Mediated Gene Delivery by Nonspecifically-Bound Complexes

Angela K. Pannier

Biological Systems Engineering, University of Nebraska-Lincoln,
231 L.W. Chase Hall, Lincoln, NE 68583-0726

Lonnie D. Shea

Chemical and Biological Engineering, Northwestern University,
2145 Sheridan Rd/ TECH E136, Evanston, IL 60208.

Developing systems capable of controlled and efficient gene transfer is a fundamental goal of biotechnology, with applications including functional genomics, gene therapy, and tissue engineering. Substrate-mediated delivery, also termed solid phase delivery, describes the immobilization of DNA, complexed with nonviral vectors, to a biomaterial or substrate through specific or nonspecific interactions. Cells cultured on the substrate are exposed to elevated DNA concentrations within the local microenvironment, which enhances transfection. We investigated transfection resulting from DNA complexes immobilized to a substrate through specific interactions introduced through complementary functional groups on the vector and surface or through nonspecific interactions. Self-assembled monolayers (SAMs) of alkanethiols on gold were used to provide a controlled surface to investigate transfection following specific and non-specific immobilization. DNA, complexed with polyethylenimine (PEI), was immobilized to SAMs through nonspecific mechanisms or covalently linked to SAMs presenting appropriate functional groups through a fraction of the func-

tional groups available on the PEI present in the complex. Nonspecific immobilization of DNA complexes and subsequent transfection was mediated by the hydrophilicity and ionization of the substrate, while covalent tethering resulted in immobilized quantities similar to nonspecific conditions, but provided no transfection. Subsequent studies incorporated polyethylene glycol (PEG)-terminated alkanethiols into the SAMs to reduce nonspecific complex adsorption. Covalent tethering of complexes to PEG/carboxylic acid monolayers resulted in statistically less complex immobilization and no transfection. Nonspecific immobilization to monolayers containing 40% PEG resulted in statistically less DNA complexes immobilized, but substantially greater transfection. Cell adhesion was not affected at this percentage of PEG. Similarly, the addition of Pluronic block copolymers (of polyethylene oxide and polypropylene oxide) to surfaces also enhanced transfection. We hypothesize that the presence of PEG in the monolayer may better preserve complex conformation upon binding to substrates, thereby enhancing the activity of substrate-mediated delivery of DNA complexes.

