

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Wildlife Damage Management Conferences --  
Proceedings

Wildlife Damage Management, Internet Center for

---

October 2000

## Effectiveness of Anise as a coyote lure

Coren P. Jagnow  
*Penn State University*

Russ Mason  
*Utah State University*

Follow this and additional works at: [http://digitalcommons.unl.edu/icwdm\\_wdmconfproc](http://digitalcommons.unl.edu/icwdm_wdmconfproc)



Part of the [Environmental Sciences Commons](#)

---

Jagnow, Coren P. and Mason, Russ, "Effectiveness of Anise as a coyote lure" (2000). *Wildlife Damage Management Conferences -- Proceedings*. 50.

[http://digitalcommons.unl.edu/icwdm\\_wdmconfproc/50](http://digitalcommons.unl.edu/icwdm_wdmconfproc/50)

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Wildlife Damage Management Conferences -- Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

### **Effectiveness of Anise as a coyote lure**

Coren P. Jagnow, Penn State University, 9 Ferguson Bldg., University Park, PA 16802, USA

Russ Mason, USDA-APHIS-WS-National Wildlife Research Center, UMC 5295, Utah State University, Logan UT 84322, USA

*Abstract:* Odors are frequently used to attract coyotes (*Canis latrans*) to control devices. Coyote olfactory perception is vital to effectively manage depredation of livestock by coyotes because of the importance of volatile cues to this species. Lures are usually chosen to reflect the biological needs of coyotes as well as relying on food-seeking behaviors to elicit attraction. Anise (*Illicium verum*) was chosen as the stimulus because of its attractiveness to dogs and its common use as an ingredient in animal foods. Plants comprise a large part of coyotes' diets during warm weather; therefore, a plant odor was chosen to find an effective warm weather coyote lure. Anise was presented in concentrations of 0.0625, 0.125, and 0.250%. The intermediate concentration was the most preferred followed by the highest concentration. The conclusion is that anise may be useful as an odor lure and additional tests are recommended.

*Key words:* Anise, *Canis latrans*, coyote, coyote lure, *Illicium verum*