

1994

A Novel Protein, CSG2p, Is Required for Ca^{2+} Regulation in *Saccharomyces cerevisiae*

Troy Beeler

Uniformed Services University of the Health Sciences

Kenneth Gable

Uniformed Services University of the Health Sciences

Chun Zhao

Uniformed Services University of the Health Sciences

Teresa Dunn

Uniformed Services University of the Health Sciences

Follow this and additional works at: <http://digitalcommons.unl.edu/usuhs>

 Part of the [Medicine and Health Sciences Commons](#)

Beeler, Troy; Gable, Kenneth; Zhao, Chun; and Dunn, Teresa, "A Novel Protein, CSG2p, Is Required for Ca^{2+} Regulation in *Saccharomyces cerevisiae*" (1994). *Uniformed Services University of the Health Sciences*. 50.

<http://digitalcommons.unl.edu/usuhs/50>

This Article is brought to you for free and open access by the U.S. Department of Defense at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Uniformed Services University of the Health Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

A Novel Protein, CSG2p, Is Required for Ca²⁺ Regulation in *Saccharomyces cerevisiae**

(Received for publication, June 28, 1993, and in revised form, November 29, 1993)

Troy Beeler, Kenneth Gable, Chun Zhao, and Teresa Dunn‡

From the Department of Biochemistry, Uniformed Services University of the Health Sciences, Bethesda, Maryland 20814

Nineteen mutants that lost the ability to grow in 100 mM Ca²⁺ (but remained insensitive to 50 mM Sr²⁺) were identified in a screen of approximately 60,000 mutagenized yeast colonies. Cells carrying mutations in the CSG2 gene grow normally in low Ca²⁺ medium but have decreased growth rates when the Ca²⁺ concentration is above 10 mM. The *csg2* mutant cells accumulate much higher levels of Ca²⁺ in a compartment that is exchangeable with extracellular Ca²⁺ but the nonexchangeable Ca²⁺ pool which predominates in wild-type cells is not influenced. Sr²⁺ influx is not increased in the *csg2* mutant cells. Mg²⁺ decreases the amount of Ca²⁺ in the non-exchangeable pool without influencing the *csg2*-induced exchangeable Ca²⁺ pool. The data indicate that the *csg2* mutation causes a selective increase in Ca²⁺ accumulation into a pool which is distinct from the vacuolar pool. The CSG2 protein consists of 410 amino acids, contains nine putative transmembrane segments, four potential sites for N-linked glycosylation, and a sequence with homology to the EF-hand Ca²⁺-binding site.

It has been difficult to demonstrate that *Saccharomyces cerevisiae* requires Ca²⁺ for vegetative growth using Ca²⁺-depleted rich media (YPD-Ca²⁺) or Ca²⁺-free synthetic media. Nonetheless, results from numerous lines of investigation indicate that Ca²⁺ plays an important regulatory role in *S. cerevisiae* (1, 2). Many proteins known to be regulated (directly or indirectly) by Ca²⁺ are present in *S. cerevisiae* including calmodulin (3), calmodulin-dependent protein kinase (4), calmodulin-regulated protein phosphatase (5), kex2 protease (6), trehalase (7), glycogen phosphorylase (8), phospholipase C (9), vacuolar K⁺ channel (10), and protein kinase C (11). Furthermore, numerous Ca²⁺-sensitive (12–14) and Ca²⁺-dependent (15, 16) mutants have been isolated.

Since *S. cerevisiae* has become an experimental model system for investigating the many eukaryotic processes that are controlled by Ca²⁺, (3, 17) it is of interest to identify the genes and proteins that are important in regulating the cytosolic Ca²⁺ concentration of *S. cerevisiae*. The resting cytosolic Ca²⁺ concentration (100–200 nM (17, 18)) is apparently maintained by the active transport of Ca²⁺ from the cytosol into internal organelles such as the vacuole, and out of the cell through the

plasma membrane (19–21). For Ca²⁺ to serve as a regulatory signal, mechanisms must also exist for Ca²⁺ to flow into the cytosol in response to the appropriate stimuli.

One approach toward identifying the genes (and the proteins) which are important in regulating the cytosolic Ca²⁺ is to isolate mutants that lose the ability to grow in medium containing high Ca²⁺ concentrations, identify those mutants with alterations in the regulation of cellular Ca²⁺, and then clone and characterize the genes which complement the mutations. Anraku and co-workers (13) have generated a collection of Ca²⁺-sensitive mutants that represent 18 complementation groups. Nine of the genes are required for vacuolar morphogenesis or are subunits of the vacuolar proton ATPase (14). One of the genes is allelic to *CDC24* (22).

We have recently generated a collection of Ca²⁺-sensitive mutants using a screening protocol similar to that of Anraku and co-workers (13) except that the mutagenesis and prescreen growth was done in media containing 1 mM EGTA and the primary screen was done at 37 °C to allow identification of temperature-sensitive, Ca²⁺-sensitive mutants. Because defects in the vacuolar transport system are expected to cause Sr²⁺ sensitivity, we focused on the mutants that are specifically sensitive to Ca²⁺ but resistant to Sr²⁺ hoping to identify candidates for novel Ca²⁺ transport genes not related to vacuole function. Here we report on our collection of *csg* (Calcium Sensitive Growth) mutants and the molecular/biochemical analysis of one of the isolates, *csg2*, which has identified a novel protein that may regulate Ca²⁺ accumulation by a non-vacuole organelle.

EXPERIMENTAL PROCEDURES

Materials—Arsenazo III was obtained from Aldrich. ⁴⁵Ca²⁺ was purchased from DuPont-New England Nuclear. Zymolyase 100T was supplied by Seikagaku Kogyo, Rockville, MD. All other chemicals were purchased from Sigma.

Strains, Growth Media, and Transformations—The strains used in this study are DBY947(α *ade2-101 ura3-52*), TDY2003 (α *ade2-101 ura3-52 csg2-1*), TDY2040 (α *ade2-101 trp1Δ leu2Δ ura3-52 csg2::LEU2**), and TDY821 *mata/α ura3-52/ura3-52 lys2/+ +ade2-101 trp1Δ/trp1Δ leu2Δ/leu2Δ can^R/can^S* or were derived from them by standard genetic crosses. Yeast media for growth and sporulation were prepared according to Sherman *et al.* (23). Yeast were transformed by the method of Ito (24) except that 5 μg of sonicated salmon sperm DNA was routinely added to each reaction.

Bacterial strains SCS1, JM109, and JM110 were used for the propagation of plasmids and M13 phage. M13mp18 and mp19 phage and the pRS316 phagemid (25) were used to generate single-stranded template DNA for sequence determination.

Mutagenesis—DBY947 or an isogenic *a*-mating strain was mutagenized by a modification of the method described by Lindgren *et al.* (26). Cells (10⁶/ml) in YPD + 1 mM EGTA were treated with 1.0% EMS¹ for 90 min at 26 °C which induced about 70% killing. The cells were diluted 1000-fold into YPD + 1 mM EGTA and allowed to recover for 4 h. Aliquots (200 μl) were removed and spread on YPD agar plates contain-

* This work was supported by Grant GM46495 from the National Institutes of Health and Grants CO71bK and R071AG from the Uniformed Services University of the Health Sciences. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

The nucleotide sequence(s) reported in this paper has been submitted to the GenBank™/EMBL Data Bank with accession number(s) L24113.

‡ To whom correspondence should be addressed: Dept. of Biochemistry, Uniformed Services University of the Health Sciences, 4301 Jones Bridge Rd., Bethesda, MD 20814. Tel.: 301-295-3592; Fax: 301-295-3512.

¹ The abbreviations used are: EMS, ethylmethanesulfonate; PIPES, 1,4-piperazinebis(ethanesulfonic acid); ORF, open reading frame.

-309		GCCTGCAGA	
-300	GCAGCGGCCAAGCAGAAGGAGGCATGGTACTCCTTCTTATTCAAATAAGATTGTAATATATGCCTACTTCTTCT		
-225	CTTGTATATACATATATACTTATAACCGCTATAACCGTCTTGTAAATATCGGCTATCACCGCGGAAGGTGTATC		
-150	GCAAGAAAAAAAAGCTCTCTCACTCTAAGGAGGCTATGTGAACGTTGGGAAGATCAAGTGAAGAAAATGTTTG		
-75	TAGGCCATTTTCTCCAGAACAGATCCGCTCTTGAGCTGCTGAGTTAGCACGATAACAAACAAGATACAGCGTCC		
+1	ATGTCTACCACACTACTTTGGTTTTCAAGTCTAATAGGCTACGCTGATTCAAACAAAATGTTTGTCTAACATACAA		
	M S T T L L W F S S V I G Y V I Q T K C L S N I Q		+25
+76	TCTAAAAAGGAAATCTCCGTGGGGCCCAATGGTACAATGCAACGGCTGAAACTAACGGCGACAACGGAAACTCA		
	S K K E I S V G P N G T I A T P E T N G D N G N S		+50
+101	AGTTCATTAACCTTCTATCTGACCTTTATGTATTTGCTTCTGGTGGCTCTGGGTGCCTGCAATCGACTTTGG		
	S S L T F Y L T F M Y F A S W L L W V P A S R L W		+75
+176	GAGAAGATGAGACCGATGTTTGTCTGACTCAGACTCGAACAGGAATCTCAGTTTGACAACAACAACACCGGG		
	E K M R P M F V S D S D S N R N S Q F <u>D N N N S G</u>		+100
+251	TCTGTGACAAACGAAGATGTCGATACGTTCTCGCAGGTGTTGGATGATCCTCAACCGGATTCCAGCCCAACAG		
	<u>S V T N E D</u> V D T F S H V L D D P Q P R I P A Q Q		+125
+326	CAGAAGCAAAAATCATATCCGTGGCTACCTTCAAATATGTGGCTAAGCTAACAGTGTGGCTCTCATAATGATT		
	Q K Q K I I S V A T F K Y V A K L T V L A L I M I		+150
+401	GTCGGTGATTGACTTATAACATGGCTTTGTCATGTCACCGGCATTGATGTTGCTTTGATGCAAAAATCTGCC		
	V A D L T Y N M A L S L S P A F D V A L M Q N T A		+175
+476	ATTTTCGAAATGCTCACTTTACTATATGGTGTGCGAATCTCCAGGAAGAACTACGTTTCCGTAATTTCTC		
	I F E I V T L L Y G V C G I S R K N Y V F R N F L		+200
+551	ATCATGATGAACCGGCTATTGGAATTTGATCATCTCATAACAGGAGGCTACCTGTGACATGCTTCCGGAAAG		
	I M M N A V I G I L I I S Y T K A T C D M L A G K		+225
+626	CTGTCGCTCAACCTAAGACGGGTGAACCTTTGACCCATCTGTTGATAGGTTGAAAGTGCTCTGATTGGC		
	L S V N P N T G E L S D P F L F D R L K G A L I C		+250
+701	GGCCTTGGTGCTTTGATTATGGTCCCTTTGGCGTGTATGGAACCGTTGCTTTGCACTAACATTTCCAAGAAC		
	G L G A L I M G P F A V L W N R W F C S N I S K N		+275
+776	GAAAACTCTGCTAGTCTTGGTTAAGCAGAGCACCCAGATGGCCCTAATCGGTATTATTGGCATGTAATACTT		
	E N S A V V L V K Q S T H M A L I G I I G M V I L		+300
+851	TTGCCATTTATCTCAAATTCCTCCCGTGAAGTCTGTTGGAATCCATTTCGTTGTTCTATAATGACAAGAGCTTT		
	L P F I P K F P S R E S V E S I S L F Y N D K S F		+325
+926	TGTTCTCTCTACTAGGCTCGATTATCTTTGGTTCCTTCCGAGCTTGATTCCGATATTAGAGTTGAATCGCAAG		
	W F S L L G S I I F G S L P S L I S I L E L N R K		+350
+1001	GCCCTGCTGAGTATTGACGACGTCGAACCTGGGAGCTATTATCTTTATCGGGTTAGCTCAGTGGGTTTCCGAA		
	A P A E Y L T T C N L G A I I F M G L A E W V C E		+375
+1076	CCTACGCAACCAATTTGAGATGGGAAGTCATAGGATACATAATGCTAACGGTAAGTTTGTGGTCTATCA		
	P T Q T T I V R W E V I G Y I M L T V S L L V L S		+400
+1151	GTAACACTCGGGGAAGTAAATACCACATTAGTACGGACAACATACTTTATATGACTGATGATGTAACAAG		
	V T L G E G K Y H H U		+410
+1226	CATATGCAATTATTGGTTTCCCTAATAAAAAATCCCAATCAATATCTTGATTATTTTCAAGACAATTACTAGG		
+1301	ATGTTCTTCTATTTTTTATTTTATTTGTATATTGTATATTCTAAAAAGAGAAGCCATTTGGTGGCCTTATT		
+1376	ATAAATATTAAGAGGCATACCTCCGCTATCCGCTATCCCTGCTATGAACAATCAATTAAGACGGCTTATAAG		
+1451	CGATATACAAG		

FIG. 2. Nucleotide and translated amino acid sequence of CSG2. The sequence was determined as described under "Experimental Procedures." The sequence of a putative Ca²⁺-binding site is underlined.

these *csg* mutants identifies two major complementation groups and eight independent isolates (Table I).

Among the Ca²⁺-specific mutants, those in the *csg1* and *csg2* complementation groups have an average 315 and 1090% increase in the cellular Ca²⁺ relative to wild-type when incubated for 1.5 h in 10 mM Ca²⁺ at 37 °C. All other Ca²⁺-specific mutant isolates show wild-type levels of Ca²⁺ accumulation except for the mutant strain TDY 2022 which shows a 680% increased Ca²⁺ accumulation.

Because mutants in the *csg2* complementation group show the most dramatic Ca²⁺ over-accumulation phenotype and have an extremely tight Ca²⁺-sensitive growth phenotype, this group was characterized first. A heterozygous *csg2-1/CSG2+* diploid grows normally on YPD + 100 mM Ca²⁺ indicating that the mutation is recessive. Tetrad analysis of the heterozygous diploid shows 2 Ca²⁺-sensitive:2 Ca²⁺-resistant spore colonies for each 4-spored tetrad indicating that the mutant phenotype is due to a single nuclear mutation.

The wild-type *CSG2+* gene was cloned by selecting for its ability to complement the *csg2-1* Ca²⁺-sensitive phenotype as described under "Experimental Procedures." The *CSG2* gene

was localized within the complementing fragment of genomic DNA by subcloning various restriction fragments and testing their ability to complement the *csg2-1* Ca²⁺-sensitive phenotype.

Nucleotide Sequence of CSG2—The sequence from the *EcoRI* site in both directions was determined. The *EcoRI* site was found to lie within a large open reading frame (ORF). The entire nucleotide and deduced amino acid sequence of the predicted 45,579-dalton protein is presented in Fig. 2.

A comparison of the *CSG2* ORF to sequences in the NBRF and EMBL GenBank databases shows that an amino-terminal portion of the *CSG2* ORF is identical to an unidentified ORF at the end of the cloned 1.7-kilobase DNA fragment that also contained the full *SCO1* gene. *SCO1* was mapped to chromosome II (33, 34).

The hydropathy analysis of *CSG2* indicates the existence of up to nine transmembrane sequences (Fig. 3A). A possible topological representation of *CSG2p* is shown in Fig. 3B. A potential signal sequence is found at the amino terminus. It contains a sequence of 12 hydrophobic amino acids followed by two potential signal peptidase cleavage sites (TKC ↓ LS ↓ N) (35, 36).





FIG. 3. Identification of membrane spanning segments and potential glycosylation and Ca²⁺-binding site on CSG2p. Panel A, Kyte and Doolittle hydropathy plot of CSG2p. Each point on the curve represents a hydropathy average of a 14-amino-acid window centered at that point. Proposed transmembrane domains are numbered 1-9. Panel B, schematic representation of CSG2p showing possible signal peptidase cleavage sites, a putative Ca²⁺-binding site, 2 N-linked glycosylation sites, and transmembrane segments.



Four consensus N-linked glycosylation sites are found; however, two of the four are placed on the cytoplasmic side of the membrane in our model suggesting that only two of the sites are available for glycosylation. A 12-amino-acid segment with homology to a consensus Ca²⁺-binding loop of the EF-hand type is found in the first cytoplasmic loop (Fig. 3B). The comparison of this site to the consensus Ca²⁺-binding site as determined by Marsden *et al.* (37, 38) is shown in Fig. 4.

Disruption of the Chromosomal CSG2 Locus—The four independent non-complementing alleles of *csg2* identified in our screen for Ca²⁺-sensitive mutants all show similar biochemical phenotypes. To determine the phenotype of a null mutation at the *CSG2* locus, the cloned *CSG2* gene was used to construct a deletion/substitution at the *CSG2* locus as described under "Experimental Procedures." The disrupted allele, designated *csg2::LEU2+*, was substituted at the chromosomal locus by a

one-step gene replacement (32). Spore colonies having the *csg2::LEU2+* null allele are viable and grow comparably to wild-type on YPD medium but fail to grow on the same medium containing 50 mM Ca²⁺. The null allele is similar to the EMS-derived *csg2* alleles with regard to growth properties and biochemical phenotypes. Hence it is the lack of functional CSG2p that causes Ca²⁺ sensitivity. The Ca²⁺ transport properties of the *csg2* null mutant are the same as those of the EMS-induced *csg2* mutants. Accordingly, all the biochemical data presented below is for the *csg2::LEU2+* null mutant strain.

Effect of Ca²⁺ on the Growth Rate of *csg2* Mutant Cells—The rate of wild-type cell growth is influenced little by 50 mM Ca²⁺. The growth rate of *csg2::LEU2+* cells in the absence of added Ca²⁺ is similar to that of wild-type, but in the presence of 50 mM Ca²⁺, *csg2::LEU2+* cells fail to grow. Ca²⁺ concentrations as low as 10 mM decrease the growth of *csg2::LEU2+* at 26 °C (Fig. 5).

CSG2 Sequence 92-103											
Asp	Asn	Asn	Asn	Ser	Gly	Ser	Val	Thr	Asn	Glu	Asp
Amino acids that occur in the Ca ²⁺ -binding loop of EF-hands with the percentage of total amino acids found											
Position in Ca ²⁺ -Binding Loop											
1	2	3	4	5	6	7	8	9	10	11	12
ASP	Lys	Asp	Gly	Asp	GLY	Phe	Ile	Asp	Phe	GLU	Glu
98	28	73	51	56	89	16	70	33	16	31	86
Asn	Ala	ASN	Lys	SER	Asp	Tyr	VAL	Ser	Val	Asp	ASP
0.6	10	24	14	22	2	15	13	19	12	27	9
Tyr	Thr	Glu	ASN	Asn	Asn	Lys	Leu	Glu	Ala	Ala	Val
0.6	10	2.5	9	17	2	13	9	14	10	10	2
Glu	Gln	Ser	Arg	Gly	Lys	Thr	Met	THR	Glu	Lys	Asn
0.6	9	0.6	6	2	2	13	2.5	11	10	8	1
-	Val	-	Ala	Gln	Arg	Glu	Cys	Gly	Leu	Gln	Gln
-	8	-	5	0.6	1	6	2	10	10	5	0.6

FIG. 4. Comparison of the amino acid sequence of a putative Ca²⁺-binding site and the consensus Ca²⁺-binding site of the EF hand. The amino acids typed in **bold** are the ones found in CSG2p.




FIG. 5. Effect of Ca²⁺ on the growth rate of wild-type and *csg2* mutant cells. **Panel A**, wild-type (○, ●) and *csg2::LEU2+* (▽, ▼) cells were grown in YPD buffered to pH 4.7 with citrate at 26 °C to an OD_{600 nm} of 0.2. The cells were then split to YPD, pH 4.7, with (●, ▼) or without (○, ▽) 50 mM CaCl₂. At the indicated time aliquots were removed, diluted 1:1 in 0.1 M K-EDTA, pH 7.6, and the OD_{600 nm} was determined. Dilutions were made as necessary to maintain the OD_{600 nm} between 0.1 and 0.3 throughout the experiment. **Panel B**, wild-type (○) and *csg2::LEU2+* (▽) cells were grown to an OD_{600 nm} of 0.1 in YPD, pH 4.7, at 26 °C, split to six tubes, and 0, 1, 3, 10, 30, or 100 mM CaCl₂ was added. Aliquots were removed at 0 and 8 h, diluted, and spread on YPD plates to determine the number of viable cells. The number of cells at 8 h is plotted as a percent of the original cell number at *t* = 0.

Effect of *csg2* Mutant on Cellular Ca²⁺ Levels—The rate as well as the amount of Ca²⁺ accumulated by *csg2::LEU2+* cells in 10 mM Ca²⁺ is greater than that of wild-type. Sr²⁺ accumulation by *csg2::LEU2+* cells was increased 2.5 times while Ca²⁺ accumulation increased 22 times indicating that Ca²⁺ accumulation was more influenced by the *csg2* mutation (Fig. 6).

In wild-type cells, most cellular Ca²⁺ is localized in the vacuole where it forms a stable, nonexchangeable complex with polyphosphate (18). To determine if the increased Ca²⁺ accumulated by *csg2::LEU2+* cells is also sequestered in a nonexchangeable Ca²⁺ pool, cells were incubated for 3 h in YPD medium containing 30 mM Ca²⁺ with 1 μCi/ml ⁴⁵Ca²⁺ and then shifted into YPD without ⁴⁵Ca²⁺ (Fig. 7).

For the *csg2::LEU2+* mutant the amount of nonexchangeable Ca²⁺ (~400 nmol Ca²⁺/mg) is similar to that in wild-type, but the large increase in an exchangeable Ca²⁺ pool (~800 nmol




FIG. 6. Ca²⁺ and Sr²⁺ accumulation by wild-type and *csg2* mutant cells. Wild-type (▽, ▽) and *csg2* mutant cells (●, ○) were in exponential growth at 27 °C. At zero time, 10 mM Ca²⁺ (▼, ●) or Sr²⁺ (▽, ○) was added, and at various times aliquots were removed to determine viability and the amount of Ca²⁺ or Sr²⁺ accumulated. To determine the Ca²⁺ or Sr²⁺ loading level, cells were washed by centrifugation and resuspended to a density of 10⁷/ml in KCl solution containing 50 μM arsenazo III. Accumulated Ca²⁺ or Sr²⁺ was released by digitonin (1 mg/ml) addition and measured spectrophotometrically by monitoring the difference absorbance of the arsenazo III-Ca²⁺ or arsenazo III-Sr²⁺ complex at 660–685 nm using an AMINCO-SLM dualbeam spectrophotometer. Viability (*inset*) was determined by diluting aliquots 10⁴-fold and spreading 200 μl onto YPD plates.




FIG. 7. The *csg2*-induced Ca²⁺ pool is exchangeable. Wild-type (○) and *csg2::LEU2+* (●) cells were grown to an OD_{600 nm} of 0.5 in YPD and diluted into YPD + 30 mM Ca²⁺ (1 μCi/ml) to a OD_{600 nm} of 0.1. After a 3-h incubation at 26 °C, the cells were washed three times by centrifugation with cold YPD and resuspended in YPD without ⁴⁵Ca²⁺. At the indicated times, the ⁴⁵Ca²⁺ remaining in the cells was determined by filtering 5-ml aliquots, washing the filters with 3 × 5 ml of cold 20 mM Mg²⁺, drying the filters, and then measuring the ⁴⁵Ca²⁺ on the filter by scintillation counting. At 5, 6, and 18 h after ⁴⁵Ca²⁺ release was initiated, the wild-type and *csg2::LEU2+* cells had essentially the same amount of ⁴⁵Ca²⁺.

Ca²⁺/mg) suggests Ca²⁺ accumulation by a nonvacuolar organelle.

Another distinction between the *csg2*-induced Ca²⁺ pool and the vacuolar Ca²⁺ pool is the response of Ca²⁺ loading to increasing the Mg²⁺ concentration of the growth medium. Ca²⁺ accumulation by wild-type cells is blocked by extracellular Mg²⁺ (Fig. 8). Similarly the nonexchangeable Ca²⁺ pool in the *csg2* mutant is eliminated by Mg²⁺, but the exchangeable Ca²⁺ pool (Fig. 8, *inset*) is not affected by Mg²⁺, further indicating that the *csg2* mutant has normal vacuolar Ca²⁺ accumulation which is blocked by Mg²⁺, but increased accumulation into a non-vacuolar organelle which is not influenced by Mg²⁺. Normal vacuolar Ca²⁺ transport in *csg2::LEU2+* cells is further indicated by the observation that the rate and amount of vacuolar Ca²⁺ accumulation by permeabilized cells and isolated vacuole membrane vesicles is the same for *csg2::LEU2+* mutant cells as for wild-type (data not shown).

DISCUSSION

The data indicate that CSG2p is involved in the regulation of Ca²⁺ accumulation by an organelle which sequesters Ca²⁺ in an exchangeable form. The cytosolic Ca²⁺ concentration in both




FIG. 8. The effect of Mg^{2+} on the cellular Ca^{2+} content of wild-type and *csg2* mutants. Wild-type (\circ) and *csg2::LEU2+* mutant (\bullet) cells were grown to an OD_{600} of 1.2 and diluted 10-fold into YPD medium containing 50 mM CaCl_2 and the indicated Mg^{2+} concentration (YPD medium has about 0.6 mM Mg^{2+}). After a 2-h incubation, the amount of Ca^{2+} accumulated was measured as described in Fig. 6. The inset shows the A23187-stimulated Ca^{2+} -release rate from *csg2::LEU2+* cells that were loaded in YPD (top trace) or YPD + 5 mM Mg^{2+} (bottom trace).

the wild-type and the *csg2::LEU2+* cells is apparently similar since vacuolar Ca^{2+} accumulation (polyphosphate dependent, Mg^{2+} inhibitable, and nonexchangeable) is not significantly influenced, and the cytosolic Ca^{2+} concentration as measured with fura 2 (data not shown) is not changed. The increased loading of *csg2::LEU2+* cells must, therefore, arise from altered Ca^{2+} fluxes across the membrane of the affected organelle.

The Ca^{2+} -induced killing and over-accumulation of Ca^{2+} by *csg2::LEU2+* cells is caused by the lack of CSG2 protein as opposed to an alteration in the properties of the protein. One model to explain the effect of the *csg2* mutation on cellular Ca^{2+} is that eliminating CSG2p directly causes an organelle to over-accumulate Ca^{2+} . If CSG2p directly mediates Ca^{2+} translocation across the organelle membrane, it must function in efflux. Alternatively, it might act to negatively regulate a Ca^{2+} pump. Because CSG2p is a membrane protein with a putative Ca^{2+} -binding site on the cytoplasmic side of the membrane, it is reasonable to suppose that the action of the protein may be regulated by cytosolic Ca^{2+} . The regulation of Ca^{2+} release from endoplasmic and sarcoplasmic reticulum by cytosolic Ca^{2+} (Ca^{2+} -induced Ca^{2+} release) has been demonstrated in other organisms. CSG2p is not homologous to the IP_3 -activated Ca^{2+} channel or the ryanodine-activated Ca^{2+} channel found in muscle cells; however, it might be a member of a new family of intracellular Ca^{2+} channel proteins. Although in wild-type cells, the amount of Ca^{2+} in the CSG2p-regulated Ca^{2+} pool is much less than that in the vacuole, it may still be important in regulating cytosolic Ca^{2+} levels in response to physiological signals since mobilization of only 1% of the cellular Ca^{2+} would raise the cytosolic Ca^{2+} concentration to 10 μM .

There are two alternative models which could explain the Ca^{2+} sensitivity of *csg2* mutants. 1) CSG2p is important in regulating a Ca^{2+} -dependent process, so that in the absence of functional CSG2p and in the presence of high Ca^{2+} concentra-

tions, this Ca^{2+} -regulated process is inappropriately activated leading to cell death. 2) CSG2p mediates an important cellular process, for which another system can substitute in low Ca^{2+} concentrations; however, the alternative system is blocked by Ca^{2+} making CSG2p essential at high Ca^{2+} concentrations. Both models assume that the Ca^{2+} -overloading phenotype is a consequence of Ca^{2+} -induced cell damage. Although our data do not eliminate these two alternative models, we favor the model described above in which the *csg2* mutant directly causes an organelle to overaccumulate Ca^{2+} since specific Ca^{2+} accumulation by yeast cells is not generally associated with cell damage or death.

REFERENCES

- Davis, T. N., and Thorner, J. (1986) in *Yeast Cell Biology* (Hicks, J., ed) pp. 477-503, Alan R. Liss, New York
- Iida, H., Sakaguchi, S., Yagawa, Y., and Anraku, Y. (1990) *J. Biol. Chem.* **265**, 21216-21222
- Davis, T., Urdea, M. S., Masiarz, F. R., and Thorner, J. (1986) *Cell* **47**, 423-431
- Ohya, Y., Kawasaki, H., Suzuki, K., Londesborough, J., and Anraku, Y. (1991) *J. Biol. Chem.* **266**, 12784-12794
- Cyert, M., Kunisawa, R., Kaim, D., and Thorner, J. (1991) *Proc. Natl. Acad. Sci. U. S. A.* **88**, 7376-7380
- Mizuno, K., Nakamura, T., Ohshima, T., Tanaka, S., and Matsuo, H. (1989) *Biochem. Biophys. Res. Commun.* **159**, 305-311
- Neves, M.-J., and Francois, J. (1992) *Biochem. J.* **288**, 859-864
- Francois, J., and Hers, H.-G. (1988) *Eur. J. Biochem.* **174**, 561-567
- Yoko-o, T., Matsui, Y., Yagisawa, H., Nojima, H., Uno, I., and Toh-e, A. (1993) *Proc. Natl. Acad. Sci. U. S. A.* **90**, 1804-1808
- Berl, A., and Slayman, C. L. (1990) *Proc. Natl. Acad. Sci. U. S. A.* **87**, 7824-7828
- Levin, D. E., Fields, F. O., Kunisawa, R., Bishop, J. M., and Thorner, J. (1990) *Cell* **62**, 213-224
- Ohya, Y., Miyamoto, H., Suzuki, K., Londesborough, J., and Anraku, Y. (1991) *J. Bacteriol.* **165**, 28-33
- Ohya, Y., Ohsumi, Y., and Anraku, Y. (1986) *J. Gen. Microbiol.* **132**, 979-988
- Ohya, Y., Umemoto, N., Tanida, I., Ohta, A., Iida, H., and Anraku, Y. (1991) *J. Biol. Chem.* **266**, 13971-13977
- Ohya, Y., Ohsumi, Y., and Anraku, Y. (1984) *Mol. Gen. Genet.* **193**, 389-394
- Schmitt, H. D., Pazicha, M., and Gallwitz, D. (1988) *Cell* **53**, 635-647
- Iida, H., Yagawa, Y., and Anraku, Y. (1990) *J. Biol. Chem.* **265**, 13391-13399
- Dunn, T., Gable, K., and Beeler, T. (1994) *J. Biol. Chem.* **269**, 7273-7278
- Eilam, Y. (1982) *Microbios* **35**, 99-110
- Ohsumi, Y., and Anraku, Y. (1983) *J. Biol. Chem.* **258**, 5614-5617
- Rudolph, H. K., Antebi, A., Fink, G. R., Buckley, C. M., Dorman, T. E., LeVitre, J., Davidow, L. S., Mao, J., and Moir, D. T. (1989) *Cell* **58**, 133-145
- Miyamoto, S., Ohya, Y., Ohsumi, Y., and Anraku, Y. (1987) *Gene (Amst.)* **54**, 125-132
- Sherman, F., Fink, G., and Lawrence, C. (1974) *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY
- Ito, H., Fukuda, Y., Murata, K., and Kimura, A. (1983) *J. Bacteriol.* **153**, 163-168
- Sikorski, R. S., and Hieter, P. (1989) *Genetics* **122**, 19-27
- Lindgren, G., Hwang, Y. L., Oshima, Y., and Lindgren, C. C. (1965) *Can. J. Gen. Cyto.* **7**, 491-499
- Holm, C., Meeks-Wagner, D., Fangman, W., and Botstein, D. (1986) *Gene (Amst.)* **42**, 169-173
- Holmes, D. S., and Quigley, M. (1981) *Anal. Biochem.* **114**, 193-197
- Rose, M. D., Novick, P., Thomas, J. H., and Fink, G. R. (1987) *Gene (Amst.)* **60**, 237-243
- Boeke, J. D., LaCrute, F., and Fink, G. R. (1984) *Mol. Gen. Genet.* **197**, 345-346
- Southern, E. (1975) *J. Mol. Biol.* **98**, 503-517
- Rothstein, R. J. (1983) *Methods Enzymol.* **101**, 202-211
- Schulze, X., and Rödel, X. (1988) *Mol. Gen. Genet.* **211**, 492-498
- Schulze, X., and Rödel, X. (1989) *Mol. Gen. Genet.* **216**, 37-43
- von Heijne, G. (1983) *Eur. J. Biochem.* **116**, 17-21
- von Heijne, G. (1988) *Nucleic Acids. Res.* **14**, 4683-4690
- da Silva, A. C. R., and Reinach, F. C. (1991) *Trends Biochem. Sci.* **16**, 53-57
- Marsden, B., Shaw, G. S., and Sykes, B. D. (1990) *Biochem. Cell Biol.* **68**, 587-601
- Kyte, J., and Doolittle, R. F. (1982) *J. Mol. Biol.* **157**, 105-132
- Difco Manual* (1953) Ninth Edition, Difco Laboratories, Detroit, MI