

February 1971

Peptidoglycan Differences in Strains of *Bacillus cereus* Constitutive and Inducible for Penicillinase Production

Kenneth Nickerson
UNL, knickerson1@unl.edu

Richard A. Day
University of Cincinnati, Cincinnati, Ohio

Follow this and additional works at: <http://digitalcommons.unl.edu/bioscimicro>

 Part of the [Microbiology Commons](#)

Nickerson, Kenneth and Day, Richard A., "Peptidoglycan Differences in Strains of *Bacillus cereus* Constitutive and Inducible for Penicillinase Production" (1971). *Papers in Microbiology*. 53.
<http://digitalcommons.unl.edu/bioscimicro/53>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Microbiology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Peptidoglycan Differences in Strains of *Bacillus cereus* Constitutive and Inducible for Penicillinase Production

KENNETH W. NICKERSON¹ AND RICHARD A. DAY

Department of Chemistry, University of Cincinnati, Cincinnati, Ohio 45221

Received for publication 6 March 1970

Strains of *Bacillus cereus* differing in penicillinase production were shown to possess cell walls of differing muramic acid contents.

Although it is generally agreed (1, 9, 11) that penicillinase (EC 3.5.2.6) is somehow related to a cell wall biosynthetic enzyme, the exact nature of this relation is still obscure. There appear to be two main viewpoints. Pollock (9) envisions an ancient gene duplication, followed by separate evolution, in response to the reputed existence of penicillin as a selective force in nature. In this case, the two enzymes would contain only partially the same amino acid sequence. Alternatively, penicillinase may be a direct, limited step mutation from that biosynthetic enzyme (1). If so, the question arises whether the original biosynthetic capacity has been retained, lost, or altered. The two cell wall biosynthetic enzymes most commonly suggested to be related to penicillinase are D-alanine carboxypeptidase and transpeptidase (9, 11).

We studied cell walls from *Bacillus cereus* NRRL 569 and *B. cereus* 569/H (NCTC 9945), inducible and constitutive, respectively, for penicillinase production. These strains differ only by a point mutation (6). Nonconstitutive revertants can be isolated from strain 569/H which again possess basal levels of penicillinase identical to 569. Observed differences in cell wall structure are interpreted in terms of an altered biosynthetic capacity. Ampicillin-resistant strains of *Escherichia coli* were previously shown (2) to possess "weakened" cell walls.

The amino sugar content of acetone- and ether-washed whole cells is a quantitative measure of cell wall content (7). Two methods were used to make this determination. The average of three Elson-Morgan determinations (7) showed 569/H to possess 40% more amino sugars of all types than did 569. Application of hydrolyzed samples to an amino acid analyzer showed 569/H to possess 37% more glucosamine.

Hydrolysates of mechanically purified cell walls yielded the results shown in Table 1. The purity of the samples is apparent in the absence of any peaks other than those indicated. The walls appear to contain a typical diaminopimelic acid peptidoglycan with the carboxyls of both glutamic acid and diaminopimelic acid present as amides. There is at least one teichoic acid or other glucosamine-galactosamine containing polymer and no attached whole protein.

The most interesting difference between these two strains is the lowered muramic acid value of 569/H. We interpret this to indicate the presence of an enhanced number of extended-chain, peptide cross-links (5, 8, 10), in which the L-alanine at the start of one tetrapeptide sequence is no longer attached to muramic acid but is instead attached to the D-alanine at the end of another tetrapeptide. The observed muramic acid difference occurs without a concurrent drop in the amino acids which comprise the peptide cross-links. The lack of such a 1:1 relationship has been observed (3, 4) in many other bacteria and may be a common consequence of extended peptide chain formation.

Significantly, no extended-chain, peptide cross-links have been observed (4) to contain two adjacent D-alanines, indicating that one has been lost in the chain extension process. A transpeptidase has been hypothesized (10) to be the agent of such lengthening. It is not yet clear whether this transpeptidase is identical to the transpeptidase known (11) to form the D-alanyl-diaminopimelic acid cross-links, but the decreased muramic acid content of strain 569/H appears to be due to the enhanced production or activity of a chain-extending transpeptidase. We cannot yet say for certain whether this enhanced activity is due to derepression of the enzymes responsible concomitantly with that of penicillinase in strain 569/H, or whether penicillinase itself still possesses such a biosynthetic function. For reasons presented in

¹ Present address: Department of Genetics, University of Wisconsin, Madison, Wis. 53706.

TABLE 1. *Composition of mechanically purified Bacillus cereus cell walls*^a

Component	Strain ^b	
	569	569/H
Diaminopimelic acid	29	28
Glutamic acid	27	25
Muramic acid	21	10
Ammonia	54	44
Glycine	3	1
Alanine	47	46
Glucosamine	82	84
Galactosamine	24	26

^a Eighteen-hour cultures were harvested and washed with water. Cells were broken in a 9-kc sonic oscillator. The cell walls were collected by centrifugation and washed repeatedly with phosphate buffer (pH 7.0). They were then incubated at 37 C for 3 hr in phosphate buffer (pH 7.6) with 0.1 mg of trypsin per ml and again washed repeatedly with buffer. Samples were hydrolyzed overnight at 92 C in 6 N HCl and analyzed on a Technicon automatic amino acid analyzer.

^b Values expressed as micromoles ($\pm 1 \mu\text{mole}$) per 100 mg of cell wall.

detail elsewhere (Nickerson, Ph.D. Thesis, University of Cincinnati, 1969), we favor the latter view.

This research was supported by Public Health Service grant A1-06375 from the National Institute of Allergy and Infectious Diseases. K.W.N. was the recipient of Public Health Service predoctoral fellowship 5-F1-GM-29, 756-01 from the National Institute of General Medical Sciences. We also thank Susan Durand for her kind assistance in the use of the Technicon autoanalyzer.

LITERATURE CITED

1. Boman, H. G., K. G. Eriksson-Grennberg, J. Földes, and B. Lindström. 1967. The regulation and possible evolution of a penicillinase-like enzyme in *Escherichia coli*, p. 366-372. In V. V. Koningsberger and L. Bosch (ed.), Regulation of nucleic acid and protein biosynthesis, vol. 10. Elsevier Publishing Co., Amsterdam.
2. Boman, H. G., K. Eriksson-Grennberg, S. Normark, and E. Matsson. 1968. Resistance of *Escherichia coli* to penicillins. IV. Genetic study of mutants resistant to D,L-ampicillin concentrations of 100 $\mu\text{g/ml}$. Genet. Res. 12:169-185.
3. Campbell, J. N., M. Leyh-Bouille, and J.-M. Ghuysen. 1969. Characterization of the *Micrococcus lysodeikticus* type of peptidoglycan in walls of other *Micrococcaceae*. Biochemistry 8:193.
4. Ghuysen, J.-M. 1968. Use of bacteriolytic enzymes in determination of wall structure and their role in cell metabolism. Bacteriol. Rev. 32:425-464.
5. Ghuysen, J.-M., E. Bricas, M. Lache, and M. Leyh-Bouille. 1968. Structure of the cell walls of *Micrococcus lysodeikticus*. III. Isolation of a new peptide dimer, N⁶-[L-alanyl- γ -(α -D-glutamylglycine)]-L-lysyl-D-alanyl-N⁶-[L-alanyl- γ -(α -D-glutamylglycine)]-L-lysyl-D-alanine. Biochemistry 7:1450-1460.
6. Kogut, M., M. R. Pollock, and E. J. Tridgell. 1956. Purification of penicillin-induced penicillinase of *Bacillus cereus* NRRL 569: a comparison of its properties with those of a similarly purified penicillinase produced spontaneously by a constitutive mutant strain. Biochem. J. 62:391-401.
7. Litwack, G. 1960. Experimental biochemistry—a laboratory manual, John Wiley & Sons, Inc., New York.
8. Pickering, B. T. 1966. Components of the cell wall of *Clostridium welchii* (type A). Biochem. J. 100:430-440.
9. Pollock, M. R. 1967. Origin and function of penicillinase: a problem in biochemical evolution. Brit. Med. J. 4:71-77.
10. Schleifer, K. H., and Kandler. 1967. *Micrococcus lysodeikticus*: a new type of cross-linkage of the murein. Biochem. Biophys. Res. Commun. 28:965-972.
11. Tipper, D. J., and J. L. Strominger. 1968. Biosynthesis of the peptidoglycan of bacterial cell walls. XII. Inhibition of cross-linking by penicillins and cephalosporins: studies in *Staphylococcus aureus* in vivo. J. Biol. Chem. 243:3169-3179.