

9-1-2006

Genome of Horsepox Virus

Gustavo A. Delhon

University of Nebraska - Lincoln, gdelhon3@Unl.edu

C. L. Afonso

Agricultural Research Service, United States Department of Agriculture

Z. Lu

Agricultural Research Service, United States Department of Agriculture

L. Zsak

Agricultural Research Service, United States Department of Agriculture

N. T. Sandybaev

Scientific Research Agricultural Institute Zhambylskaya Oblast, Kordaiskiy Rayon, Gvardeiskiy 485444, Republic of Kazakhstan

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/virologypub>

 Part of the [Virology Commons](#)

Delhon, Gustavo A.; Afonso, C. L.; Lu, Z.; Zsak, L.; Sandybaev, N. T.; Kerembekova, U. M.; Zaitsev, V. L.; Kutish, G. F.; and Rock, D. L., "Genome of Horsepox Virus" (2006). *Virology Papers*. Paper 58.
<http://digitalcommons.unl.edu/virologypub/58>

This Article is brought to you for free and open access by the Virology, Nebraska Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Virology Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Gustavo A. Delhon, C. L. Afonso, Z. Lu, L. Zsak, N. T. Sandybaev, U. M. Kerembekova, V. L. Zaitsev, G. F. Kutish, and D. L. Rock

Genome of Horsepox Virus

E. R. Tulman,^{1,2,3} G. Delhon,^{1,4,5} C. L. Afonso,^{1,6} Z. Lu,¹ L. Zsak,¹ N. T. Sandybaev,⁷
U. Z. Kerembekova,⁷ V. L. Zaitsev,⁷ G. F. Kutish,^{1,5,6} and D. L. Rock^{1,5*}

Plum Island Animal Disease Center, Agricultural Research Service, United States Department of Agriculture, Greenport, New York 11944¹; Department of Pathobiology and Veterinary Science² and Center of Excellence for Vaccine Research,³ University of Connecticut, Storrs, Connecticut 06269; Area of Virology, School of Veterinary Sciences, University of Buenos Aires, Buenos Aires, Argentina⁴; Department of Pathobiology, College of Veterinary Medicine, University of Illinois, Urbana, Illinois 61802⁵; Southeast Poultry Research Laboratory, Agricultural Research Service, United States Department of Agriculture, Athens, Georgia 30605⁶; and Scientific Research Agricultural Institute Zhambylskaya Oblast, Kordaiskiy Rayon, Gvardeiskiy 485444, Republic of Kazakhstan⁷

Received 9 May 2006/Accepted 30 June 2006

Here we present the genomic sequence of horsepox virus (HSPV) isolate MNR-76, an orthopoxvirus (OPV) isolated in 1976 from diseased Mongolian horses. The 212-kbp genome contained 7.5-kbp inverted terminal repeats and lacked extensive terminal tandem repetition. HSPV contained 236 open reading frames (ORFs) with similarity to those in other OPVs, with those in the central 100-kbp region most conserved relative to other OPVs. Phylogenetic analysis of the conserved region indicated that HSPV is closely related to sequenced isolates of vaccinia virus (VACV) and rabbitpox virus, clearly grouping together these VACV-like viruses. Fifty-four HSPV ORFs likely represented fragments of 25 orthologous OPV genes, including in the central region the only known fragmented form of an OPV ribonucleotide reductase large subunit gene. In terminal genomic regions, HSPV lacked full-length homologues of genes variably fragmented in other VACV-like viruses but was unique in fragmentation of the homologue of VACV strain Copenhagen B6R, a gene intact in other known VACV-like viruses. Notably, HSPV contained in terminal genomic regions 17 kbp of OPV-like sequence absent in known VACV-like viruses, including fragments of genes intact in other OPVs and approximately 1.4 kb of sequence present only in cowpox virus (CPXV). HSPV also contained seven full-length genes fragmented or missing in other VACV-like viruses, including intact homologues of the CPXV strain GRI-90 D2L/I4R CrmB and D13L CD30-like tumor necrosis factor receptors, D3L/I3R and C1L ankyrin repeat proteins, B19R kelch-like protein, D7L BTB/POZ domain protein, and B22R variola virus B22R-like protein. These results indicated that HSPV contains unique genomic features likely contributing to a unique virulence/host range phenotype. They also indicated that while closely related to known VACV-like viruses, HSPV contains additional, potentially ancestral sequences absent in other VACV-like viruses.

The genus *Orthopoxvirus* includes members of the family *Poxviridae* historically relevant to human health—variola virus (VARV), the etiologic agent of smallpox, and vaccinia virus (VACV), the vaccine virus used to eradicate smallpox (32). Other orthopoxviruses (OPVs), similar to VACV, are zoonotic and significant for human health, including monkeypox virus (MPXV) and cowpox virus (CPXV) (33). Still others, similar to VARV, remain restricted to specific, albeit nonhuman, hosts, including camelpox virus (CMLV) in camels and ectromelia virus (ECTV) in mice. Recent developments have heightened interest in OPV virulence and host range, including the threats of deliberate VARV reintroduction, virulence associated with preemptive smallpox vaccination and use of VACV-based recombinant vaccines, and the introduction of MPXV into the United States (16, 28, 69, 83). Isolation of OPV from infected animals and humans during limited disease outbreaks or from animals in the wild suggests that additional OPVs circulating in nature could represent an emerging disease threat (24, 25, 27, 32, 46, 49, 50, 90).

Given their importance, OPVs have been extensively studied as models of poxviral molecular biology, genomics, genetics, and virus-host interaction (19, 33, 59). Research has revealed that OPVs contain approximately 170 to 230 genes, with those in central genomic regions generally involved in poxviral intracytoplasmic replication and those in terminal genomic regions involved or potentially involved in virus-host interactions, including manipulation of host immune or cellular apoptotic responses (4, 19, 59, 60, 82, 87).

Comparative analysis of completely sequenced OPV genomes, including most known OPV species and several strains of VARV, VACV and the closely related rabbitpox virus (RPXV), MPXV, CMLV, and CPXV has begun to reveal the degree of variability within the genus *Orthopoxvirus*, verifying that terminal genomic regions are the most variable and thus likely to contribute to the virulence and host range characteristics of different OPVs (2, 9, 21, 22, 36, 39, 51, 52, 54, 58, 78, 80, 81). The precise roles and contributions of many variable genes and gene complements in OPV virulence and host range, however, remain to be fully characterized. It is likely that complete genomic data from uncharacterized OPV isolates will aid in OPV gene identification and functional characterization, while also providing information regarding the pathogenic potential of the virus.

* Corresponding author. Mailing address: 2522 Veterinary Medicine Basic Science Building, MC-002, 2001 S. Lincoln Ave., Urbana, IL 61802. Phone: (217) 244-0533. Fax: (217) 244-7421. E-mail: dlrock@uiuc.edu.

Horsepox virus (HSPV) is an OPV causing horsepox, classically known as a poxviral disease of horses. Although common before the 20th century, horsepox is rare today to the point of being considered extinct (14, 44). Multiple clinical forms of horsepox have been described, including a benign, localized form involving lesions in the muzzle and buccal cavity known previously as contagious pustular stomatitis and a generalized, highly contagious form known as equine papular stomatitis (44, 94). Horsepox has also been associated with an exudative dermatitis of the pasterns described as "grease" or grease heel, a clinical syndrome also associated with other infectious and environmental agents (14, 33, 94). Horsepox is differentiated clinically from two other poxviral diseases of horses, equine molluscum contagiosum and Uasin Gishu disease. Equine molluscum contagiosum is a mild, self-limiting cutaneous disease similar to the human disease and is associated with a virus similar to molluscum contagiosum virus (88, 94). Uasin Gishu disease has been described in nonindigenous horses of eastern Africa and is associated with a poorly characterized OPV; however, generalized skin lesions are proliferative and papillomatous and the disease may be chronic in nature (33, 88, 94). HSPV is yet to be characterized molecularly, with no DNA sequence information available. Given the interest in understanding the genetic basis of viral host range and virulence and the relationships between OPVs, we have sequenced and analyzed the genome of a pathogenic field isolate of HSPV.

MATERIALS AND METHODS

Viral DNA isolation, cloning, sequencing, and sequence analysis. The HSPV strain MNR-76 was isolated from sick horses in Bayan-somon of Khentii aimak, Mongolia, in 1976. MNR-76 causes severe disease in horses of the Mongolian breed, including pyrexia, pustular stomatitis with occasional lesions on udders and ears, and especially severe disease in foals and mares, in which death was noted (S. M. Mamadaliyev, personal communication). Viruses were passaged twice in sheep kidney cells, from which viral genomic DNA was extracted as previously described (93). Random DNA fragments were obtained by incomplete enzymatic digestion with Tsp509I endonuclease, cloned into the dephosphorylated EcoRI site of pUC19 plasmids, and grown in *Escherichia coli* DH10B cells (Gibco BRL, Gaithersburg, Md.). Double-stranded DNA templates were purified and sequenced from both ends with M13 forward and reverse primers using dideoxy chain terminator sequencing chemistries and the Applied Biosystems PRISM 3700 automated DNA sequencer (Applied Biosystems, Foster City, CA). Chromatogram traces were base called with Phred (30), which also produced a quality file containing a predicted probability of error at each base position. The sequences were assembled with Phrap (29) and CAP3 (43) using quality files and default settings to produce a consensus sequence with some subsequent manual editing using the Consed sequence editor (37). Gap closure was achieved by primer walking of gap-spanning clones and sequencing of PCR products. Final DNA consensus sequences represented on average sevenfold redundancy at each base position, contained no obvious polymorphisms, and demonstrated a Consed estimated error rate of less than 0.01 error per 10 kb.

Sequence analysis was conducted essentially as previously described (1). Briefly, DNA composition, structure, repeats, and restriction enzyme patterns were analyzed and open reading frame (ORF) maps created using EMBOSS (70), GCG v.10 (Accelrys, Inc., San Diego, CA), and MacVector (Accelrys, Inc) software packages. ORFs longer than 30 amino acids with a methionine start codon were evaluated for coding potential using the GLIMMER (71) computer program, and those greater than 60 amino acids were subjected to similarity searches against nonredundant protein databases and redundant viral protein databases using BLAST (8) and against viral nucleotide databases using TFasta and TFASTX (65, 66). Here, 236 ORFs were annotated and numbered from left to right, with alphabetic subordering given to indicate multiple potential fragments of larger OPV ORFs. Given the predicted nature of all HSPV genes and gene products, ORF names were used throughout the text to indicate both the predicted gene and its putative protein product. Genomic, subgenomic,

and protein alignments and comparisons were done using DIALIGN v2.2.1 (57) using anchors as generated by CHAOS (17), Multi-LAGAN (18), CLUSTAL W (89), BLAST, FASTA (64), SEAVIEW (34), and DOTTER (84) programs. Phylogenetic analyses were conducted on whole-genome sequences and genomic subregions, including a central region used previously for OPV phylogenetic analysis (positions 26800 to 170171) (22, 51), using PHYLIP (31); PHYLO_WIN (34), TREE-PUZZLE (73), and PHYML (40) programs, with evolutionary models selected using MrModeltest 2.2 (62) and additional analyses conducted on alignments in which poorly aligned regions were removed with Gblocks (20).

Nucleotide sequence accession number. The HSPV MNR-76 genome sequence has been deposited in GenBank under accession no. DQ792504.

RESULTS AND DISCUSSION

Organization of the HSPV genome. HSPV MNR-76 genome sequences were assembled into a contiguous sequence of 212,633 bp. The leftmost nucleotide was arbitrarily designated base 1. Similar to other OPVs, the HSPV genome contained 69% A+T nucleotide composition and a central coding region bounded by two identical inverted terminal repeat (ITR) regions.

HSPV ITRs were 7,527 bp and contained elements similar to repetitive and nonrepetitive sequences characterized in other OPVs, including a portion of the terminal hairpin loop-like sequence (positions 1 to 15 from each terminus) and nonrepetitive region 1 (NR1) (positions 21 to 101 from each terminus) and concatemer resolution (position 21 to 40 from each terminus) sequences identical to those present in VACV strain Copenhagen (CPN) (11, 36, 55). Notably, HSPV lacked extensive tandem repetition of terminally located sequences, containing only single copies of the 69-bp (positions 102 to 170, 100% identical to CPN) and 54-bp (positions 518 to 571, 96% identical to CPN) motifs repeated 8.5 to 42 times in VACV strains and RPXV (9, 10, 36, 51). Incomplete copies of 69-bp (positions 171 to 188), 54-bp (positions 572 to 601), and VACV 125-bp repeat-like (positions 494 to 517) motifs flanked complete 69-bp and 54-bp motifs, which were also separated from each other by an NR2-like sequence (positions 189 to 493, 92% identity to CPN positions 2867 to 3171). The HSPV ITR contained eight ORFs initiating and terminating in the ITR, with HSPV001/HSPV207 encompassing the 54-bp and 125-bp motif region (Table 1). These data indicate that while similar to VACV in regions of the ITR, HSPV organizationally resembles other OPVs such as VARV, MPXV, and ECTV which contain fewer or single complete tandem repeat units in their termini (21, 53, 81).

HSPV contained 236 ORFs potentially encoding proteins of 53 to 1,920 amino acids and sharing similarity with those in previously described OPV genomes (Tables 1 and 2). Of these 236 annotated ORFs, 54 were significantly smaller or fragmented forms of 25 larger ORFs present in other OPVs, leaving 182 potentially full-length OPV gene homologues. The HSPV central genomic region contained genes colinear and highly conserved among other OPV genomes, with ORFs HSPV041 to HSPV145 sharing an average 98% amino acid identity with VACV CPN ORFs F1L to A24R and with CPXV GRI-90 ORFs G1L to A25R (Table 2 and data not shown). Genes in this conserved region included those involved in basic replicative functions such as viral transcription and transcript modification, DNA replication, and assembly of intracellular mature and extracellular enveloped virions (IMVs and EEVs,

TABLE 1. HSPV ORFs in terminal genomic regions compared to best-matching ORFs annotated in VACVs, RPXV, and CPXVst

HSPV ORF	Position (length ^b)	VACV ^c				RPXV ^d				CPXV ^e				Putative function/similarity ^h		
		ORF	CPN	WR		Tian		MVA		mO		GRI			BRI	Length
				Length	% Id ^f	ORF Length	ORF Length	ORF Length	ORF Length	ORF Length	ORF Length	% Id ^f	ORF Length			
Left-terminal genomic region																
HSPV001	688-473 (72)	C23L	244	86	001	244	C20L	244	001L	136	L10L	258	001	64	Chemokine binding protein TNFR II-like protein, Crmb	
HSPV002	1547-804 (248)	C22L	122	91	002	61		002L	176	L09L	34	63	003	246		
HSPV003	2723-1676 (349)			004	122					L08L	122	122	005	355		
HSPV004	4570-2810 (587)	C21L	113	100	005	48				L07L	48	113	006	619	Ankyrin repeat protein	
		C20L	103	69	006	64				L06L	128	109				
		C19L	259	90	007	109				L04L	109	77				
				008	112							149				
HSPV005a	5051-4779 (91)	C18L	150	98						L03L	93	93	008	672	Ankyrin repeat protein	
HSPV005b	5530-5081 (150)	C17L	386	92				003L	102	L02L	416	163	008	672		
HSPV005c	6797-5607 (397)	C16L	181	97				004L	233			385	008	672		
								189R	188	L01L	147	184	009	153		
HSPV006	7419-6961 (153)												010	215		
HSPV007	8041-7589 (151)												011	215		
HSPV008	9327-8509 (273)												012	202	BTB/POZ domain protein	
HSPV009	10133-9681 (151)												013	111	Chemokine binding domain protein	
HSPV010	10529-10197 (111)												014	110	TNFR, CD30-like protein	
HSPV011a	11413-10625 (263)												015	110	Ankyrin repeat protein	
HSPV011b	12197-11334 (288)												016	764		
HSPV011c	12637-12191 (149)												017	764		
HSPV012	14485-13175 (437)												018	171	Ankyrin repeat protein	
HSPV013	15127-14852 (92)												019	796	Ankyrin repeat protein	
HSPV014a	15498-15220 (93)												020	170	VACV C7L-like protein	
HSPV014b	16216-15473 (248)												021	139	Growth factor	
HSPV014c	16975-16367 (203)												022	331	ECTV p28-like host range protein	
HSPV014d	17605-17111 (165)												023	242	Secreted IL-18 binding protein	
HSPV015a	17913-17695 (73)												024	126	Ankyrin repeat host range protein	
HSPV015b	18205-17960 (82)												025	668		
HSPV016	18365-18784 (140)	C11R	142	95	009	140	C18R	140	005R	140	005R	140	026	668		
HSPV017	19934-18942 (331)	C10L	331	96	010	331	C17L	331	006L	326	007L	331	027	632	Ankyrin repeat protein	
HSPV018	20448-20765 (106)			011	181	C16R	44	007R	91	008R	239	008	028	242		
HSPV019	21707-21330 (126)			013	126	C15L	68	008L	120	009L	124	009	029	126		
HSPV020a	22038-21769 (90)			014	237	C14L	142	010L	142	011L	142	142	030	409		
HSPV020b	22479-22054 (142)			015	137	C13L	115	011L	135	012L	137	137	031	409		
HSPV020c	22999-22571 (143)			016	77	C12L	77	012L	90	013L	77	77	032	668		
HSPV020d	23508-23278 (77)			017	71	C11L	76	013L	71	014L	71	71	033	668		
HSPV020e	23745-23533 (71)			018	60	C10L	59	014L	59		59	59	034	668		
HSPV021	24072-23884 (63)			019	634	C9L	634	014L	109	016L	634	634	035	668		
HSPV022	26149-24248 (634)	C9L	634	99	019	634	C9L	634	015L	96		634	036	68		
								016L	297				037	614	Ankyrin repeat protein	
HSPV023	26725-26195 (177)	C8L	184	89	020	177	C8L	177	017L	177	019L	177	038	186		
HSPV024	27249-26800 (150)	C7L	150	97	021	150	C7L	150	018L	150	020L	150	039	150	Host range protein	
HSPV025	27949-27485 (155)	C6L	151	89	022	151	C6L	151	019L	157	021L	151	040	155		
HSPV026a	28280-28089 (64)	C5L	204	98	023	204	C5L	204			022L	204	041	69		
HSPV026b	28690-28328 (121)	C5L	204	98	023	204	C5L	204			022L	204	042	125	BTB/POZ domain protein	
HSPV027a	29175-28765 (137)	C4L	316	97	024	316	C4L	255			023L	316	043	316		
HSPV027b	29689-29489 (67)	C4L	316	98	024	316	C4L	255			023L	316	044	316		
HSPV028	30547-29759 (263)	C3L	263	96	025	263	C3L	263			025L	263	045	263	Complement binding protein	
HSPV029	32153-30618 (512)	C2L	512	97	026	512	C2L	512			026L	512	046	512	Kelch-like protein	
HSPV030	32891-32223 (223)	C1L	224	97	027	229	C1L	224			027L	224	047	231		
HSPV031	33231-32881 (117)	N1L	117	100	028	117	N1L	117	020L	113	028L	117	048	117	TLR/IL-1R/TNFR signaling inhibitor	
HSPV032	33895-33371 (175)	N2L	175	97	029	175	N2.L*	175	021L	170	029L	175	049	177	Alpha-amanitin-sensitive protein	

TABLE 1—Continued

HSPV ORF	Position (length ^b)	VACV ^c			MVA			RPXV ^d			CPXV ^e			Putative function/similarity ^h						
		CPN	WR		Tian	ORF ^g	Length		ORF	Length	ORF	Length	% Id ^f		BRI					
			Length	% Id ^f			Length	ORF								Length	ORF	Length	ORF	
HSPV163	161656–161000 (219)	A41L	219	166	219	A52L	219	153L	219	212L	219	149	219	A43L	219	97	178	218	Secreted immunomodulatory protein	
HSPV164	161827–162225 (133)	A42R	133	167	133	A53R	133	154R	128	213R	133	150	133	A44R	133	100	179	133	Profilin-like protein	
HSPV165	162266–162850 (195)	A43R	194	168	194	A54R	194	155R	190	214R	194	151	194	A45R	196	95	180	194	Hydroxysteroid dehydrogenase	
HSPV166	164226–163189 (346)	A44L	346	170	346	A55L	346	157L	346	216L	346	153	346	A47L	346	98	182	346	Superoxide dismutase-like protein	
HSPV167	164273–164647 (125)	A45R	125	171	125	A56R	125	158R	121	217R	125	154	125	A48R	125	98	183	125	TLR/IL-1R signaling inhibitor	
HSPV168	164640–165359 (240)	A46R	214	100	172	240	A57R	210	159R	240	218R	240	155	240	A49R	240	97	184	242	Thymidylate kinase
HSPV169	166183–165452 (244)	A47L	244	99	173	252	A58L	252	160L	238	220L	252	156	244	A50L	244	97	185	244	
HSPV170	166282–166893 (204)	A48R	204	100	174	227	A59R	204	161R	204	221R	204	157	204	A51R	227	99	186	227	DNA ligase
HSPV171	166944–167429 (162)	A49R	162	96	175	162	A60R	162	162R	162	222R	162	158	162	A52R	162	95	187	162	
HSPV172	167464–169119 (552)	A50R	552	98	176	552	A61R	552	163R	552	223R	552	159	552	A53R	552	97	188	554	
HSPV173a	169175–169414 (80)	A51R	334	88	177	334	A62R*	334	164R	310	226R	73	84	A54R	334	92	189	334		
HSPV173b	169359–170168 (270)	A51R	334	96	177	334	A62R	334	164R	310	227R	266	126	A54R	334	96	189	334		
HSPV174	170241–170810 (190)	A52R	190	98	178	190	A63R*	190	170R	190	228R	190	160	190	A55R	190	96	190	190	TLR/IL-1R signaling inhibitor
HSPV175	171134–171439 (102)	A53R	108	91	179	103	*	137	171R	177	242R	558	166	102	A56R	186	88	191	186	TNFR, CrmC
HSPV176	171969–173660 (564)	A55R	564	99	180	564	A65R*	564	165R	315	232R	564	162	564	A57R	564	97	193	563	Kelch-like protein
HSPV177	173713–174654 (314)	A56R	315	98	181	314	A66R	315	165R	315	233R	310	163	206	A58R	314	94	194	297	EEV hemagglutinin
HSPV178a	174675–174860 (62)	A57R	151	93	182	151	A67R	151	166R	97	235R	151	164	151	A59R	197	96	195	197	Guanylate kinase
HSPV178b	174975–175265 (97)	A57R	151	100	182	151	A67R	151	167R	300	236R	300	165	300	B1R	300	99	196	209	Ser/Thr protein kinase, DNA replication
HSPV179	175419–176318 (319)	B1R	300	99	183	300	B1R	300	168R	96	238R	219	169R	143	B2R	503	93	197	505	
HSPV180a	176412–177068 (210)	B2R	219	96	184	219	B2R	219	172R	409					B2R	503	93	197	505	Schlafen-like protein
HSPV180b	177107–177478 (124)	B3R	124	96	185	167	B3R	124	170R	179	240R	124	124	124	B2R	503	91	197	558	Ankyrin repeat protein
HSPV181	178138–179811 (558)	B4R	558	98	186	558	B4R	558	171R	177	242R	558	166	558	B3R	558	95	198	558	
HSPV182	179917–180867 (317)	B5R	317	97	187	317	B5R	317	173R	317	243R	317	167	317	B4R	317	96	199	317	EEV host range protein
HSPV183a	180953–181285 (111)	B6R	173	99	188	173	B6R	173	174R	173	244R	173	168	173	B5R	183	93	200	179	
HSPV183b	181288–181482 (65)	B6R	173	100	188	173	B6R	173	174R	173	244R	173	168	173	B5R	183	93	200	179	
HSPV184	181523–182068 (182)	B7R	182	100	189	182	B7R	182	175R	177	246R	182	169	182	B6R	182	97	201	181	Chemokine binding domain
HSPV185	182149–182964 (272)	B8R	272	97	190	272	B8R	272	176R	226	247R	272	170	272	B7R	271	97	202	266	IFN-γ receptor
HSPV186	183054–183284 (77)	B9R	77	100	191	77	B9R	77	177R	72	248R	77	77	61	B8R	221	97	203	225	MYXV M-T4-like protein
HSPV187	183384–183746 (121)	B10R	166	99	192	166	B10R	166	178R	158	249R	166	172	166	B9R	501	94	204	501	Kelch-like protein
HSPV188	183821–184054 (78)	B11R	88	96	193	72	B11R	76	179R	74	250R	72	173	72	B10R	105	96	205	90	
HSPV189	184124–184972 (283)	B12R	283	98	194	283	B12R	283	180R	283	251R	283	174	283	B11R	283	98	206	285	Ser/Thr protein kinase
HSPV190	185074–186108 (345)	B14R	222	97	195	345	B14R	222	181R	116	253R	222	175	345	B12R	345	93	207	341	Serpin, SPI-2
HSPV191	186186–186632 (149)	B15R	149	97	196	149	B15R	149	183R	143	254R	149	176	149	B13R	149	98	208	149	
HSPV192	186736–187713 (326)	B16R	290	97	197	326	B16R	290	184R	326	255R	326	207	134	B14R	326	95	209	326	IL-1 receptor
HSPV193	188784–187765 (340)	B17L	340	97	198	340	B17L	340	185L	340	257L	340	177	340	B15L	340	96	210	340	
HSPV194	188921–190642 (574)	B18R	574	98	199	574	B18R	574	186R	574	258R	413	178	574	B16R	574	95	211	574	Ankyrin repeat protein
HSPV195	190711–191775 (355)	B19R	353	98	200	351	B19R	353	187R	234			179	351	B17R	351	91	212	366	IFN-α/β binding protein
HSPV196	191850–194222 (791)	B20R	127	96	202	53	B20R*	613	203	309	180	791	180	791	B18R	795	94	213	800	Ankyrin repeat protein
HSPV197	194331–195971 (547)			203	309										B19R	557	94	215	557	Kelch-like protein
HSPV198	196272–197342 (357)	C12L	353	97	205	353	C19L	353	004L	353	005	357	004L	353	B20R	375	95	217	372	Serpin, SPI-1
HSPV199	197515–198090 (192)	C13L	65	90	206	190			003L	190	004	192	003L	190	B21R	190	94	218	198	Chemokine binding domain protein
HSPV200	198347–204106 (1,920)			82	98				188R	70	002L	89	002L	89	B22R	1,033	97	219	1,019	VARV B22R-like protein
HSPV201a	204447–204806 (120)	B21R	91	96					189R	188			91	K1R	581	96	220	579	Ankyrin repeat protein	
HSPV201b	204961–205161 (67)														K1R	581	90	220	579	

HSPV202 205215-205673 (153)	B22R	181	97	189R	188	L01L	147	002	184	I1R	153	97	222	153
HSPV203a205837-207027 (397)	B23R	386	92	190R	233	L02L	416		385	I2R	672	94	223	672
				191R	102									
HSPV203b207104-207553 (150)	B24R	150	98			L03L	98		163	I2R	672	92	223	672
HSPV203c207583-207855 (91)						L03L	93			I2R	672	97	223	672
HSPV204 208064-209824 (587)	B25R	259	90		211	L04L	100		140	I3R	586	95	225	619
	B26R	103	69		212	L06L	128		109					
	B27R	113	100		213	L07L	48		113					
					214				113					
HSPV205 209912-210958 (349)	B28R	122	91	192R	176	L08L	122		122	I4R	351	95	226	355
						L09L	34		63					
HSPV206 211087-211830 (248)	B29R	244	86	193R	136	L10L	258		258	I5R	255	87	227	246
HSPV207 211946-212161 (72)													229	64

^a Boldface indicates ORFs >10% different in length from intact orthologues from CPXV GRI-90 or BRI. Names of ORF homologues have been abbreviated here for simplicity and lack the following prefixes for the following viruses: VACV WR, WR; T, Tian Tan; MVA, MVA; m0LTR, ORFs in the m0 long terminal repeat indicated here with prefix L; m0, unique m0 ORFs; RPXV, RPXV; CPXV, BRI.
^b All lengths are in amino acids.
^c VACV strains (accession numbers): CPN (M35027); WR (AY243312); Tian, Tian Tan (AF095689); MVA (U94848); m0, LCJ6m0 (AY678277). Larger ORFs matching multiple HSPV ORFs are VACV CPN ORFs C5L, C4L, MIL, A26L, A51R, A57R, and B6R; VACV WR ORFs 014, 023, 024, 030, 177, 182, and 188; VACV Tian Tan ORFs C5L, C4L, MIL, A27L, A62R, A67R, and B6R; VACV MVA ORFs 164R, 174R, and 189R; and VACV m0 ORFs L03L, 022L, 023L, 030L, 185L, and 244R.
^d RPXV strain Utrecht. RPXV ORF lengths lacking an ORF designation indicate ORFs lacking translation products annotated in sequence AY484669. Larger ORFs matching multiple HSPV ORFs are the 409-amino-acid-long ORF, ORF 015, ORF 016, ORF 022, the 233-amino-acid-long ORF, ORF 164, and ORF 168.
^e CPXV strains (accession numbers): GRI, GRI-90 (X94355); BRI (AF482758). Larger ORFs matching multiple HSPV ORFs are GRI-90 ORFs D4L, D14L, C3L, C4L, C9L, C15L, C16L, A54R, A59R, B2R, B3R, K1R, and L2R and BRI ORFs 008, 016, 019, 020, 025, 033, 039, 158, 189, 195, 197, 200, 220, and 223.
^f % Id, percent amino acid identity in local BLAST match.
^g Asterisks indicate ORFs resequenced/reannotated by Upton et al. (92) as present, intact, or fused to a subsequent ORF in the Tian Tan genome.
^h Abbreviations: IL-18, interleukin-18; TLR, Toll-like receptor; PKR, double-stranded RNA-dependent protein kinase; IEV, intracellular enveloped viron; IFN- γ , gamma interferon; MYXX, myxoma virus; dsRNA, double-stranded RNA.

respectively), indicating that HSPV is similar to other OPVs in these functions (59) (Table 2).

HSPV terminal genomic regions were similar to other OPVs in that they contained a homologous subset of the sequence and intact ORFs present in various strains of CPXV, viruses found to contain a relatively complete OPV genotype and thus thought to be viruses from which other OPV lineages are derived following gene fragmentation and loss (Table 1; Fig. 1) (75, 79). Many of these ORFs have been characterized in other OPVs as affecting viral virulence, host range, and modification of host responses, including apoptosis and innate and adaptive immune mechanisms (59, 60, 82). However, the specific subset of genes present in HSPV was unique relative to other OPVs, containing terminal genomic sequences not characteristic of currently known OPVs and including approximately 1.4 kb of sequence found only in CPXV (located between positions 15453 and 16985) (Fig. 1).

Phylogenetic analysis. Phylogenetic analysis of OPV genomic regions, including the highly conserved central region and parts of the more variable terminal regions, indicated that HSPV is closely related to sequenced strains of VACV and RPXV, falling very close to or within this VACV subgroup (referred to here as VACV-like viruses) relative to other OPVs (Fig. 2). These results are consistent with those obtained previously for OPVs, with VACV-like viruses closely related to each other compared to other OPV species, and they indicated that HSPV is a VACV-like virus (21, 38, 51). As a VACV-like virus, HSPV also shares a closer relationship with CPXV strain GRI-90 than with CPXV strain Brighton Red (BRI), consistent with previous OPV phylogenetic analyses and indicating the distinct nature of CPXV species despite the relative conservation in gene content (Fig. 1 and 2) (21, 38, 51). Similarly, a close relationship was observed between HSPV and VACV using concatenated right terminal OPV gene sequences used previously for OPV phylogenetic analysis (HSPV177, HSPV179, HSPV182, and HSPV191; data not shown) (38). These results indicate that HSPV and VACV are very similar phylogenetically and share a relatively recent common ancestor. Notably, HSPV had a slightly greater estimated distance to VACV-like isolates than they demonstrated to each other, with HSPV tending to fall outside the rest of the VACV-like cluster (Fig. 2). These data suggested that, while very closely related, HSPV is phylogenetically distinct from other characterized VACV-like viruses.

Comparison of HSPV with VACV-like viruses. Given the close phylogenetic relationship between HSPV and VACV-like viruses, HSPV ORFs were compared to VACV-like homologues in the more variable terminal genomic regions which tend to contain genes dispensable for basic replicative processes but important for specific virus-host interaction and aspects of virulence and host range (Fig. 1; Table 1). While HSPV maintained a high level of amino acid identity where homologous terminal region ORFs were present (average of 95% amino acid identity to CPN), we focused here on comparison of HSPV and VACV in genes likely fragmented relative to CPXV and other OPVs. Overall, these differences often involved genes that are members of multigene families and/or homologues of genes shown or thought to affect OPV virulence or host range, among them those that code for ankyrin repeat proteins, kelch-like proteins, and tumor necrosis factor recep-

TABLE 2. HSPV ORFs in central genomic regions compared to orthologues annotated in VACV CPN^a

HSPV ORF	Position (length ^c)	VACV CPN		Putative function/similarity
		ORF	Length	
HSPV063	61662–63362 (567)	E6R	567	
HSPV064	63447–63944 (166)	E7R	166	
HSPV065	64072–64890 (273)	E8R	273	Virion core protein
HSPV066	67919–64902 (1,006)	E9L	1,006	DNA polymerase
HSPV067	67951–68235 (95)	E10R	96	IMV redox protein
HSPV068	68622–68236 (129)	E11L	129	Virion core protein
HSPV069	70609–68612 (666)	O1L	666	
HSPV070	70983–70660 (108)	O2L	108	Glutaredoxin
HSPV071	72067–71132 (312)	I1L	312	DNA binding virion core protein
HSPV072	72304–72077 (76)	I2L	73	
HSPV073	73114–72308 (269)	I3L	269	DNA binding phosphoprotein
HSPV074a	73439–73200 (80)	I4L ^b	771	Ribonucleotide reductase large subunit
HSPV074b	74885–73566 (440)	I4L	771	
HSPV074c	75213–74842 (124)	I4L	771	
HSPV074d	75503–75216 (96)	I4L	771	
HSPV075	75770–75534 (79)	I5L	79	IMV membrane protein
HSPV076	76937–75792 (382)	I6L	382	Telomere binding protein
HSPV077	78201–76933 (423)	I7L	423	Virion core proteinase
HSPV078	78207–80234 (676)	I8R	676	RNA helicase NPH-II
HSPV079	82016–80244 (591)	G1L	591	Metalloprotease
HSPV080	82342–83001 (220)	G3L	220	
HSPV081	82348–82016 (111)	G2R	111	Transcriptional elongation factor
HSPV082	83348–82977 (124)	G4L	124	Glutaredoxin 2
HSPV083	83351–84652 (434)	G5R	434	Virion core protein
HSPV084	84663–84851 (63)	G5.5R	63	RNA polymerase subunit RPO7
HSPV085	84856–85350 (165)	G6R	166	
HSPV086	86433–85321 (371)	G7L	371	Virion core protein
HSPV087	86464–87243 (260)	G8R	260	Late transcription factor VLTF-1
HSPV088	87266–88285 (340)	G9R	340	Myristylated protein
HSPV089	88289–89038 (250)	L1R	250	Myristylated IMV envelope protein
HSPV090	89073–89333 (87)	L2R	87	
HSPV091	90378–89329 (350)	L3L	350	
HSPV092	90403–91155 (251)	L4R	251	DNA binding virion core protein
HSPV093	91168–91551 (128)	L5R	128	IMV membrane protein
HSPV094	91511–91969 (153)	J1R	153	IMV membrane protein
HSPV095	91988–92518 (177)	J2R	177	Thymidine kinase
HSPV096	92587–93585 (333)	J3R	333	Poly(A) polymerase small subunit
HSPV097	93503–94057 (185)	J4R	185	RNA polymerase subunit RPO22
HSPV098	94585–94187 (133)	J5L	133	
HSPV099	94692–98549 (1,286)	J6R	1,286	RNA polymerase subunit RPO147
HSPV100	99064–98552 (171)	H1L	171	Tyr/Ser protein phosphatase
HSPV101	99078–99644 (189)	H2R	189	IMV membrane protein
HSPV102	100624–99653 (324)	H3L	324	IMV envelope protein
HSPV103	103012–100628 (795)	H4L	795	RNA polymerase-associated protein
HSPV104	103198–103830 (211)	H5R	203	Late transcription factor VLTF-4
HSPV105	103834–104775 (314)	H6R	314	DNA topoisomerase IB
HSPV106	104815–105252 (146)	H7R	146	
HSPV107	105299–107830 (844)	D1R	844	mRNA capping enzyme large subunit
HSPV108	108225–108935 (237)	D3R	237	Virion core protein
HSPV109	108232–107795 (146)	D2L	146	Virion core protein
HSPV110	108938–109591 (218)	D4R	218	Uracil DNA glycosylase
HSPV111	109626–111980 (785)	D5R	785	NTPase, DNA replication
HSPV112	112024–113934 (637)	D6R	637	Early transcription factor small subunit
HSPV113	113964–114446 (161)	D7R	161	RNA polymerase subunit RPO18
HSPV114	115326–114415 (304)	D8L	304	IMV membrane protein, cell binding
HSPV115	115368–116006 (213)	D9R	213	MutT motif
HSPV116	116006–116749 (248)	D10R	248	MutT motif
HSPV117	118648–116756 (631)	D11L	631	NPH-I, transcription termination factor
HSPV118	119546–118686 (287)	D12L	287	mRNA capping enzyme small subunit
HSPV119	121232–119580 (551)	D13L	551	Rifampin resistance protein
HSPV120	121708–121259 (150)	A1L	150	Late transcription factor VLTF-2
HSPV121	122403–121732 (224)	A2L	224	Late transcription factor VLTF-3
HSPV122	122630–122403 (76)	A2.5L	76	Virion redox protein
HSPV123	124579–122648 (644)	A3L	644	Virion core protein P4b
HSPV124	125477–124635 (281)	A4L	281	Virion core protein
HSPV125	125515–126006 (164)	A5R	164	RNA polymerase subunit RPO19

Continued on facing page

TABLE 2—Continued

HSPV ORF	Position (length ^c)	VACV CPN		Putative function/similarity
		ORF	Length	
HSPV126	127124–126009 (372)	A6L	372	
HSPV127	129280–127151 (710)	A7L	710	Early transcription factor large subunit
HSPV128	129334–130197 (288)	A8R	288	Intermediate transcription factor VITF-3
HSPV129	130501–130196 (102)	A9L	99	IMV membrane protein
HSPV130	133177–130505 (891)	A10L	891	Virion core protein P4a
HSPV131	133192–134145 (318)	A11R	318	Nonstructural protein
HSPV132	134725–134153 (191)	A12L	192	Virion core protein
HSPV133	134961–134752 (70)	A13L	70	IMV membrane protein
HSPV134	135341–135072 (90)	A14L	90	IMV membrane protein
HSPV135	135519–135361 (53)	A14.5L	53	IMV membrane protein
HSPV136	135793–135512 (94)	A15L	94	Virion core protein
HSPV137	136913–135780 (378)	A16L	378	Myristylated IMV membrane protein
HSPV138	137527–136919 (203)	A17L	203	Phosphorylated IMV membrane protein
HSPV139	137542–139020 (493)	A18R	493	DNA helicase, transcriptional elongation
HSPV140	139237–139007 (77)	A19L	77	
HSPV141	139590–140867 (426)	A21L	426	DNA polymerase processivity factor
HSPV142	139591–139241 (117)	A20R	117	IMV membrane protein
HSPV143	140833–141360 (176)	A22R	176	Holliday junction resolvase
HSPV144	141383–142528 (382)	A23R	382	Intermediate transcription factor VITF-3
HSPV145	142528–146019 (1,164)	A24R	1,164	RNA polymerase subunit RPO132

^a Boldface indicates ORFs >10% different in length from intact orthologues from CPXV GRI-90 or Brighton Red.

^b 14L is a larger ORF matching multiple HSPV ORFs.

^c Lengths are in amino acids.

tors (TNFRs) (4, 45, 77). While terminal-region genotypes vary both among OPVs and between known VACV-like viruses, HSPV contained features similar to known VACV-like viruses relative to other OPVs and features that were quite novel (Table 1; Fig. 1).

HSPV genetic features similar to VACV. Genotypic similarity between HSPV and other VACV-like viruses included a number of genes that were fragmented relative to CPXV and occasionally relative to other OPVs. These genes included several which were fragmented or arranged in a similar fashion between HSPV and VACV-like viruses, commensurate with their close phylogenetic relationship (Table 1; Fig. 2). HSPV genes sharing similar ORF fragments with those in certain VACVs include HSPV005/HSPV203 and HSPV020, genes encoding ankyrin proteins and fragmented or missing in most OPVs (Fig. 1A). HSPV005b/HSPV203b in the ITR represents the same fragment of GRI-90 D4L/I2R as CPN C18L/B24R. HSPV020a to -e and similar ORFs in VACV are homologous fragments of CPXV CHOhr, a gene which enables replication of VACV in the normally nonpermissive CHO cell line and affects eukaryotic initiation factor 2 α (eIF2 α) phosphorylation in HeLa cells (41, 85). Other HSPV ORFs with similar VACV fragments included HSPV146d, HSPV180, and HSPV186. HSPV146d encodes the same 725-amino-acid amino-terminal fragment of the A-type inclusion (ATI) protein present in several VACV-like viruses and expressed in some as a soluble 94-kDa protein (26). HSPV186 is a VACV-like ORF fragment homologous to the amino-terminal region of the OPV homologue of myxoma virus M-T4, a protein important for virulence and infection of lymphocytes by myxoma virus (12). The HSPV186 homologue is expressed in VACV strain Western Reserve (WR); however, deletion mutants were not affected for viral growth in vitro or virulence in mice (68). While amino-terminal M-T4-like fragments are also present in certain

strains of MPXV (22, 52), the large nucleotide deletion affecting HSPV186 was characteristic of VACV (Fig. 1C). Also characteristic of VACV are homologues of HSPV180a and HSPV180b (CPN B2R and B3R, respectively), apparent fragments of a larger ORF intact in all OPV species other than VACV and VARV and previously annotated as similar to cellular Schlafen, a family of variably sized proteins with the prototypical 337-amino-acid murine Schlafen 1 recently shown to target cyclin D1 pathways during induction of cellular mid-G₁ cell cycle arrest (15, 39). Notably, HSPV180a and HSPV180b revealed the bipartite nature of the larger OPV homologue, with Schlafen similarity present in the HSPV180b-like (carboxyl-terminal) region and the HSPV180a-like (amino-terminal) region sharing similarity with the putative B2R homologue of *Melanoplus sanguinipes* entomopoxvirus (MSV237) and limited similarity with ORFs of unknown function (p26) from nucleopolyhedrosis viruses (data not shown). While maintenance of these two domains as separate ORFs in HSPV and VACV conceivably suggests function, HSPV180b and VACV orthologues lack carboxyl-terminal sequences both present in the intact OPV ORF and similar to the carboxyl terminus of cellular Schlafen. Overall, similar fragmentation patterns between HSPV and VACV potentially represent shared, derived characters.

Several genes fragmented in HSPV were also fragmented in certain VACV-like isolates but intact in others (Table 1). HSPV ORF fragments with intact homologues in certain VACVs included HSPV018, HSPV161, HSPV173a and -b, and HSPV175. HSPV018 is an amino-terminal fragment homologue of the ECTV p28 ubiquitin ligase, a protein critical for ECTV virulence and macrophage host range and having intact homologues in all other OPV species (74, 87) (Fig. 1). While this gene is also fragmented in several VACV strains, intact homologues have been identified in VACV strains IHD-W and




FIG. 1. Schematic comparison of HSPV left (A) and right (B and C) terminal genomic regions to those of other orthopoxviruses. Virus names were abbreviated as follows and correspond to sequences from the following GenBank accession numbers in parentheses: VACV CPN (M35027); VACV WR (AY243312); Tian, VACV Tian Tan (AF095689); VACV MVA (U94848); mO, VACV Lister isolate LC16m0 (AY678277); RPXV, RPXV Utrecht (AY484669); GRI, CPXV GRI-90 (X94355); MPXV, MPXV Zaire-96-1-16 (AF380138); ECTV, ECTV Moscow (AF012825); CMLV, CMLV M-96 (AF438165); VARV, VARV Bangladesh-1975 (L22579); CPXV BRI (AF482758). Heavy lines indicate nucleotide sequences; boxes indicate ORFs matching those annotated in HSPV and those in genomic regions absent in HSPV. ORF names and genomic positions in kilobase pairs (K) are indicated for HSPV and CPXV Brighton Red, as are names of ORFs absent in these two species. Hatching indicates ORFs different in length (>10%) from intact orthologues from CPXV GRI-90 or Brighton Red. Red ORFs indicate HSPV ORFs intact or carried on sequences that are absent relative to VACV-like viruses. Asterisks indicate ORFs resequenced/reannotated by Upton et al. (92) as present or intact in the Tian Tan genome. Large solid-lined boxes indicate sequences matching the global alignment only at the opposite genomic terminus; dashed boxes indicate where sequences located in the opposite genomic terminus match the global alignment. Red lines indicate ITR sequence in each virus and are unaligned on the terminal side of HSPV002/HPSV206. Panel A is presented at a different scale relative to panels B and C.




FIG. 1—Continued.

Lister and in RPXV (51, 58, 91). Similarly, HSPV173a and -b resembled homologous ORFs in VACV Lister and RPXV and fragments of the CPN A51R gene intact in other VACV strains and all other OPVs. HSPV161 was a homologue of CPN A39R, a secreted semaphorin affecting viral virulence and host inflammatory responses during infection, but, similarly to ho-

mologues in WR and other VACV strains, contained a carboxyl-terminal truncation that may predict a nonfunctional product (35). HSPV175, similar to several VACV-like viruses, encoded a truncated copy of the intact CrmC TNFR-like protein encoded by VACV strains Lister, Evans, and USSR (5). HSPV039 and HSPV187 were fragmented genes with homo-




FIG. 2. Phylogenetic analysis of HSPV central genomic regions. Conserved HSPV central genomic nucleotide sequences (positions 26800 to 170171) corresponding to regions used previously for OPV phylogenetic analysis (51) were aligned with homologous OPV sequences using DIALIGN, and gapped regions were realigned with CLUSTAL W and trimmed with Gblocks. The unrooted tree for 124,677 aligned characters was generated using maximum likelihood with general time reversible correction for multiple substitutions, four-category discrete gamma model, estimation for proportion of invariant residues, and 100 bootstrap replicates as implemented in PHYML. Bootstrap values greater than 70 are indicated at appropriate nodes; dots indicate values of 100. Homologous nucleotide sequences from the following viruses and accession numbers were compared: VACV strain CPN, M35027; VACV WR, AY243312; VACV Lister (Elstree) vaccine consensus (Lis), AY678276; VACV Lister-derived LC16m0 (m0), AY678277; VACV Tian Tan (Tian), AF095689; VACV MVA, U94848; RPXV Utrecht (RPXV), AY484669; CPXV strain GRI-90 (X94355); CPXV BRI, AF482758; MPXV strain Zaire-96-I-16 (MPXV ZAI), AF380138; MPXV WRAIR7-61 (MPXV W61), AY603973; MPXV USA_2003_039 (MPXV U39), DQ011157; CMLV strain M-96 (CMLV M96), AF438165; CMLV CMS, AY009089; VARV strain Bangladesh-1975 (VARV BAN), L22579; VARV India-1967 (VARV IND), X69198; VARV Garcia-1966 (VARV GAR), Y16780; ECTV strain Moscow (ECTV MOS), AF012825. The scale indicates estimated distance. Identical topologies at supported nodes were obtained using additional maximum likelihood analyses as implemented in TREE-PUZZLE, using neighbor-joining and maximum parsimony as implemented in PHYLO_WIN and PHYLIP, respectively, and using an unedited alignment (146,439 characters) (data not shown). Similar topologies were also obtained using similar analyses on whole-genomic alignments (data not shown).

logues fragmented in all VACV-like viruses but with VACV-like homologues fragmented in a pattern distinct from those in HSPV. HSPV039 was similar to both CPN K5L and K6L fragments of the OPV monoglyceride lipase-like gene but was much closer in size to the intact CPXV homologue, and HSPV187 was a smaller fragment of the CPXV GRI-90 B9R kelch-like protein. While HSPV175, HSPV039, and HSPV187 homologues were fragmented in both HSPV and most VACV-like viruses, these genes were also disrupted in most other OPVs (Fig. 1).

HSPV also contained intact genes whose homologues were intact in certain VACV-like viruses but disrupted in others, similar to genes recently described in the RPXV genome (Table 1) (51). HSPV002/HSPV206 in the ITR encoded the OPV 35-kDa secreted chemokine binding protein and, similarly to the functional, full-length protein expressed by VACV Lister

and other OPVs, lacked the amino-terminal mutation preventing expression of functional protein in CPN, WR, and VACV strain Tian Tan (6). HSPV147 was an intact copy of the gene encoding P4c, a protein involved with direction of IMV to insoluble ATIs but with homologues fragmented or absent in CPN, Tian Tan, and modified vaccinia Ankara (MVA). HSPV190 was only the third intact VACV-like orthologue of the serine proteinase inhibitor (serpin) 2 (SPI-2) to be identified, and HSPV198 was an intact orthologue of the SPI-1 gene intact in most VACV-like viruses but transposed to the opposite terminus in RPXV and VACV CPN, Tian Tan, and Lister and absent in MVA (Fig. 1C). Intact SPI-1 and SPI-2 exhibit antiapoptotic and/or anti-inflammatory activity through inhibition of caspases and have been shown to affect viral virulence and/or host range (48, 59, 82, 87). HSPV196 encodes an intact ankyrin repeat protein truncated by deletion in all VACV-like viruses except RPXV, where the homologue was recently identified as unique among VACV-like viruses in that the entire nucleotide region encompassing the gene was present (51). Similarly, HSPV199 encodes an intact homologue of the BRI CPXV218 chemokine binding protein, with intact homologues also encoded in the right terminus of WR and in the left terminus of VACV Lister and RPXV (Fig. 1) (7). Overall, different fragmentation patterns or gene loss between HSPV genes and VACV homologues may indicate sequence divergence after functional gene loss or, alternatively, could conceivably reflect independent loss of gene function in different VACV-like lineages during convergent adaptation toward similar virulence or host range phenotypes. Gene loss near ITR boundaries may reflect loss during terminal transposition events (47, 61). These phenomena would help explain gene fragmentation that is variable both within the VACV-like lineage and between OPV species.

HSPV genetic features distinct from VACV. Despite sharing specific genomic and genotypic features with some or all known VACV-like viruses within the range of VACV-like genotypic heterogeneity, HSPV contained many features that were unique. These included genes uniquely intact in HSPV but for which homologous nucleotide sequence was present in other VACVs, and they included HSPV genes, both intact and fragmented, that were associated with nucleotide sequences completely novel among VACV-like viruses, resulting in terminal genomic regions encoding additional proteins and protein fragments resembling those in CPXV (94% average amino acid identity to CPXV GRI-90 orthologues) (Fig. 1; Table 1). Finally, HSPV demonstrated unique fragmentation of several genes, including those that were intact in all or most other known VACV-like viruses.

HSPV contained in the ITRs intact genes that are fragmented or absent in all other VACVs (Table 1). HSPV003/HSPV205 is an intact homologue of the secreted CPXV Brighton Red CrmB TNFR II-like protein (CPXV GRI-90 D2L/I2R), a protein which interacts with and inhibits TNF and lymphotoxin alpha and whose orthologue in VARV has been recently shown to contain a novel carboxyl-terminal chemokine binding domain also present and active in several other OPV proteins (4, 7, 42). HSPV004/HSPV204 encodes an intact homologue of the ankyrin repeat protein encoded by CPXV GRI D3L/I3R and intact homologues in MPXV, ECTV, and CMLV (Fig. 1).

HSPV contains approximately 17 kbp of sequence in three distinct genomic regions (positions 7527 to 18195 in the left terminal region and 194379 to 195517 and 198775 to 204285 in the right terminal region) absent in known VACV-like viruses but homologous to sequences in sequenced strains of CPXV and other OPVs (Fig. 1). HSPV also contains approximately 1.4 kbp of sequence absent not only in VACV but also in all known OPVs except CPXV. For this region, located between positions 15453 and 16985, only MPXV contains a fragment (approximately 75 bp) of homologous sequence. Notably, sequences near this region reflect ITR and/or terminal translocations in several OPVs (Fig. 1), and repetitive sequence near this locus in ECTV has been suggested to be a dynamic genomic region (21). Conceivably, the presence of this 1.4-kbp sequence in HSPV is consistent with retention of adjacent and relatively significant amounts of CPXV-like sequence in this left terminal region relative to other OPVs (Fig. 1).

HSPV sequence in the left terminal region absent in other VACV-like viruses corresponds to the D7L loci of CPXV GRI-90 and the CPXV014 to CPXV020 region of CPXV BRI (Fig. 1A). These sequences relative to other VACV-like viruses essentially extend from the ITR boundary region to the region upstream of the HSPV016 viral growth factor homologue (CPN C11R), replacing the OPV-like sequence that is transposed from the right terminal region to the left terminal region in other VACV-like viruses. HSPV sequences in this region include 15 ORFs representing three intact OPV genes (HSPV008, HSPV010, and HSPV012) and six potentially truncated or fragmented genes (HSPV007, HSPV009, HSPV011a to -c, HSPV013, HSPV014a to -d, and HSPV015a and -b) (Table 1; Fig. 1). HSPV008 encodes an intact protein orthologous only to CPXV GRI-90 D7L and ECTV strain Moscow EVM004 (21, 79). These proteins contain amino-terminal BTB/POZ domains, evolutionarily conserved domains important for oligomerization and ordering of protein complexes and often present in amino-terminal regions of both cellular and poxviral kelch-like proteins, but in these smaller HSPV008 orthologues the BTB/POZ domain is not associated with kelch repeat domains (3, 75). HSPV009 encodes a truncated orthologue of CPXV GRI-90 D12L product, a protein similar to the CrmB carboxyl terminus and whose orthologue in ECTV was recently characterized as a secreted chemokine binding protein (7). HSPV010 encodes an intact orthologue of CD30 TNFR-like proteins present in CPXV and ECTV, proteins able to bind CD30 ligands and/or have immunomodulatory effects (63, 72). HSPV left-end sequences also contain genes for three ankyrin repeat proteins absent in VACV. While HSPV012 encodes an intact ankyrin repeat protein also intact in CPXV and MPXV, HSPV011a to -c and HSPV014a to -d encode fragments of intact ankyrin repeat proteins encoded only in ECTV and/or CPXV, with HSPV014b and -c encoded within the region containing 1.4 kbp of sequence found only in CPXV. Finally, HSPV015a and -b appeared to encode fragments of a paralogue of CPN C7L, a VACV host range protein which enables viral replication in human cells (67). While all OPVs appear to encode intact C7L orthologues (HSPV024), intact HSPV015 orthologues are encoded only in CPXV and CMLV, with fragmented ORFs annotated in MPXV and VARV (Table 1).

HSPV sequence in the right terminal region absent in other

VACV-like viruses essentially bound the region homologous to the VACV WR SPI-1 (HSPV198) locus, a region transposed to the opposite terminus in several other VACVs (Fig. 1). Unique sequence upstream of HSPV198 includes HSPV197, an intact kelch-like protein also intact in CPXV and ECTV but fragmented or absent in MPXV, CMLV, and VARV. Unique sequences downstream of HSPV198 contain an intact orthologue of the VARV strain Bangladesh B22R gene (HSPV200). B22R homologues represent the largest poxviral genes, encoding proteins of approximately 2,000 amino acids and with no known function but predicted to contain carboxyl-terminal transmembrane domains and cysteine residues which conceivably mediate disulfide bond formation (54, 56, 76). B22R homologues are intact in all OPV species except VACV-like viruses, making the presence of HSPV200 notable (Fig. 1).

Despite containing additional sequence not present in other VACV-like viruses, HSPV did lack sequences homologous to several larger regions in other OPVs. These include from GRI-90 the D8L to D11L locus, a region encoding ankyrin repeat, kelch-like, and lectin-like proteins with homologous sequence only in ECTV (79) (Table 1), and most of the K1R to S1R/T1R locus, a region encoding ankyrin repeat, CrmD TNFR, and CrmE TNFR proteins and with homologous sequence present in MPXV, ECTV, and CMLV and, notably, in VACV Lister (Fig. 1C). HSPV also lacks any remnant of the second VARV B22R-like gene identified in certain strains of CPXV and of which remnants remain in VARV and CMLV lineages (HSPV185-HSPV186 locus [Fig. 1C]) (56).

Finally, HSPV contains fragmented genes intact in all or nearly all other VACV-like viruses. Within the central conserved region, HSPV074a to -d represented fragments of the CPN I4L ribonucleotide reductase large subunit gene, while HSPV044 encoded an intact small subunit (Table 2). Ribonucleotide reductase is a heterodimeric protein involved in redox reactions that are key to synthesis of deoxyribonucleotides, an activity for which various poxviruses encode different enzyme complements, potentially adapted to replication in specific host cell types lacking adequate nucleotide pools (59). Experimental disruption of the VACV ribonucleotide reductase large subunit has been shown previously to have no effect on virus replication *in vitro* and a mild effect on virulence in mice (23). Although I4L homologues are not encoded in all other poxviral genera, to our knowledge this is the first example of its natural disruption in an OPV genome. Similarly, HSPV183 is unique among VACV-like homologues (CPN B6R) as the only form of the gene to be fragmented, although a fragmented form is also found in VARV (Fig. 1B) and an isolate of MPXV (accession no. AAY97373). Notably, HSPV contained fragmented genes intact in all VACVs except MVA, a virus that has accumulated numerous mutations and extensive nucleotide deletions through extensive passage *in vitro* and concomitant attenuation and restriction of host range (9). These include HSPV026, orthologue of CPN C5L BTB domain protein, and the HSPV033 ankyrin repeat protein. In addition, HSPV178, similar to MVA, demonstrates a smaller fragmented form of the guanylate kinase gene than do other VACV-like viruses.

Perspective on relationship of HSPV to VACV. Genomic sequence analysis of HSPV MNR-76 indicates that it is a novel VACV-like OPV that contains unique features not present in known VACVs. Although MNR-76 is unique in the comple-

ment of OPV genes remaining intact in HSPV, the pattern of terminal gene loss/fragmentation is commensurate with genotypes observed in other VACV-like viruses. Notably, the majority of left terminal HSPV sequence absent in VACV appears to contain gene fragments, with HSPV conceivably in the process of losing this sequence similarly to other VACV-like viruses.

The close phylogenetic and genotypic relationship between HSPV and other VACV-like viruses and the presence of additional CPXV-like sequences in HSPV are notable given previous speculations involving horsepox and the origins of VACV (14). While the origins of current VACV-like strains have been heavily debated and remain obscure, current knowledge affirms that VACV-like viruses constitute an OPV lineage independent of known CPXV and VARV species from which VACV has been speculated to be derived (14, 32, 33, 38) (Fig. 2). It is likely that a once naturally circulating but now rare VACV-like virus(s) from which current strains are derived was introduced as a vaccine virus, and the agent of horsepox has been surmised as a likely candidate (14). Indeed, apparently Edward Jenner believed that his vaccine originated from the "grease" infection found in the heels of horses, and the use of horse-derived material for use as vaccines is documented (14, 33). In addition, phenotypic similarity of certain vaccines transmitted between cows, humans, and horses has been noted, and experimental infection of horses with VACV can produce clinical signs of horsepox (14, 44, 86). The data presented here indicate that the HSPV MNR-76 genome contains features consistent with such a hypothesis, a phylogenetically VACV-like virus isolated from a horse and containing additional OPV-like terminal sequences, sequences likely ancestral and absent in other VACV-like viruses yet in certain regions appearing to be undergoing gene fragmentation and loss commensurate with transition toward a VACV-like genotype.

Despite speculation as to what role horsepox played in the development of smallpox vaccines, it is clear that HSPV MNR-76 does not represent a direct ancestral genotype to all known VACVs, given the disruption of many HSPV genes intact in certain VACV isolates (Table 1). It is unclear what constitutes the genotypic diversity of all the viruses historically used for smallpox vaccine, especially considering the potential for disparate source material and passage histories of VACV-like vaccine viruses (14, 33). Indeed, phenotypic and genotypic diversity is observed between and within strains of VACV (14, 33, 58) (Fig. 1). This diversity does include sequence unique to a given strain, such as the presence of CPXV GRI K3R and S1R/T1R-like genes in the historically important Lister vaccine strain (Fig. 1C), making the presence of HSPV MNR-76-like sequences in uncharacterized vaccine strains a possibility. Isolated in 1976, HSPV was causing disease in horses while smallpox vaccines were still being distributed during the World Health Organization global smallpox eradication program (32). Conceivably, local or a currently uncharacterized vaccine could have been introduced into the horse population, as contact with vaccinated persons is known to have been a source of OPV disease in animals (33). Vaccine escape has been hypothesized to account for other VACV-like viruses occasionally isolated from domestic and sentinel animals, including RPXV, buffalopox in India, and viruses associated with zoonosis in South America; however, unique biological properties and/or

inability to associate the isolate with vaccine virus has also led to suggestions that they are natural VACV isolates or VACV subspecies (19, 24, 25, 27, 33, 46, 90). Similarly, HSPV MNR-76 may represent a novel, naturally circulating virus and perhaps one for which the horse was an incidental host, just as other domestic and captive animals are not thought to be the reservoir for CPXV infection despite being susceptible to infection (13, 33). Unfortunately, little is known of the prevalence of disease associated with HSPV MNR-76 in Mongolia, either in horse or in human populations. Conceivably, MNR-76 may represent a naturally circulating member of the VACV lineage, as were viruses circulating among domestic animals in the era in which current VACV-like viruses were collected as vaccine. Whatever the historical relationship between HSPV MNR-76 and characterized VACV-like viruses may be, genomic sequence analysis of other VACV-like virus isolates may add perspective to the novel nature of HSPV relative to other viruses within the VACV lineage.

ACKNOWLEDGMENTS

We thank A. Lakowitz and A. Waite Lund for excellent technical assistance.

ADDENDUM IN PROOF

Since completion of the analyses presented here, the genome sequences of several VACV clones derived from the Dryvax vaccine have become available. Preliminary analysis indicates that while most of the HSPV sequence reported here as absent in VACV was also absent in these clones, one (GenBank accession no. AY313848) contained nucleotide sequence and ORF fragments at the HSPV 197 locus, stressing the need for additional genomic sequence and analyses in examining the nature of VACV-like virus variability.

REFERENCES

- Afonso, C. L., E. R. Tulman, Z. Lu, E. Oma, G. F. Kutish, and D. L. Rock. 1999. The genome of *Melanoplus sanguinipes* entomopoxvirus. *J. Virol.* **73**:533–552.
- Afonso, C. L., E. R. Tulman, Z. Lu, L. Zsak, N. T. Sandybaev, U. Z. Kerembekova, V. L. Zaitsev, G. F. Kutish, and D. L. Rock. 2002. The genome of camelpox virus. *Virology* **295**:1–9.
- Albagli, O., P. Dhordain, C. Dewindt, G. Lecocq, and D. Leprince. 1995. The BTB/POZ domain: a new protein-protein interaction motif common to DNA- and actin-binding proteins. *Cell Growth Differ.* **6**:1193–1198.
- Alcami, A. 2003. Viral mimicry of cytokines, chemokines and their receptors. *Nat. Rev. Immunol.* **3**:36–50.
- Alcami, A., A. Khanna, N. L. Paul, and G. L. Smith. 1999. Vaccinia virus strains Lister, USSR and Evans express soluble and cell-surface tumour necrosis factor receptors. *J. Gen. Virol.* **80**:949–959.
- Alcami, A., J. A. Symons, P. D. Collins, T. J. Williams, and G. L. Smith. 1998. Blockade of chemokine activity by a soluble chemokine binding protein from vaccinia virus. *J. Immunol.* **160**:624–633.
- Alejo, A., M. B. Ruiz-Arguello, Y. Ho, V. P. Smith, M. Saraiva, and A. Alcami. 2006. A chemokine-binding domain in the tumor necrosis factor receptor from variola (smallpox) virus. *Proc. Natl. Acad. Sci. USA* **103**:5995–6000.
- Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Antoine, G., F. Scheiffinger, F. Dorner, and F. G. Falkner. 1998. The complete genomic sequence of the modified vaccinia Ankara strain: comparison with other orthopoxviruses. *Virology* **244**:365–396.
- Baroudy, B. M., and B. Moss. 1982. Sequence homologies of diverse length tandem repetitions near ends of vaccinia virus genome suggest unequal crossing over. *Nucleic Acids Res.* **10**:5673–5679.
- Baroudy, B. M., S. Venkatensan, and B. Moss. 1982. Incompletely base-paired flip-flop terminal loops link the two DNA strands of the vaccinia virus genome into one uninterrupted polynucleotide chain. *Cell* **28**:315–324.

12. Barry, M., S. Hnatiuk, K. Mossman, S. F. Lee, L. Boshkov, and G. McFadden. 1997. The myxoma virus M-T4 gene encodes a novel RDEL-containing protein that is retained within the endoplasmic reticulum and is important for the productive infection of lymphocytes. *Virology* **239**:360–377.
13. Baxby, D. 1977. Is cowpox misnamed? A review of 10 human cases. *Br. Med. J.* **1**:1379–1381.
14. Baxby, D. 1981. Jenner's smallpox vaccine: the riddle of vaccinia virus and its origin. Heinemann Educational Books Ltd., London, United Kingdom.
15. Brady, G., L. Boggan, A. Bowie, and L. A. O'Neill. 2005. Schlafen-1 causes a cell cycle arrest by inhibiting induction of cyclin D1. *J. Biol. Chem.* **280**:30723–30734.
16. Breman, J. G., and D. A. Henderson. 2002. Diagnosis and management of smallpox. *N. Engl. J. Med.* **346**:1300–1308.
17. Brudno, M., M. Chapman, B. Gottgens, S. Batzoglu, and B. Morgenstern. 2003. Fast and sensitive multiple alignment of large genomic sequences. *BMC Bioinformatics* **4**:66.
18. Brudno, M., C. B. Do, G. M. Cooper, M. F. Kim, E. Davydov, E. D. Green, A. Sidow, and S. Batzoglu. 2003. LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic DNA. *Genome Res.* **13**:721–731.
19. Buller, R. M., and G. J. Palumbo. 1991. Poxvirus pathogenesis. *Microbiol. Rev.* **55**:80–122.
20. Castresana, J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol. Biol. Evol.* **17**:540–552.
21. Chen, N., M. I. Danila, Z. Feng, R. M. Buller, C. Wang, X. Han, E. J. Lefkowitz, and C. Upton. 2003. The genomic sequence of ectromelia virus, the causative agent of mousepox. *Virology* **317**:165–186.
22. Chen, N., G. Li, M. K. Liszewski, J. P. Atkinson, P. B. Jahrling, Z. Feng, J. Schriever, C. Buck, C. Wang, E. J. Lefkowitz, J. J. Esposito, T. Harms, I. K. Damon, R. L. Roper, C. Upton, and R. M. Buller. 2005. Virulence differences between monkeypox virus isolates from West Africa and the Congo basin. *Virology* **340**:46–63.
23. Child, S. J., G. J. Palumbo, R. M. Buller, and D. E. Hraby. 1990. Insertional inactivation of the large subunit of ribonucleotide reductase encoded by vaccinia virus is associated with reduced virulence in vivo. *Virology* **174**:625–629.
24. da Fonseca, F. G., G. S. Trindade, R. L. Silva, C. A. Bonjardim, P. C. Ferreira, and E. G. Kroon. 2002. Characterization of a vaccinia-like virus isolated in a Brazilian forest. *J. Gen. Virol.* **83**:223–228.
25. Damaso, C. R., J. J. Esposito, R. C. Condit, and N. Moussatche. 2000. An emergent poxvirus from humans and cattle in Rio de Janeiro State: Cantagalo virus may derive from Brazilian smallpox vaccine. *Virology* **277**:439–449.
26. de Carlos, A., and E. Paez. 1991. Isolation and characterization of mutants of vaccinia virus with a modified 94-kDa inclusion protein. *Virology* **185**:768–778.
27. de Souza Trindade, G., F. G. da Fonseca, J. T. Marques, M. L. Nogueira, L. C. Mendes, A. S. Borges, J. R. Peiro, E. M. Pituco, C. A. Bonjardim, P. C. Ferreira, and E. G. Kroon. 2003. Aracatuba virus: a vaccinia-like virus associated with infection in humans and cattle. *Emerg. Infect. Dis.* **9**:155–160.
28. Essajee, S., and H. L. Kaufman. 2004. Poxvirus vaccines for cancer and HIV therapy. *Expert Opin. Biol. Ther.* **4**:575–588.
29. Ewing, B., and P. Green. 1998. Base-calling of automated sequencer traces using *Phred*. II. Error probabilities. *Genome Res.* **8**:186–194.
30. Ewing, B., L. Hillier, M. C. Wendl, and P. Green. 1998. Base-calling of automated sequencer traces using *Phred*. I. Accuracy assessment. *Genome Res.* **8**:175–185.
31. Felsenstein, J. 1989. PHYLIP—phylogeny inference package (version 3.2). *Cladistics* **5**:164–166.
32. Fenner, F., D. A. Henderson, I. Arita, Z. Jezek, and I. D. Ladnyi. 1988. Smallpox and its eradication. World Health Organization, Geneva, Switzerland.
33. Fenner, F., R. Wittek, and K. Dumbell. 1989. The orthopoxviruses. Academic Press, Inc., San Diego, Calif.
34. Galtier, N., M. Gouy, and C. Gautier. 1996. SEAVIEW and PHYLO WIN: two graphic tools for sequence alignment and molecular phylogeny. *Comput. Appl. Biol. Sci.* **12**:543–554.
35. Gardner, J. D., D. C. Tschärke, P. C. Reading, and G. L. Smith. 2001. Vaccinia virus semaphorin A39R is a 50–55 kDa secreted glycoprotein that affects the outcome of infection in a murine intradermal model. *J. Gen. Virol.* **82**:2083–2093.
36. Goebel, S. J., G. P. Johnson, M. E. Perkus, S. W. Davis, J. P. Winslow, and E. Paoletti. 1990. The complete DNA sequence of vaccinia virus. *Virology* **179**:247–266.
37. Gordon, D., C. Abajian, and P. Green. 1998. *Consed*: a graphical tool for sequence finishing. *Genome Res.* **8**:192–202.
38. Gubser, C., S. Hue, P. Kellam, and G. L. Smith. 2004. Poxvirus genomes: a phylogenetic analysis. *J. Gen. Virol.* **85**:105–117.
39. Gubser, C., and G. L. Smith. 2002. The sequence of camelpox virus shows it is most closely related to variola virus, the cause of smallpox. *J. Gen. Virol.* **83**:855–872.
40. Guindon, S., and O. Gascuel. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* **52**:696–704.
41. Hsiao, J. C., C. S. Chung, R. Drillien, and W. Chang. 2004. The cowpox virus host range gene, CP77, affects phosphorylation of eIF2 alpha and vaccinia viral translation in apoptotic HeLa cells. *Virology* **329**:199–212.
42. Hu, F. Q., C. A. Smith, and D. J. Pickup. 1994. Cowpox virus contains two copies of an early gene encoding a soluble secreted form of the type II TNF receptor. *Virology* **204**:343–356.
43. Huang, X., and A. Madan. 1999. CAP3: a DNA sequence assembly program. *Genome Res.* **9**:868–877.
44. Hutya, F., J. Marek, and R. Manninger. 1949. Pox (variola), p. 348–393. *In* J. R. Greig (ed.), *Special pathology and therapeutics of the diseases of domestic animals*, vol. 1. Bailliere, Tindall and Cox, London, United Kingdom.
45. Kochneva, G., I. Kolosova, T. Maksyutova, E. Ryabchikova, and S. Shchelkunov. 2005. Effects of deletions of kelch-like genes on cowpox virus biological properties. *Arch. Virol.* **150**:1857–1870.
46. Kolhapure, R. M., R. P. Deolankar, C. D. Tupe, C. G. Raut, A. Basu, B. M. Dama, S. D. Pawar, M. V. Joshi, V. S. Padbidri, M. K. Goverdhan, and K. Banerjee. 1997. Investigation of buffalopox outbreaks in Maharashtra State during 1992–1996. *Indian J. Med. Res.* **106**:441–446.
47. Kotwal, G. J., and B. Moss. 1988. Analysis of a large cluster of nonessential genes deleted from a vaccinia virus terminal transposition mutant. *Virology* **167**:524–537.
48. Legrand, F. A., P. H. Verardi, L. A. Jones, K. S. Chan, Y. Peng, and T. D. Yilma. 2004. Induction of potent humoral and cell-mediated immune responses by attenuated vaccinia virus vectors with deleted serpin genes. *J. Virol.* **78**:2770–2779.
49. Leite, J. A., B. P. Drummond, G. S. Trindade, Z. I. Lobato, F. G. da Fonseca, S. J. Dos, M. C. Madureira, M. I. Guedes, J. M. Ferreira, C. A. Bonjardim, P. C. Ferreira, and E. G. Kroon. 2005. Passatempo virus, a vaccinia virus strain, Brazil. *Emerg. Infect. Dis.* **11**:1935–1938.
50. Lewis-Jones, S. 2004. Zoonotic poxvirus infections in humans. *Curr. Opin. Infect. Dis.* **17**:81–89.
51. Li, G., N. Chen, R. L. Roper, Z. Feng, A. Hunter, M. Danila, E. J. Lefkowitz, R. M. Buller, and C. Upton. 2005. Complete coding sequences of the rabbitpox virus genome. *J. Gen. Virol.* **86**:2969–2977.
52. Likos, A. M., S. A. Sammons, V. A. Olson, A. M. Frace, Y. Li, M. Olsen-Rasmussen, W. Davidson, R. Galloway, M. L. Khristova, M. G. Reynolds, H. Zhao, D. S. Carroll, A. Curns, P. Formenty, J. J. Esposito, R. L. Regnery, and I. K. Damon. 2005. A tale of two clades: monkeypox viruses. *J. Gen. Virol.* **86**:2661–2672.
53. Massung, R. F., J. C. Knight, and J. J. Esposito. 1995. Topography of variola smallpox virus inverted terminal repeats. *Virology* **211**:350–355.
54. Massung, R. F., L.-I. Liu, J. Qi, J. C. Knight, T. E. Yuran, A. R. Kerlavage, J. M. Parsons, J. C. Venter, and J. J. Esposito. 1994. Analysis of the complete genome of smallpox variola major virus strain Bangladesh-1975. *Virology* **201**:215–240.
55. Merchlinsky, M., and B. Moss. 1989. Nucleotide sequence required for resolution of the concatemeric junction of vaccinia virus DNA. *J. Virol.* **63**:4354–4361.
56. Meyer, H., A. Totmenin, E. Gavrilova, and S. Shchelkunov. 2005. Variola and camelpox virus-specific sequences are part of a single large open reading frame identified in two German cowpox virus strains. *Virus Res.* **108**:39–43.
57. Morgenstern, B., K. Frech, A. Dress, and T. Werner. 1998. DIALIGN: finding local similarities by multiple sequence alignment. *Bioinformatics* **14**:290–294.
58. Morikawa, S., T. Sakiyama, H. Hasegawa, M. Saijo, A. Maeda, I. Kurane, G. Maeno, J. Kimura, C. Hirama, T. Yoshida, Y. Asahi-Ozaki, T. Sata, T. Kurata, and A. Kojima. 2005. An attenuated LC16m8 smallpox vaccine: analysis of full-genome sequence and induction of immune protection. *J. Virol.* **79**:11873–11891.
59. Moss, B. 2001. *Poxviridae*: the viruses and their replication, p. 2849–2883. *In* D. M. Knipe and P. M. Howley (ed.), *Fields virology*, 4th ed., vol. 2. Lippincott Williams & Wilkins, Philadelphia, Pa.
60. Moss, B., and J. L. Shisler. 2001. Immunology 101 at poxvirus U: immune evasion genes. *Semin. Immunol.* **13**:59–66.
61. Moyer, R. W., and R. L. Graves. 1981. The mechanism of cytoplasmic orthopoxvirus DNA replication. *Cell* **27**:391–401.
62. Nylander, J. A. A. 2004. MrModeltest 2.2. Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden.
63. Panus, J. F., C. A. Smith, C. A. Ray, T. D. Smith, D. D. Patel, and D. J. Pickup. 2002. Cowpox virus encodes a fifth member of the tumor necrosis factor receptor family: a soluble, secreted CD30 homologue. *Proc. Natl. Acad. Sci. USA* **99**:8348–8353.
64. Pearson, W. R. 1990. Rapid and sensitive sequence comparison with FASTP and FASTA. *Methods Enzymol.* **183**:63–98.
65. Pearson, W. R., and D. J. Lipman. 1988. Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* **85**:2444–2448.
66. Pearson, W. R., T. Wood, Z. Zhang, and W. Miller. 1997. Comparison of DNA sequences with protein sequences. *Genomics* **46**:24–36.
67. Perkus, M. E., S. J. Goebel, S. W. Davis, G. P. Johnson, K. Limbach, E. K.

- Norton, and E. Paoletti. 1990. Vaccinia virus host range genes. *Virology* **179**:276–286.
68. Price, N., D. C. Tschärke, and G. L. Smith. 2002. The vaccinia virus B9R protein is a 6 kDa intracellular protein that is non-essential for virus replication and virulence. *J. Gen. Virol.* **83**:873–878.
69. Reed, K. D., J. W. Melski, M. B. Graham, R. L. Regnery, M. J. Sotir, M. V. Wegner, J. J. Kazmierczak, E. J. Stratman, Y. Li, J. A. Fairley, G. R. Swain, V. A. Olson, E. K. Sargent, S. C. Kehl, M. A. Frace, R. Kline, S. L. Foldy, J. P. Davis, and I. K. Damon. 2004. The detection of monkeypox in humans in the Western Hemisphere. *N. Engl. J. Med.* **350**:342–350.
70. Rice, P., I. Longden, and A. Bleasby. 2000. EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet.* **16**:276–277.
71. Salzberg, S. L., A. L. Delcher, S. Kasif, and O. White. 1998. Microbial gene identification using interpolated Markov models. *Nucleic Acids Res.* **26**:544–548.
72. Saraiva, M., P. Smith, P. G. Fallon, and A. Alcami. 2002. Inhibition of type 1 cytokine-mediated inflammation by a soluble CD30 homologue encoded by ectromelia (mousepox) virus. *J. Exp. Med.* **196**:829–839.
73. Schmidt, H. A., K. Strimmer, M. Vingron, and A. von Haeseler. 2002. TREE-PUZZLE: maximum likelihood phylogenetic analysis using quartets and parallel computing. *Bioinformatics* **18**:502–504.
74. Senkevich, T. G., E. J. Wolfe, and R. M. L. Buller. 1995. Ectromelia virus RING finger protein is localized in virus factories and is required for virus replication in macrophages. *J. Virol.* **69**:4103–4111.
75. Shchelkunov, S., A. Totmenin, and I. Kolosova. 2002. Species-specific differences in organization of orthopoxvirus kelch-like proteins. *Virus Genes* **24**:157–162.
76. Shchelkunov, S. N., V. M. Blinov, S. M. Resenchuk, A. V. Totmenin, L. V. Olenina, G. B. Chirikova, and L. S. Sandakhchiev. 1994. Analysis of the nucleotide sequence of 53 kbp from the right terminus of the genome of variola major virus strain India-1967. *Virus Res.* **34**:207–236.
77. Shchelkunov, S. N., V. M. Blinov, and L. S. Sandakhchiev. 1993. Ankyrin-like proteins of variola and vaccinia viruses. *FEBS Lett.* **319**:163–165.
78. Shchelkunov, S. N., R. F. Massung, and J. J. Esposito. 1995. Comparison of the genome DNA sequences of Bangladesh-1975 and India-1967 variola viruses. *Virus Res.* **36**:107–118.
79. Shchelkunov, S. N., P. F. Safronov, A. V. Totmenin, N. A. Petrov, O. I. Ryazankina, V. V. Gutorov, and G. J. Kotwal. 1998. The genome sequence analysis of the left and right species-specific terminal region of a cowpox virus strain reveals unique sequences and a cluster of intact ORFs for immunomodulatory and host range proteins. *Virology* **243**:432–460.
80. Shchelkunov, S. N., A. V. Totmenin, V. N. Loparev, P. F. Safronov, V. V. Gutorov, V. E. Chizhikov, J. C. Knight, J. M. Parsons, R. F. Massung, and J. J. Esposito. 2000. Alastrim smallpox variola minor virus genome DNA sequences. *Virology* **266**:361–386.
81. Shchelkunov, S. N., A. V. Totmenin, P. F. Safronov, M. V. Mikheev, V. V. Gutorov, O. I. Ryazankina, N. A. Petrov, I. V. Babkin, E. A. Uvarova, L. S. Sandakhchiev, J. R. Sisler, J. J. Esposito, I. K. Damon, P. B. Jahrling, and B. Moss. 2002. Analysis of the monkeypox virus genome. *Virology* **297**:172–194.
82. Shisler, J. L., and B. Moss. 2001. Immunology 102 at poxvirus U: avoiding apoptosis. *Semin. Immunol.* **13**:67–72.
83. Smith, G. L., and G. McFadden. 2002. Smallpox: anything to declare? *Nat. Rev. Immunol.* **2**:521–527.
84. Sonhammer, E. L., and R. Durbin. 1995. A dot-matrix program with dynamic threshold control suited for genomic DNA and protein sequence analysis. *Gene* **167**:GC1–GC10.
85. Spehner, D., S. Gillard, R. Drillien, and A. Kirn. 1988. A cowpox virus gene required for multiplication in Chinese hamster ovary cells. *J. Virol.* **62**:1297–1304.
86. Studdert, M. J. 1989. Experimental vaccinia virus infection of horses. *Aust. Vet. J.* **66**:157–159.
87. Taylor, J. M., and M. Barry. 2006. Near death experiences: poxvirus regulation of apoptotic death. *Virology* **344**:139–150.
88. Thompson, C. H., J. A. Yager, and I. B. Van Rensburg. 1998. Close relationship between equine and human molluscum contagiosum virus demonstrated by in situ hybridisation. *Res. Vet. Sci.* **64**:157–161.
89. Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
90. Trindade, G. S., F. G. da Fonseca, J. T. Marques, S. Diniz, J. A. Leite, S. De Bodt, Y. Van der Peer, C. A. Bonjardim, P. C. Ferreira, and E. G. Kroon. 2004. Belo Horizonte virus: a vaccinia-like virus lacking the A-type inclusion body gene isolated from infected mice. *J. Gen. Virol.* **85**:2015–2021.
91. Upton, C., L. Schiff, S. A. Rice, T. Dowdeswell, X. Yang, and G. McFadden. 1994. A poxvirus protein with a RING finger motif binds zinc and localizes in virus factories. *J. Virol.* **68**:4186–4195.
92. Upton, C., S. Slack, A. L. Hunter, A. Ehlers, and R. L. Roper. 2003. Poxvirus orthologous clusters: toward defining the minimum essential poxvirus genome. *J. Virol.* **77**:7590–7600.
93. Wesley, R. D., and A. E. Tuthill. 1984. Genome relatedness among African swine fever virus field isolates by restriction endonuclease analysis. *Prev. Vet. Med.* **2**:53–62.
94. Yager, J. A., D. W. Scott, and B. P. Wilcock. 1993. The skin and appendages, p. 531–738. *In* K. V. F. Jubb, P. C. Kennedy, and N. Palmer (ed.), *Pathology of domestic animals*, 4th ed., vol. 1. Academic Press, San Diego, Calif.