

5-1-2013

Getting Hooked on Farmed Salmon: A Good Source of Omega-3s

Susan Raatz

USDA-ARS, susan.raatz@ars.usda.gov

Matthew Picklo

USDA-ARS, matthew.picklo@ars.usda.gov


William Wolters

USDA-ARS, bill.wolters@ars.usda.gov

Rosalie Marion Bliss

USDA-ARS

Follow this and additional works at: <http://digitalcommons.unl.edu/usdaagresmag>

 Part of the [Agriculture Commons](#), [Animal Sciences Commons](#), [Food Science Commons](#), and the [Plant Sciences Commons](#)

Raatz, Susan; Picklo, Matthew; Wolters, William; and Bliss, Rosalie Marion, "Getting Hooked on Farmed Salmon: A Good Source of Omega-3s" (2013). *Agricultural Research Magazine*. Paper 61.

<http://digitalcommons.unl.edu/usdaagresmag/61>

This Article is brought to you for free and open access by the USDA Agricultural Research Service --Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agricultural Research Magazine by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

GETTING HOOKED ON FARMED SALMON

A Good Source of Omega-3s

Nearly 85 percent of seafood enjoyed by U.S. consumers is imported, and almost half of that is farm raised through a process called “aquaculture,” according to the National Aquaculture Association, based in Pine Bluff, Arkansas. In 2011, the United States imported \$16.6 billion of seafood and exported \$5.4 billion, a difference of \$11.2 billion.

As the world’s population grows and demand for seafood increases, many seafood species are overfished or fully exploited. That means the world’s oceans cannot turn out more fish than are currently being produced. As the ceiling is being reached on seafood availability from the wild, U.S. producers of farm-raised seafood are working hard to help fill today’s growing demand for seafood in a sustainable way. Aquaculture is a process for raising aquatic species—both marine and freshwater—in a captive environment under controlled conditions.

Salmon: A Great Catch for Nutrition

Agricultural Research Service scientists at the Grand Forks [North Dakota] Human Nutrition Research Center (GFHNRC) conducted studies involving one popular U.S. farmed fish, Atlantic salmon. They wanted to learn more about the omega-3 fatty acids in the fish.



TANIA DAVIS (D2866-1)

A salmon fillet (high in omega-3 fatty acids), purple potatoes (loaded with antioxidants), and asparagus.

Two omega-3 fatty acids, EPA (eicosapentaenoic acid) and DHA (docosahexaenoic acid), are abundant in oily fish such as salmon, tuna, mackerel, and herring. Some data has shown that consuming 250 milligrams of EPA and DHA—the amount in less than 3 calories of polyunsaturated

culture Center in Franklin, Maine.

“Atlantic salmon is an excellent source of EPA and DHA,” says Raatz, “and people can easily increase their consumption of omega-3 PUFAs by including more seafood in their diets.” These new studies provide data of interest to those who have

concerns that farm-raised salmon has less available omega-3s than salmon caught in the wild.

Retaining Healthy Omega-3s in Fish

While eating seafood rich in omega-3 fatty acids is known to reduce risk of heart disease, it has not been known whether the baking process causes loss of omega-3s in farm-raised Atlantic salmon. The researchers at Grand Forks demonstrated that baking salmon to the proper temperature does not decrease the content of beneficial omega-3 fatty acids in salmon.

TROUTLODGE, INC. (D1376-1)



Atlantic salmon is one of the major species grown in aquaculture, a process for raising aquatic species in a captive environment under controlled conditions. Recent ARS studies should help alleviate concerns about whether farm-raised salmon have less available omega-3s than salmon caught in the wild.



At the Grand Forks [North Dakota] Human Nutrition Research Center, physiologist Matthew Picklo performs fatty acid analysis of a sample of human blood plasma to determine the effects of salmon consumption.

ounces), and 270 grams (about 9.5 ounces).

The Institute of Medicine (IOM) and the *Dietary Guidelines for Americans* recommend consuming 8 ounces of seafood weekly, but at this time the IOM has not established a recommended dietary allowance for EPA or DHA.

Blood was collected from each of the 19 subjects to mark fatty acid levels and other heart disease risk indicators at the beginning and end of each treatment.

The results showed that EPA blood levels doubled after the volunteers consumed the 6.3-ounce portions and increased nearly threefold after they consumed the 9.5-ounce portions. Also, based on the blood indicators, DHA levels were elevated by about 50 percent, regardless of portion size.

“We found that eating 6 ounces twice weekly was adequate when it comes to raising blood levels of EPA and DHA,” says Raatz. “The volunteers did not have to eat a lot of fish to see the benefit.”

“We showed that consuming farm-raised salmon was an excellent way to increase omega-3 fatty acids in the blood to levels that corresponded to reduced heart disease risk,” says Picklo.

The American Heart Association recommends consuming fish twice a week to reduce risk of heart disease.

At this time, only the U.S. states of Maine and Washington produce farm-raised salmon, says Sebastian Belle, executive director of the Maine Aquaculture Association based in Hallowell, Maine.

“On average during the past 5 years, Maine has produced

about 26 million pounds of farm-raised salmon,” says Belle. Significantly, in 2011, both U.S.-imported wild and farmed Atlantic salmon reached nearly 400 million pounds.

“It’s important to understand the health benefits of Atlantic salmon produced through aquaculture,” says Wolters, who collaborated with GFHNRC on both of these salmon studies. “A lot of the seafood consumers find at local grocery stores is farm raised.”

The study was published in the *Journal of the Academy of Nutrition and Dietetics* in February 2013.—By **Rosalie Marion Bliss, ARS.**

This research is part of Human Nutrition, an ARS national program (#107) described at www.nps.ars.usda.gov.

Susan Raatz and Matthew Picklo are with the USDA-ARS Grand Forks Human Nutrition Research Center, 2420 2nd Ave., Grand Forks, ND 58202; (701) 795-8294 [Raatz], (701) 795-8380 [Picklo], susan.raatz@ars.usda.gov, matthew.picklo@ars.usda.gov.

*William Wolters is with the USDA-ARS National Cold Water Aquaculture Center, 25 Salmon Farm Rd., Franklin, ME 04634; (207) 422-2713, bill.wolters@ars.usda.gov. **

MATTHEW PICKLO (D2867-1)



ARS nutritionist Susan Raatz (right) and cook Doris Zidon prepare farm-raised salmon fillets for a research study at the Grand Forks nutrition center.

The study also examined the extent to which the baking process for Atlantic salmon produces oxidized fatty acids. “Some people think that when you bake fish, it’s possible to oxidize the healthful fatty acids, which would leave compounds that are not good for you, like toxic omega-3 oxidation byproducts,” says Picklo. “We looked at the extent to which baking would alter the fatty acids in salmon.”

The researchers found that baking actually decreases the presence of fatty acid oxidation products. “Preparing the fish based on restaurant and safety guidelines—to a tender-but-safe 145°F—rather than overcooking to a tough, chewy fish was key,” says Raatz.

The 2011 paper was published in the *Journal of Agricultural and Food Chemistry*.

Measuring Blood Levels of Omega-3s

In another study, the scientists evaluated a group of 19 healthy human volunteers who were provided 3 different portion sizes of farm-raised Atlantic salmon. Each volunteer was assigned to consume two weekly servings of one of the three portion sizes of salmon for a 4-week period. After a “blood-clearing” break of 4 to 6 weeks, a different portion size was served, followed by another break, after which the third portion size was served, so that each volunteer had consumed all three portion sizes.

The raw weights of the salmon portions fed to the volunteers were 90 grams (about 3.2 ounces), 180 grams (about 6.3