University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Anthony F. Starace Publications

Research Papers in Physics and Astronomy

11-1-1997

Dispersion coefficients for highly excited molecular states of K₂

Mircea Marinescu University of Nebraska - Lincoln

Anthony F. Starace University of Nebraska-Lincoln, astarace1@unl.edu

Follow this and additional works at: http://digitalcommons.unl.edu/physicsstarace



Part of the Physics Commons

Marinescu, Mircea and Starace, Anthony F., "Dispersion coefficients for highly excited molecular states of K2" (1997). Anthony F. Starace Publications. Paper 62.

http://digitalcommons.unl.edu/physicsstarace/62

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at Digital Commons@University of Nebraska -Lincoln. It has been accepted for inclusion in Anthony F. Starace Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Dispersion coefficients for highly excited molecular states of K₂

Mircea Marinescu and Anthony F. Starace

Department of Physics and Astronomy, The University of Nebraska, Lincoln, Nebraska 68588-0111

(Received 29 May 1997)

We report results of theoretical calculations for the dispersion coefficients associated with the long-range interaction of two 39 K atoms in the states 39 K(4s)- 39 K(5d,6d,7s). These results have been found by Wang *et al.* [Phys. Rev. Lett. **78**, 4173 (1997)] to be in a good agreement with their measurements. [S1050-2947(97)10010-5]

PACS number(s): 34.20.Cf, 31.15.Md

I. INTRODUCTION

Progress in ultracold spectroscopy has made possible experimental investigation of weakly bound highly excited molecular states with an unprecedent level of accuracy. In particular, the first observation of optical-optical doubleresonance photoassociative spectroscopy of ultracold ³⁹K atoms near the highly excited 39 K(4s) + ³⁹K(4d,5d,6d,6s,7s) asymptotes in a high-density magneto-optical trap has recently been reported [1]. Based on the measurements of the higher excited vibrational levels, it was possible to estimate the values of the dominant dispersion coefficients for the long-range interaction of two K atoms with one of them in a highly excited state [1]. Theoretical calculations of the dispersion coefficients for the 39 K(4s)- 39 K(6s,4d) dissociation limits of K₂ have been reported previously [2]. There is good agreement with new experimental data for these states [1]. However, no theoretical predictions exist for the higher excited states measured recently [1]. In this Brief Report we present theoretical calculations of the dispersion coefficients for the $^{39}{\rm K}(4s)$ - $^{39}{\rm K}(7s,5d,6d)$ dissociation limits and compare with the experimental measurements [1]. Throughout this paper we use atomic units ($\hbar = m_e = e = 1$).

II. MATHEMATICAL BACKGROUND

Derivations of the formal expressions for the dispersion coefficients associated with the S-S and S-D dissociation limits were presented in Ref. [2], based on a model potential description of the active valence electron of the alkali-metal atoms [3]. Here we therefore only present the final results needed for our analysis. Briefly, in the long-range limit, the molecular potential interaction may be written as a series of inverse powers of the internuclear distance R. The coefficients of this series are the dispersion coefficients. They may be computed using perturbation theory, where the perturbation is given by the Coulomb interaction between the atomic charge distributions. For the S-S asymptote, where in the dissociation limit one of the atoms is in the ground state $\{n_{\alpha},0,0\}$ and the other in an excited S state $\{n_{\alpha},0,0\}$, the long-range potential interaction is given by the following expression:

$$V^{\beta}(R) = -\frac{C_6^{\beta}}{R^6} - \frac{C_8^{\beta}}{R^8} - \frac{C_{10}^{\beta}}{R^{10}} - \cdots, \tag{2.1}$$

where the coefficients C_n^{β} are the dispersion coefficients, which are obtained from second-order perturbative corrections to the molecular energies. In Eq. (2.1) we have $\beta = p\sigma$, where $p = \pm 1$ for even (g) and odd (u) molecular states, and $\sigma = \pm 1$ for singlet and triplet states. The dispersion coefficients C_6^{β} , C_8^{β} , and C_{10}^{β} are given by the following expressions [2]:

$$C_6^{\beta} = \frac{2}{3} (S_1^0 + \beta S_2^0),$$
 (2.2)

$$C_8^{\beta} = S_3^0 + S_4^0 + 2\beta S_5^0,$$
 (2.3)

$$C_{10}^{\beta} = \frac{14}{5} S_6^0 + \frac{4}{3} (S_7^0 + S_8^0) + \beta (\frac{14}{5} S_9^0 + \frac{8}{3} S_{10}^0),$$
 (2.4)

where S_j^0 for $j=1,\ldots,10$ are sums of different products of atomic radial matrix elements divided by appropriate energy denominators. S_j^0 are given by Eqs. (46), (47), (50)–(52) and (55)–(59) of Ref. [2].² Similarly, for the S-D asymptote, where in the dissociation limit one atom is in the ground state and the other is in an excited D state $\{n_e, 2, m_e\}$, the long-range interaction potential is given by

$$V^{m_e\beta}(R) = -\frac{C_5^{m_e\beta}}{R^5} - \frac{C_6^{m_e\beta}}{R^6} - \cdots, \qquad (2.5)$$

where the C_5 coefficients are obtained in the first order of perturbation and the C_6 coefficients are obtained in the second order of perturbation. They are given by the following expressions:

$$C_5^{m_e\beta} = \frac{(-1)^{m_e+1}\beta}{5} \binom{4}{2+m_e} (n_e 2|r^2|n_g 0)^2, \quad (2.6)$$

¹In general, for dissociation limits where one atom is in the ground state and the other is in an excited state $\{n_e, l_e, m_e\}$, we have $\beta = (-1)^{l_e} p \sigma$.

²Note that there is a misprint in Eq. (56) of Ref. [2]: E_{n_30} in the denominator should be E_{n_20} .

$$C_6^{m_e\beta} = A_1^{m_e} S_1^2 + A_2^{m_e} S_2^2 + \beta A_3^{m_e} S_3^2,$$
 (2.7)

where the algebraic coefficients $A_j^{m_e}$ for j=1,2,3 are given in Table II of Ref. [2], the sums S_j^2 for j=1,2,3 are given by Eqs. (78)–(80) of Ref. [2], and the $\binom{n}{k}$ are the binomial coefficients. The main task in computing the dispersion coefficients consists in evaluating each of the double sums S_1^0,\ldots,S_{10}^0 and S_1^2,S_2^2,S_3^2 . They are sums over the principal quantum numbers of both of the atoms. For example, the sum S_1^0 is given by

$$S_1^0 = \sum_{nm} \frac{(n1|r|n_g0)^2 (m1|r|n_e0)^2}{E_{n_1} + E_{m_1} - E_{n_e0} - E_{n_g0}}.$$
 (2.8)

In general, an efficient method of computation is to separate these sums into independent contributions of each atom by using the following integral representation [4] for each of the denominators in the expressions for the S sums:

$$\frac{1}{a+b} = \frac{2}{\pi} \int_0^\infty d\omega \operatorname{Re} \left[\frac{1}{a-i\omega} \right] \operatorname{Re} \left[\frac{1}{b-i\omega} \right]. \tag{2.9}$$

This integral representation is valid only if a>0 and b>0. By inverting the sums over the principal quantum numbers and the integral over frequency, and formally computing the sums, we obtain an integral over a product of atomic radial matrix elements containing a radial Green's function for complex energy. The main advantage of this procedure is that the two-center molecular problem is reduced to the one-center atomic problem.

An example will serve to illustrate the theoretical procedure just described. Consider again the S_1^0 sum given by Eq. (2.8). For the specific order of the energy levels of K, i.e., 4s, 4p, 5s, 3d, 5p, 4d, 6s, 4f, 6p, 5d, 7s, 5f, 5g, 7p, 6d, etc., it is not always possible to arrange the denominators in the expression of the S sums such that a>0 and b>0, as required in order to use Eq. (2.9). The cases where the condition a>0 and b>0 cannot be satisfied must be treated separately. Thus for the energies in S_1^0 , if we chose $a=E_{n1}-E_{4S}$ and $b=E_{m1}-E_{7S}$, then for $\{m1\}$ $\equiv 4P$, 5P, and 6P we have b<0. In those cases where b is smaller than 0, we replace 1/(a+b) by

$$\frac{1}{a+b} + \frac{1}{a-b} + \frac{2}{\pi} \int_0^\infty d\omega \operatorname{Re} \left[\frac{1}{a-i\omega} \right] \operatorname{Re} \left[\frac{1}{b-i\omega} \right], \tag{2.10}$$

where the last two terms in Eq. (2.10) cancel each other.³ This procedure enables us to obtain a complete sum over the atomic states in the integral. The first two terms resulting from Eq. (2.10) are treated separately. Thus, the final expression for S_1^0 is given by

TABLE I. Dispersion coefficients in a.u. for the $^{39}K(4s)$ - $^{39}K(7s)$ asymptote.

β	$C_6 \times 10^{-5}$	$C_8 \times 10^{-9}$	$C_{10} \times 10^{-12}$
+1	3.9195	1.0217	2.8799
	3.7 ± 0.2^{a}		
-1	3.9226	1.0219	2.8800

^aMeasurement of Wang et al. [1].

$$\begin{split} S_1^0 &= \frac{2}{\pi} \int_0^\infty d\omega \ \text{Re}[(4S|rg_P(E_{4S} + i\omega)r|4S)] \\ &\times \text{Re}[(7S|rg_P(E_{7S} + i\omega)r|7S)] \\ &+ (4P|r|7S)^2[(4S|rg_P(E_{4S} - E_{4P} + E_{7S})r|4S) \\ &+ (4S|rg_P(E_{4S} + E_{4P} - E_{7S})r|4S)] \\ &+ (5P|r|7S)^2[(4S|rg_P(E_{4S} - E_{5P} + E_{7S})r|4S) \\ &+ (4S|rg_P(E_{4S} + E_{5P} - E_{7S})r|4S)] \\ &+ (4S|rg_P(E_{4S} + E_{5P} - E_{7S})r|4S)] \\ &+ (6P|r|7S)^2[(4S|rg_P(E_{4S} - E_{6P} + E_{7S})r|4S) \\ &+ (4S|rg_P(E_{4S} + E_{6P} - E_{7S})r|4S)], \end{split}$$

where g_P is the radial Green's function for angular momentum l=1. The last three terms in Eq. (2.11) stem from the first two terms in Eq. (2.10) for the three cases where b < 0. Alternatively, if we choose $a = E_{m1} - E_{4S}$ and $b = E_{n1} - E_{7S}$, then we obtain

$$S_{1}^{0} = \frac{2}{\pi} \int_{0}^{\infty} d\omega \operatorname{Re}[(4S|rg_{P}(E_{7S} + i\omega)r|4S)] \operatorname{Re}[(7S|rg_{P}(E_{4S} + i\omega)r|7S)] + (4P|r|4S)^{2}[(7S|rg_{P}(E_{4S} - E_{4P} + E_{7S})r|7S) + (7S|rg_{P}(E_{4S} + E_{4P} - E_{7S})r|7S)] + (5P|r|4S)^{2}[(7S|rg_{P}(E_{4S} - E_{5P} + E_{7S})r|7S)] + (7S|rg_{P}(E_{4S} + E_{5P} - E_{7S})r|7S)] + (6P|r|4S)^{2}[(7S|rg_{P}(E_{4S} - E_{6P} + E_{7S})r|7S) + (7S|rg_{P}(E_{4S} + E_{6P} - E_{7S})r|7S)].$$

$$(2.12)$$

Equations (2.11) and (2.12) are two alternative expressions for the same sum S_1^0 . Thus they provide a good numerical test of our codes, which give results using these two equa-

TABLE II. Dispersion coefficients in a.u. for the $^{39}{\rm K}(4s)$ - $^{39}{\rm K}(5d)$ asymptote.

	β	$C_5 \times 10^2$	$C_6 \times 10^{-5}$
m=0	+1	-86.150	4.0148
	-1	86.150	4.0164
$m = \pm 1$	+1	57.433	3.5620
			3.3 ± 0.3^{a}
	-1	-57.433	3.5609
$m = \pm 2$	+1	-1.4358	2.1986
	-1	1.4358	2.1992

^aMeasurement of Wang et al. [1].

³This cancellation follows for b < 0 from the fact that $(a-b)^{-1} = (a+|b|)^{-1}$ and the fact that $\operatorname{Re}(b-i\omega)^{-1} = -\operatorname{Re}(|b|-i\omega)^{-1}$.

TABLE III. Dispersion coefficients in a.u. for the $^{39}\text{K}(4s)$ - $^{39}\text{K}(6d)$ asymptote.

	β	$C_5 \times 10^2$	$C_6 \times 10^{-5}$
m=0	+1	-14.282	9.1057
	-1	14.282	9.1068
$m = \pm 1$	+1	9.5212	8.0870
			10.1 ± 0.7^{a}
	-1	-9.5212	8.0862
$m = \pm 2$	+1	-2.3803	5.0275
	-1	2.3803	5.0279

^aMeasurement of Wang et al. [1].

tions which agree to seven digits of accuracy. In a similar way, each of the sums may be reduced to an integral representation. One then must evaluate only atomic matrix elements containing a radial Green's function. This step may be done easily by using the Dalgarno-Lewis method [5] even in cases involving complex energies [2,3].

III. NUMERICAL RESULTS AND DISCUSSION

The numerical results for the C_6 , C_8 , and C_{10} coefficients corresponding to the $^{39}\mathrm{K}(4s)$ - $^{39}\mathrm{K}(7s)$ asymptote are given in Table I, and those for the $^{39}\mathrm{K}(4s)$ - $^{39}\mathrm{K}(5d,6d)$ asymptotes are given in Tables II and III.⁴ The theoretical computations have an uncertainty of 4% due mainly to the 2% experimental uncertainty of the static dipole polarizability of K [6], whose value our model potential has been constructed to reproduce, following the procedure of Ref. [3]. A discussion of the influence of the uncertainty of the measured static

dipole polarizability on the values of the calculated dispersion coefficients was given in Ref. [7] for the case of longrange interactions among three alkali-metal atoms. A similar discussion also holds for the dispersion coefficients of K dimers presented here. We notice that for the $^{39}\mathrm{K}(4s)$ - $^{39}\mathrm{K}(5d,6d)$ asymptote, the C_5 coefficients have very small values in comparison with the C_6 coefficients. Thus the dominant contributions to the $^{39}\mathrm{K}(4s)$ - $^{39}\mathrm{K}(5d,6d)$ asymptotes for K_2 are given by the $1/R^6$ term in Eq. (2.5) for $R\!<\!10^7$ a.u. Beyond 10^7 a.u. the C_5 coefficients may begin to dominate. However, at such large distances the Casimir effect may become important and hence one expects a smooth transformation of the $1/R^5$ term in Eq. (2.5) to one varying as the $1/R^6$ power.

The experimental values [1] of the dispersion coefficients are obtained by analyzing the spectroscopic data of high vibrational levels. For the $^{39}K(4s)$ - $^{39}K(7s)$ asymptote the measured vibrational spectrum has ${}^{3}\Sigma_{g}^{+}$ symmetry, while for the ${}^{39}\text{K}(4s)$ - ${}^{39}\text{K}(5d,6d)$ asymptotes the vibrational spectrum has 1_u symmetry. The experimental values for the dispersion coefficients C_6 have been determined by using the single leading term potential model (cf. Ref. [8]). As shown in the tables, we find good agreement between theory and experiment for the leading terms of the ${}^{39}K(4s)$ - ${}^{39}K(7s)$ and ${}^{39}K(4s)$ - ${}^{39}K(5d)$ asymptotes. The disagreement between theory and experiment for the case of 39 K(4s)- 39 K(6d) may be the result of neglect of finestructure contributions as well as the exchange interaction and Casimir effect in the long-range form of the potential curve used to extract the dispersion coefficients from the experimental data.

ACKNOWLEDGMENT

This work was supported by the U.S. Department of Energy, Office of Basic Energy Sciences, under Grant No. DE-FG03-96ER14646.

⁴The coefficients C_k have dimensions of energy×(length)^k and are given in the tables in a.u., where the energy unit in a.u. equals 27.211 396 eV and the length unit in a.u. equals 5.291 772 × 10⁻⁹ cm.

^[1] H. Wang, X. T. Wang, P. L. Gould, and W. C. Stwalley, Phys. Rev. Lett. 78, 4173 (1997).

^[2] M. Marinescu and A. Dalgarno, Phys. Rev. A 52, 311 (1995).

^[3] M. Marinescu, H. R. Sadeghpour, and A. Dalgarno, Phys. Rev. A 49, 982 (1994).

^[4] C. Mavroyannis and M. J. Stephen, Mol. Phys. 5, 629 (1962);
A. D. McLachlan, Proc. R. Soc. London, Ser. A 271, 387 (1963); see also the review by A. Dalgarno and W. D. Davi-

son, Adv. At. Mol. Phys. 2, 1 (1966).

^[5] A. Dalgarno and J. T. Lewis, Proc. R. Soc. London, Ser. A 233, 70 (1955).

^[6] W. D. Hall and J. C. Zorn, Phys. Rev. A 10, 1141 (1974).

^[7] M. Marinescu and A. F. Starace, Phys. Rev. A 55, 2067 (1997).

^[8] R. J. LeRoy and R. B. Bernstein, J. Chem. Phys. 52, 3869 (1970).