University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Roman L. Hruska U.S. Meat Animal Research Center

USDA Agricultural Research Service -- Lincoln, Nebraska

1-1-1985

Variation in Sensory Properties of Meat as Affected by Sex Condition, Muscle, and Postmortem Aging

Steven C. Seideman MARC

John D. Crouse **MARC**

Follow this and additional works at: http://digitalcommons.unl.edu/hruskareports



Part of the Animal Sciences Commons

Seideman, Steven C. and Crouse, John D., "Variation in Sensory Properties of Meat as Affected by Sex Condition, Muscle, and Postmortem Aging" (1985). Roman L. Hruska U.S. Meat Animal Research Center. Paper 66. http://digitalcommons.unl.edu/hruskareports/66

This Article is brought to you for free and open access by the USDA Agricultural Research Service -- Lincoln, Nebraska at Digital Commons@University of Nebraska - Lincoln. It has been accepted for inclusion in Roman L. Hruska U.S. Meat Animal Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Variation in Sensory Properties of Meat as Affected by Sex Condition, Muscle, and Postmortem Aging

Steven C. Seideman and John D. Crouse¹

Introduction

For several decades, the sensory properties of beef, particularly tenderness, have been of interest to the meat industry. Variations in sensory properties of beef have been attributed to muscle cut or muscle and postmortem aging. The objective of this study was to examine the sensory properties of five beef muscles, determine the contribution of connective tissue (i.e., collagen) to tenderness, and investigate the response of various muscles to postmortem aging.

Procedure

Eight bulls and eight steers of similar backgrounds were slaughtered. The longissimus dorsi (LD; ribeye), psoas major (PM; tenderloin), semitendinosus (ST; eye of round), semimembranosus (SM; top round), and biceps femoris (BF; bottom round) muscles were removed from the right and left sides of each carcass 24 h postmortem. The muscles from the right sides of all carcasses were immediately frozen (24 h postmortem) while the muscles from the left sides of all carcasses were aged at refrigeration temperatures for 7 days prior to

After freezing, all muscles were cut into steaks. Steaks were used for sensory panel evaluations, shear force determinations, and compositional properties. Sensory panel evaluations were conducted using an 8-member panel. Panelists rated steaks for juiciness, ease of fragmentation, amount of connective tissue, tenderness, and flavor intensity. Cores (1.3 cm) from cooked steaks were also sheared on an Instron Universal testing machine equipped with a Warner-Bratzler shear device. One steak from each carcass and each muscle was powdered in liquid nitrogen and analyzed for percentage fat, amount of collagen (expressed on a wet basis and fat-free basis), and percentage soluble collagen.

Sensory and compositional properties of five bovine muscles are shown in Table 1. The psoas major muscle was the most juicy and most tender of all muscles. Muscles were ranked in order of sensory ratings; PM>ST>LD>BF>SM. The amount of fat (intramuscular) was highest in the PM muscle and lowest in the SM muscle. The amount of collagen on a fat-free basis was ranked LD>ST>BF>PM>SM. The percentage collage solubility was ranked LD>PM>ST>BF>SM.

Simple correlation coefficients between compositional conponents and tenderness and shear force at 1 and 7 days posmortem are presented in Table 2. The percentage of fat within a muscle was significantly correlated to tenderness at 1 and 7 days postmortem and to shear force at 7 days postmorten The amount of collagen was negatively correlated to tende ness and shear force at 1 and 7 days postmortem, whereas the percentage soluble collagen was never significantly corelated to tenderness or shear force.

Mean values for tenderness and shear force stratified by muscle and postmortem aging period are shown in Table 3 Sensory tenderness ratings were not affected by aging period; however, shear force values were lower after 7 days of agiru as compared to samples aged for only 1 day.

The results suggest that muscles vary considerably in tenderness, but neither the amount or the percentage collagen solubility are solely responsible for differences in the tenderness between muscles, although the amount of collagen was, by far, the more closely correlated to tenderness differences. The sensory panel tenderness ratings did not appear to reflect any differences in aged muscle samples; whereas shear force values were substantially lower for muscles aged 7 days, as opposed to muscles aged for only 1 day. Research is continuing to determine why muscles differ in tenderness.

Table 1.—Sensory and compositional properties of five bovine muscles

	Muscle					
Property	Longissimus dorsi	Psoas major	Semitendinosus	Semimembranosus	Biceps femoris	
Sensory properties						
Juiciness ^e	5.3b	5.8a	5.2b	4.9°	5.2b	
Ease of fragmentationf		5.9ª	4.9b	4.5°	4.6°	
Amount of connective						
tissue ^g	4.6°	5.9ª	4.8b	4.4°	4.5°	
Tendernessh	4.8bc	5.9a	4.96	4.7°	4.8bc	
Flavor intensity	5.7d	5.9ª	5.7 ^{cd}	5.8bc	5.9ab	
Shear force (lb)		5.49d	8.83°	9.37bc	10.10b	
Compositional properties						
Fat (pct)	2.67⁵	3.43a	2.27⁵	1.36°	2.170	
Amount of collagen (mg/g)		3.18°	7.53a	5.05b	7.43a	
Collagen solubility (pct) Amount of collagen (mg/g)		16.6b	15.9b	9.1°	14.5b	
fat free	1.54ª	0.56°	1.25ab	0.51°	1.10b	

abod Means on the same line followed by a common superscript are not different.

Results

Seideman is a research food technologist and Crouse is the research leader, Meats Unit, MARC.

Means based on an 8-point scale (8 = extremely juicy; 1 = extremely dry).

Means based on an 8-point scale (8 = extremely easy; 1 = extremely difficult).

Means based on an 8-point scale (8 = extremely easy; 1 = extremely difficult).

hMeans based on an 8-point scale (8 = extremely tender; 1 = extremely tough).

Means based on an 8-point scale (8 = extremely intense; 1 = extremely bland)

Table 2.—Simple correlation coefficients between compositional components and tenderness and shear force at 1 and 7 days postmortem

Compositional components	Tende	erness	Shear force		
	1 day	7 days	1 day	7 days	
Fat (pct)	0.39***	0.59***	-0.19	-0.40***	
Amount of collagen	-0.42***	-0.33**	-0.42***	0.32**	
Collagen solubility	-0.02	0.16	0.08	0.13	

"*P<0.01

Table 3.—Mean values for tenderness and shear force stratified by muscle and postmortem aging period

Tinderness purameter	Postmortem aging period (days)	Muscle Muscle					
		Longissimus dorsi	Psoas major	Semitendinosus	Semimembranosus	Biceps femoris	
Sensory tenderness	1	4.6	5.8	5.1	4.9	4.7	
	7	4.8	5.9	4.9	4.7	4.8	
Shear force (lb)	1	17.5	6.9	9.1	10.6	10.8	
	7	9.6	5.4	8.1	8.3	9.1	