

September 2000

A role for calcineurin in *Dictyostelium discoideum* phagocytosis

Aidong Yuan

University of Nebraska-Lincoln

Catherine P. Chia

University of Nebraska-Lincoln, cchia1@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/bioscimicro>



Part of the [Microbiology Commons](#)

Yuan, Aidong and Chia, Catherine P, "A role for calcineurin in *Dictyostelium discoideum* phagocytosis" (2000). *Papers in Microbiology*. 69.

<http://digitalcommons.unl.edu/bioscimicro/69>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Microbiology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Yuan, A. D. and C. P. Chia 2000.

A role for calcineurin in Dictyostelium discoideum phagocytosis. Session # 1696.

Published in Abstracts: 40th American Society for Cell Biology Annual Meeting, in *Molecular Biology of the Cell* **11** (supplement): p. 327a. Copyright © 2000 American Society for Cell Biology. Used by permission.

A role for calcineurin in Dictyostelium discoideum phagocytosis

Aidong Yuan¹, Catherine P. Chia², ¹School of Biological Sciences, University of Nebraska-Lincoln, 348 Manter Hall, Lincoln, Nebraska 68588-0118, ²University of Nebraska

The Ca²⁺/calmodulin-dependent protein phosphatase calcineurin is involved in the development of the cellular slime mold *Dictyostelium discoideum*. Because of its interactions with Ca²⁺, which appear to influence *D. discoideum* phagocytosis (Yuan and Chia, 1999, *Mol. Biol. Cell* 10, 220a), we undertook studies to test whether calcineurin also plays a role in *Dictyostelium* phagocytosis. The immunosuppressants cyclosporin A and FK506, through the formation of cyclosporin A-cyclophilin A and FK506-FK506-binding protein complexes, respectively, inhibited calcineurin activity. These two calcineurin inhibitors suppressed phagocytosis of fluorescently labeled yeast in a dose-dependent manner. Although it inhibited phagocytosis, cyclosporin A had an insignificant effect on the macropinocytosis of the fluid-phase marker fluorescein isothiocyanate-dextran. Furthermore, trifluoperazine, a calmodulin antagonist that indirectly inhibits calcineurin, also suppressed phagocytosis in a dose-dependent fashion and induced the formation of giant intracellular vacuoles. Fluorescence microscopy of cyclosporin A-treated (for 30 min.) cells stained with rhodamine-phalloidin had cytoplasmic chunks of F-actin that were not present in control cells, while cells treated with FK506 and trifluoperazine (also for 30 min.), displayed less cortical but more cytoplasmic F-actin staining than normal cells. Typically, drug-treated cells were smaller and rounder than untreated cells. Our data suggest calcineurin may play a role in *D. discoideum* phagocytosis, either through the dephosphorylation of actin-regulating proteins or other cytoskeletal proteins such as the heavy chain subunit of nonmuscle myosin II since dephosphorylation of the latter promotes filament assembly.