University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Human–Wildlife Interactions

Wildlife Damage Management, Internet Center for

2008

Sharpshooting suburban white-tailed deer reduces deer–vehicle collisions

Anthony J. DeNicola *White Buffalo Inc.*

Scott C. Williams Connecticut Agricultural Experiment Station

Follow this and additional works at: http://digitalcommons.unl.edu/hwi Part of the <u>Environmental Health and Protection Commons</u>

DeNicola, Anthony J. and Williams, Scott C., "Sharpshooting suburban white-tailed deer reduces deer-vehicle collisions" (2008). *Human-Wildlife Interactions*. Paper 69. http://digitalcommons.unl.edu/hwi/69

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Human–Wildlife Interactions by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Sharpshooting suburban white-tailed deer reduces deer-vehicle collisions

ANTHONY J. DENICOLA, White Buffalo Inc., 26 Davison Road, Moodus, CT 06469, USA wbuffaloinc@aol.com

SCOTT C. WILLIAMS, Connecticut Agricultural Experiment Station, 123 Huntington Street, New Haven, CT 06511, USA

Abstract: Too many deer-vehicle collisions (DVCs) are one of the primary reasons local governments implement lethal deer management programs. However, there are limited data to demonstrate that a reduction in deer (*Odocoileus* spp.) densities will result in a decline in DVCs. We conducted sharpshooting programs in 3 suburban communities to reduce deer numbers and to address rising DVCs. Annual or periodic population estimates were conducted using both helicopter snow counts and aerial infrared counts to assess population trends. Management efforts were conducted from 3 to 7 years. Local deer herds were reduced by 54%, 72%, and 76%, with resulting reductions in DVCs of 49%, 75%, and 78%, respectively. These projects clearly demonstrate that a reduction in local deer densities using lethal methods can significantly reduce DVCs.

Key words: deer-vehicle collisions, human-wildlife conflict, Odocoileus virginianus, sharpshooting, suburban, white-tailed deer, wildlife damage management

DEER-VEHICLE COLLISIONS (DVCs) and their deer (Storm et al. 2007). Our experience is that related public safety concerns are one of the elevated numbers of DVCs are often the only most significant conflicts that arise when whitetailed deer (Odocoileus virginianus) become abundant in urban and suburban environments (Ng et al. 2008). Conover et al. (1995) estimated that there are >1 million DVCs in the United ductions in deer densities and a reduction in States annually, and >200 human deaths are DVCs. Therefore, our objective was to examine attributed to these events.

It has been demonstrated that DVCs increase as local deer populations increase (Hygnstrom in 3 suburban counties. and VerCauteren 1999, Etter et al. 2000, Hussain et al. 2007, Grovenburg et al. 2008, Rutberg and Naugle 2008). One could logically conclude that a reduction in deer abundance would lead to a ment projects in Iowa City, Iowa; Princeton, decline in DVCs (Mastro et al. 2008). The only New Jersey; and Solon, Ohio. Management way to reduce deer numbers efficiently and efforts were conducted in Iowa City during effectively is through the removal of deer from January 2000, December 2000–January 2001, and a local population (DeNicola et al. 2000, Rutberg December 2001–January 2002. We implemented et al. 2004). In most states, live-trapping and the sharpshooting program in Princeton during relocating deer is not an option because of high February–March 2001, January–February 2002, costs, disease transmission risks (e.g., chronic February-March 2003, January-February 2004, wasting disease), unavailability of suitable re- January 2005, and February 2006. We culled lease sites, and concerns over stress to captured deer in Solon during February–March 2005 and deer. Furthermore, most relocated deer do not January-March 2006. survive a year in their new environs (Conover 2002). Therefore, only lethal management op- 15.5 km² area in Iowa City, all of Princeton tions (i.e., hunting, sharpshooting, and live cap- Township (36.3 km²), and all of Solon (51.8 km²). ture followed by euthanasia) can potentially These communities were typical suburban dereduce deer densities in the short term.

sociated with discussions about how to ad- intermingled small agricultural plots and undress conflicts associated with an abundance of developed open spaces. Public safety concerns

conflict that local politicians feel comfortable using to justify the authorization of lethal deer management options. However, no one has demonstrated a clear correlation between reif the implementation of a sharpshooting management program reduced the number of DVCs

Study areas

We implemented sharpshooting manage-

Management activities were focused in a velopments that were composed of a matrix of There is often considerable controversy as- suburban and commercial development, with



Anthony DeNicola sharpshooting deer.

over increasing DVCs was the reason elected officials approved the use of sharpshooting to reduce the local deer herds.

Methods

We used sharpshooting techniques to kill deer (DeNicola et al. 1997). We selected bait sites throughout the area of operation before beginning sharpshooting efforts in order to maximize the efficiency and safety of removal efforts. We would attempt to have a minimum of 2 bait sites per km². Whole kernel corn was placed on the ground 3 weeks in advance of shooting efforts at select shooting locations. We would place approximately 0.5 to 1 kg of corn per deer daily at each site. Sharpshooting sites were accessed from a vehicle or from a tree stand, during the daytime and after dark. Human safety was ensured by shooting only when there was a known earthen backstop created through the shooter's relative elevation (e.g., tree stand) or topography. Deer were shot only when circumstances were safe (i.e., with no humans in the removal zone). To prevent educating deer to the procedure, we shot when fewer than 9 deer were present. Although we shot deer on a first opportunity basis, when possible antlerless deer were prioritized.

Population estimates were derived using helicopter counts over snow following methods described in Beringer et al. (1998) in Princeton (February 2002). Biologists from the Iowa Department of Natural Resources conducted less systematic helicopter counts over snow in Iowa City (1999–2002). These counts were done by the same personnel using the methods annually. Therefore, the Iowa City counts were minimum estimates and were not adjusted for detection rates. Aerial infrared (IR) censusing techniques (Naugle et al. 1996) were used over the entire management area in Princeton (December 2004) and Solon (March 2004, December 2005) to estimate population size. All IR counts were conducted by Davis Aviation (Kent, Oh.). Infrared counts were conducted using a single-engine Cessna 182 with a fuselage-mounted, highresolution Mitsubishi M-600 thermal imager. Transects were designed at 100-m intervals and flown at 500 m above ground. At this elevation 100% coverage was achieved and verified with Global Positioning System (GPS) moving map software. Flights were conducted after 2200 hours to ensure adequate ground cooling and good thermal contrast. The thermal imager output was routed through a video encoderdecoder and recorded on digital videotape for later review. Count data in Princeton and Solon were complemented by conducting simple population projections based on observed demographics (DeNicola 2008). We assumed that 60% of the populations were female, 33% of females were fawns, and recruitment rates were 1:1 (doe:fawn ratio). We then included approximations of non-culling mortality (i.e., DVC and hunter harvest data, when available, and approximate mortality rates for urban deer from the literature (Etter et al. 2002). Immigration and emigration were assumed to be equal.

DVCs were tallied using police reports in Iowa City and a combination of police reports and road-kill collection records (i.e., location of carcasses removed from the roadway) of animal control officers in Princeton and Solon. Data collection methods were consistent among years at all locations. Using Microsoft Excel, we conducted linear regression analyses comparing recorded DVCs to estimated deer densities.

Results

During 49 days over 3 years of sharpshooting at the Iowa City site, we removed 950 deer, with a culling rate of 19.3 deer per day. We culled 1,455 deer during 118 days from Princeton (12.3 deer/day) during a 6-year period. We spent 77 days sharpshooting during 2005 and 2006 to remove 1,002 deer from Solon (13.0 deer/day). At all communities, we removed more deer and at a higher culling rate per day the first year we started sharpshooting than in subsequent years (Table 1).

Following the implementation of sharpshooting program, deer numbers and the annual number of DVCs were reduced from 49% to 78% in the study sites (Table 2). The highest percentage reduction (78%) occurred in Iowa City where the deer density was reduced by 76%. Reductions in deer densities and DVCs remained suppressed during the entire period of the sharpshooting program and did not rebound in later years (Table 3). There was a direct correlation between annual deer population and DVCs in all study sites (Iowa City, $r^2 = 0.72$, F = 5.0, df = 1, 2; P > 0.05; Princeton, $r^2 = 0.98, F =$ 203.5, df = 1, 5; P < 0.05; Solon, $r^2 = 0.99$, F= 85.1, *df* = 1, 1; *P*> 0.05; Figure 1).

Discussion

DVCs become more frequent with an increase in deer densities (Etter et al. 2000, Mastro et al. 2008). However, our study is the only one we are aware of that

by accounting for harvest numbers, recruitment rates, and natural mortality occurring within the community. This further supports the observations by McNulty et al. (1997) of local deer management effects on deer movements.

In this study we also found that the percentage

TABLE 1. Number of days sharpshooting was conducted and deer shot and removed from 3 suburban communities during different winters (a winter may include December of the prior year).

Location	Winter	Days of effort	Num- ber deer killed	Number deer killed/ day
Iowa City, Ia.	2000	10	360	36.0
	2001	21	340	16.2
	2002	18	250	13.8
Princeton, N.J.	2001	15	322	21.5
	2002	27	303	11.2
	2003	21	280	13.3
	2004	27	276	10.2
	2005	13	124	9.5
	2006	15	150	10.0
Solon, Oh.	2005	37	602	16.3
	2006	40	400	10.0

TABLE 2. Number of deer and annual DVCs, both before (pre-cull) and after (post-cull) implementation of a deer sharpshooting program within 3 suburban communities.

Site location	Winter culling period	No. deer pre-cull	No. DVCs pre-cull	No. deer post-cull	No. DVCs post-cull	% pop. decline	% DVC de- cline
Iowa City, Ia.	2000-2002	371	63	91	14	76	78
Princeton, N.J.	2001-2006	1600	342	450	85	72	75
Solon, Oh.	2005–2006	1400	171	650	88	54	49

demonstrates that reducing local deer densities through a culling program reduces DVCs (Table 2). Although this relationship is quite intuitive, it is important to demonstrate it because most lethal management programs of suburban deer are motivated by the desire to protect citizens from the cost and danger of injury from DVCs. In our 3 study sites, we found no indication that there was any significant level of immigration of deer into the communities from outside areas or emigration out of the communities. Instead, deer density trends could be predicted of the deer population killed annually in DVCs consistently ranged from 13% to 20% (Figure 1). Solon had the lowest percentage of the deer population killed by vehicles each year (13.2%), and Princeton had the highest (20.7%). Anecdotal observations of high traffic volumes on narrow roads with low lateral visibility in Princeton may explain why deer are more vulnerable to DVCs there than in Iowa City or Solon.

Even though a management method may be proven effective, the relative implementation



FIGURE 1. Linear regression analysis of deer population size and annual number of DVCs in Iowa City, Iowa, 1999–2002, Princeton, New Jersey, 2000–2006, and Solon, Ohio, 2004–2006.

	1999	2000	2001	2002	2003	2004	2005	2006
Iowa City, Ia. (15.5 km²)								
Deer/km ²	23.9*	20.6	7.1	5.9				
DVCs/km ²	4.1*	2.8	2.5	0.9				
Princeton, N.J. (36.3 km ²)								
Deer/km ²		44.1*	27.5	22.0	17.9	16.5	15.2	12.4
DVCs/km ²		9.4*	6.7	4.7	3.5	3.5	2.8	2.3
Solon, Oh. (51.8 km ²)								
Deer/km ²						27.0*	18.3	12.5
DVCs/km ²						3.3*	2.5	1.7

TABLE 3. Annual changes in the density of deer and recorded DVCs for 3 suburban communities prior to (pre-cull) and after implementation of sharpshooting program to cull deer (data marked with an asterisk [*] were collected pre-cull).

cost-to-benefit ratio must be considered before it can be determined to be a practical solution. For example, culling costs would be balanced by savings from preventing damage to a vehicle involved in a DVC when the cost to cull a deer equals \$354, based on data from the Princeton site (DeNicola, unpublished data). Moreover, money saved by averting accidents does not include cost benefits associated with a reduction in human injuries and fatalities associated with DVCs (Bissonette et al. 2008). Additional benefits of reducing deer densities include a

reduction in landscape damage (Ward 2000, DeNicola et al. 2000), reduced numbers of blacklegged ticks (*Ixodes scapularis*) and associated cases of Lyme disease (Stafford et al. 2003), and reduced ecological damage to forested areas (Kelty and Nyland 1983, Kittredge et al. 1992, Conover 1997).

Acknowledgments

We would like to thank the local municipal governments and police departments of Iowa City, Iowa, Princeton, New Jersey, and Solon, Ohio, as well as all the cooperating landowners. In addition, we would like to extend our gratitude to the Iowa Department of Natural Resources, the New Jersey Division of Fish and Wildlife, and the Ohio Department of Natural Resources for all their cooperation and assistance.

Literature cited

- Beringer, J., L. P. Hanson, and O. Sexton. 1998. Detection rates of white-tailed deer with a helicopter over snow. Wildlife Society Bulletin 26:24–28.
- Bissonette, J. A., C. Kassar, and L. J. Cook. 2008. An assessment of costs associated with deer– vehicle collisions: human death and injury, vehicle damage, and deer loss. Human–Wildlife Conflicts 2:17–27.
- Conover, M. R. 1997. Monetary and intangible valuation of deer in the United States. Wildlife Society Bulletin 25:298–305.
- Conover, M. R. 2002. Resolving human–wildlife conflicts: the science of wildlife damage management. Lewis, Boca Raton, Florida, USA.
- Conover, M. R., W. C. Pitt, K. K. Kessler, T. J. DuBow, and W. A. Sanborn. 1995. Review of human injuries, illnesses, and economic losses caused by wildlife in the United States. Wildlife Society Bulletin 23:407–414.
- DeNicola, A. J., D. R. Etter, and T. Almendinger. 2008. Demographics of non-hunted whitetailed deer populations in suburban areas. Human–Wildlife Conflicts 2:102–109.
- DeNicola, A. J., K. C. VerCauteren, P. D. Curtis, and S. E. Hygnstrom. 2000. Managing whitetailed deer in suburban environments. Cornell Cooperative Extension, Ithaca, New York, USA.
- DeNicola, A. J., S. J. Weber, C. A. Bridges, and J. L. Stokes. 1997. Nontraditional techniques for management of overabundant deer populations. Wildlife Society Bulletin 25:496–499.
- Etter, D. E., T. R. Van Deelen, D. R. Ludwig, S. N. Kobal, and R. E. Warner. 2000. Management of white-tailed deer in Chicago, Illinois, Forest Preserves. Vertebrate Pest Conference 19:190–196.
- Etter, D. R., T. R. Van Deelen, D. R. Ludwig, S. N. Kobal, and R. E. Warner. 2002. Survival and movements of white-tailed deer in suburban Chicago, Illinois. Journal of Wildlife Management 66:500–510.

- Grovenburg, T. W., J. A. Jenks, R. W. Klaver, K. L. Monteith, D. H. Galster, R. J. Schauer, W. W. Morlock, and J. A. Delger. 2008. Factors affecting road mortality of white-tailed deer in eastern South Dakota. Human–Wildlife Conflicts 2:48–59.
- Hussain, A., J. B. Armstrong, D. B. Brown, and J. Hogland. 2007. Land-use pattern, urbanization, and deer–vehicle collisions in Alabama. Human–Wildlife Conflicts 1:89–96.
- Hygnstrom, S. E., and K. C. VerCauteren. 1999. Ecology of white-tailed deer in the Gifford Point-Fontenelle Forest area, Nebraska. Final Report. University of Nebraska, Lincoln, Nebraska, USA.
- Kelty, D. B., and R. D. Nyland. 1983. Hardwood browse production following shelterwood cutting. Journal of Wildlife Management 47:1216– 1220.
- Kittredge, D. B., M. J. Kelty, and P. M. S. Ashton. 1992. The use of tree shelters with northern red oak natural regeneration in southern New England. Northern Journal of Applied Forestry 9:141–145.
- Mastro, L., M. R. Conover, and S. N. Frey. 2008. Deer–vehicle collision prevention techniques. Human–Wildlife Conflicts 2:80–92.
- McNulty, S. A., W. F. Porter, N. E. Mathews, and J. A. Hill. 1997. Localized management for reducing white-tailed deer populations. Wildlife Society Bulletin 25:265–271.
- McShea, W. J., C. M. Stewart, L. J. Kearns, S. Liccioli, and D. Kocka. 2008. Factors affecting autumn deer–vehicle collisions in a rural Virginia county. Human–Wildlife Conflicts 2:110–121.
- Naugle, D. E., J. A. Jenks, and B. J. Kernohan. 1996. Use of thermal infrared sensing to estimate density of white-tailed deer. Wildlife Society Bulletin 24:37–43.
- Ng, J. W., C. Nielsen, and C. C. St. Clair. 2008. Landscape and traffic factors influencing deer– vehicle collisions in an urban environment. Human–Wildlife Conflicts 2:34–47.
- Rutberg, A. T., and R. E. Naugle. 2008. Deer–collision trends at a suburban immunocontraception site. Human–Wildlife Conflicts 2:60–67.
- Rutberg, A. T., R. E. Naugle, L. A. Thiele, and I. K. M. Liu. 2004. Effects of immunocontraception on a suburban population of white-tailed deer *Odocoileus virginianus*. Biological Conservation 116:243–250.

Stafford, K. C., III, A. J. DeNicola, and H. J. Kil-

patrick. 2003. Reduced abundance of *Ixodes* scapularis (Acari: Ixodidae) and the tick parasitoid *Ixodiphagus hookeri* (Hymenoptera: Encyrtidae) with reduction of white-tailed deer. Journal of Medical Entomology 40:642–652. Storm, D. J., C. K. Nielsen, E. M. Schauber, and A. Woolf. 2007. Deer–human conflict and hunter access in an exurban landscape. Human–Wild-life Conflicts 1:53–59.

cyrtidae) with reduction of white-tailed deer. Ward, J. S. 2000. Limiting deer browse damage Journal of Medical Entomology 40:642–652. to landscape plants. Connecticut Agricultural brm, D. J., C. K. Nielsen, E. M. Schauber, and A.



ANTHONY J. DENICOLA is president of White Buffalo, Inc., a nonprofit research organization dedicated to conserving ecosystems through wildlife population control. He received a B.S. degree in biology from Trinity College of Hartford, Connecticut, an M.S. degree from the Yale School of Forestry and Environmental Studies, and a Ph.D. degree in wildlife ecology from Purdue University. His Ph.D. research involved the examination of the efficacy of contraceptives on 2 enclosed deer herds. He has conducted field trials using contraceptives on several hundred white-tailed deer in 7 study areas over the last 15 years. He also has implemented sharpshooting programs in 11 states, removing >7,000 deer from suburban communities. He is a member of the National Animal Damage Control Association, the Society for Conservation Biology, and The Wildlife Society. He holds research affiliate positions with Trinity College, the Denver Zoological Foundation, and Rutgers University. His professional interests are behavioral-ecological approaches to wildlife damage control, wildlife reproductive control, and control of introduced vertebrate species.



SCOTT C. WILLIAMS is employed at the Connecticut Agricultural Experiment Station, Department of Forestry and Horticulture. He has his B.A. degree in environmental studies from Connecticut College, and his M.E.S. degree from the Yale School of Forestry and Environmental Studies. He is a Ph.D. degree candidate in wildlife management at the University of Connecticut. He has been collaborating with White Buffalo, Inc., on his Ph.D. research, investigating behavioral changes of white-tailed deer subjected to sharpshooting, controlled hunting, and baiting. His research interests include the ecological and public health impacts of overabundant white-tailed deer herds, including deer-vehicle collisions, ecological damage, alteration of forest species composition, avoiding damage to ornamental vegetation, and deer as exotic seed dispersal agents. In his limited free time, he is an avid hunter and angler, and in early spring, he runs his own maple syrup operation.