

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Papers in Microbiology

Papers in the Biological Sciences

March 1999

Nonspecific interactions alter lipopolysaccharide patterns and protein mobility on sodium dodecyl sulfate polyacrylamide gels

Aidong Yuan

University of Nebraska-Lincoln

R L. Pardy

rpardy2@unl.edu

Catherine P. Chia

University of Nebraska-Lincoln, cchia1@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/bioscimicro>



Part of the [Microbiology Commons](#)

Yuan, Aidong; Pardy, R L.; and Chia, Catherine P., "Nonspecific interactions alter lipopolysaccharide patterns and protein mobility on sodium dodecyl sulfate polyacrylamide gels" (1999). *Papers in Microbiology*. 73.

<https://digitalcommons.unl.edu/bioscimicro/73>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Microbiology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Submitted March 8, 1999

Nonspecific interactions alter lipopolysaccharide patterns and protein mobility on sodium dodecyl sulfate polyacrylamide gels

Aidong Yuan, Roosevelt L. Pardy, and Catherine P. Chia

School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, USA

Correspondence: Dr. Aidong Yuan, School of Biological Sciences, 348 Manter Hall, University of Nebraska-Lincoln, Lincoln, NE 68588-0118, USA. Email: dyuan@unserve.unl.edu Fax: 402-472-2083

Abstract: In testing whether bacterial lipopolysaccharide (LPS) was a natural substrate for an esterase from the soil amoebae *Dictyostelium discoideum*, we observed altered banding patterns of the LPS and changed protein mobility on sodium dodecyl sulfate (SDS) polyacrylamide gels after incubation of LPS with the enzyme. The initial interpretation of these results was that the enzyme had removed ester-linked acyl chains from the LPS, leading to a change in its migration on gels. However, esterase inactivated by treatment with either dithiothreitol (DTT), heat, or SDS generated the same mobility shifts. Bovine serum albumin (BSA) also induced the same change in the electrophoretic pattern. We conclude that the altered LPS patterns and protein mobility on SDS gels were caused by nonspecific interactions between LPS and protein.

Keywords: Lipopolysaccharide, Lipopolysaccharide-protein interaction, Ladder-like pattern, Gel electrophoresis

Abbreviations: LPS, lipopolysaccharide; SPB, sample preparation buffer

1 Introduction

Bacterial lipopolysaccharides (LPS), also known as endotoxin, are the major component of the outer membrane of Gram-negative bacteria on which amoebae of the slime mold *Dictyostelium discoideum* feed. *D. discoideum* cells contain an esterase activity which specifically cleaves long-chain fatty acids from the lipid A component of bacterial lipopolysaccharides [1]. In studies of cytoskeletally associated proteins, we found a 70 kDa glycoprotein (gp70) enriched in the detergent-insoluble fraction of axenically grown cells [2]. *N*-terminal amino acid sequence of gp70 identified it to be "crystal protein" [3] and biochemical assays demonstrated its esterase activity [2]. Changes in the electrophoretic pattern of bacterial LPS in SDS polyacrylamide gels were interpreted as evidence of LPS degradation by gut juice of the snail *Helix pomatia* [4]. We used this method to test whether bacterial LPS was a natural substrate of gp70. After incubating LPS and gp70 together, we observed altered LPS banding patterns as well as changes in the mobility of gp70 on SDS gels. The initial interpretation of these results was that cleavage of ester-linked acyl chains from LPS by gp70 led to a change in the migration of the LPS. However, the same altered LPS ladder pattern and shift in gp70 mobility were generated when inactivated esterase or BSA was used. Therefore, careful interpretation of modified LPS profiles on SDS gels is warranted when LPS-protein interactions are postulated.

2 Materials and methods

2.1 Preparation of LPS and gp70

LPS was purified from late log phase cultures of *Serratia marcescens* (a gift of Dr. K. Nickerson, University of Nebraska-Lincoln) following proteinase K digestion [5]. Gp70 was purified from the cytoskeletal fraction of the transformed *D. discoideum* cell line AT-K2 that overexpresses crystal protein [2]. Briefly, cytoskeletons were prepared and solubilized from AT-K2 cells grown to 1×10^7 cells/mL at 20°C in HL5 media [6] containing 10 mg/mL G418 [2]. Gp70 was isolated from the cytoskeletal fraction either by immunoaffinity chromatography [7] using protein G-purified monoclonal antibody 130-80-2 specific for gp70 [3], or Concanavalin A affinity chromatography [2], or by centrifuging the solubilized cytoskeletons at $10,000 \times g$ for 1 h to pellet detergent-insoluble actin filaments and associated proteins. The concentration of purified gp70 was determined by comparison to known amounts of BSA run concurrently on Coomassie blue-stained SDS gels.

2.2 Assay for esterase activity of gp70

The esterase activity of gp70 was determined spectrophotometrically using the substrate α -naphthyl acetate in a general esterase assay as described previously [2]. Alternately, one milligram of LPS from *Serratia marcescens* was dissolved in 1 mL of 50 mM MOPS buffer, pH 7.5, and sonicated at 4°C for 30 min in a FS3 ultrasonic water bath (Fisher

Scientific, Hampton, NH) which made the LPS solution appear opalescent. Twenty microliters (20 μg) of the LPS solution were incubated with 5 μL (0.2 $\mu\text{g}/\mu\text{L}$) of gp70 and 15 μL of deionized water at 30°C for 30 min. In some experiments, BSA instead of gp70, or gp70 inactivated (verified using the general esterase assay) by treatment with 250 mM DTT (4°C for 30 min), 4% SDS (4°C for 30 min) or heat (10 min at 100°C) was mixed with the LPS.

2.3 Gel electrophoresis

The incubated LPS-protein mixture (40 μL) was heat-denatured (100°C for 5 min) with 10 μL of 5 \times SDS Laemmli sample preparation buffer (SPB) [8]. Ten microliters were loaded per lane of 7.5% polyacrylamide SDS (mini)-gels [8] containing 4 M urea [9]. Prestained protein standards of high molecular mass (Gibco BRL Life Technologies, Rockville, MD) were: myosin heavy chain (200 kDa), phosphorylase (97 kDa), BSA (68 kDa), ovalbumin (43 kDa), and carbonic anhydrase (29 kDa).

2.4 Silver staining

Gels were stained with silver according to Tsai and Frasch [9] with minor modifications. Briefly, gels were fixed for 30 min in 40% ethanol-5% acetic acid, and then soaked in oxidant (0.7% periodic acid in 40% ethanol-5% acetic acid) for 5 min. After three 5 min washes in water, gels were incubated for 10 min in a freshly prepared ammoniacal silver nitrate solution (12 mL of 0.1 N NaOH, 0.8 mL of NH_4OH (28.8%), 2 mL of 20% AgNO_3 and 46 mL deionized distilled water). After three water washes, gels were developed in freshly prepared 3% Na_2CO_3 containing 0.0185% formaldehyde. The development was stopped with 50 mM EDTA, pH 4.6, when the stain reached the desired intensity.

2.5 Immunoblotting

Proteins were transferred from gels onto polyvinylidene difluoride membranes (Millipore, Bedford, MA) by semidry electroblotting using Towbin buffer [10]. Blots were shaken for 15 min in 5% nonfat milk in 150 mM NaCl, 10 mM Tris, pH 7.5, to minimize nonspecific binding and washed three times (5 min each) in Tween buffer (0.1% Tween-20, 150 mM NaCl, 10 mM Tris, pH 8.0). They were incubated for 1 h in undiluted hybridoma cell culture supernatant containing a mouse monoclonal antibody, 130-80-2, specific for gp70 [2, 3]. Blots were washed again in the Tween buffer and incubated for 1 h in Tween buffer containing a 1:12,500 dilution of alkaline phosphatase-goat anti-mouse IgG (Sigma, St. Louis, MO). Blots were washed again in the Tween buffer before color development with the substrates

5-bromo-4-chloro-3-indolyl phosphate and nitroblue tetrazolium [7].

3 Results and discussion

LPS from *Serratia marcescens* had a ladder-like pattern on SDS gels (containing 4 M urea) of eleven bands (Fig. 1, lane 1), consistent with a previously published pattern [11]. The LPS molecules migrated in the range of 35–55 kDa. The protein markers were not appropriate standards for estimating the mass of LPS molecules and were used only as references for monitoring the relative mobilities of the LPS. After incubation with purified gp70, whose activity was verified using a general esterase assay, the ladder-like pattern of *Serratia marcescens* LPS partly disappeared with only two bands of 45–48 kDa remaining (indicated with +; lanes 2 and 3). An intensely staining band at roughly 180 kDa was also observed (indicated with *). The initial interpretation of the pattern alteration was that the enzyme had removed ester-linked acyl chains from the LPS, leading to a change in its migration on gels.




Figure 1. Disappearance of *Serratia marcescens* LPS bands after incubation with purified *D. discoideum* gp70 on an SDS gel containing 4 M urea. (See Section 2.2 for assay conditions.) Lane (1) LPS (4 μg), (2) LPS incubated with active gp70, (3) LPS incubated with DTT-inactivated gp70, and (4) purified gp70 (0.2 μg). In (1) the upper two white boxes outline seven more strongly staining species between 43 and 55 kDa and the lower box outlines four fainter species in the range of 35–42 kDa. In (2) and (3) the symbol (+) indicates two LPS species at approximately 45–48 kDa remaining after incubation, and the symbol (*) indicates apparent LPS-protein associations. The 65 kDa species below gp70 was a breakdown product that occurred upon storage. Sizes (in kDa) of prestained protein molecular markers are indicated to the left of (1).

Changes in the electrophoretic pattern of LPS bands in SDS gels have been interpreted as evidence of LPS degradation by the gut juice of the snail *Helix pomatia* [4]. However, LPS, incubated with gp70 completely inactivated by treatment with DTT, as judged by the esterase assay, generated the same altered ladder pattern (Fig. 1, lane 3) seen with active gp70 (Fig. 1, lane 2). Additional experiments with gp70 denatured with SDS or by boiling also generated the same altered ladder pattern (data not shown). Even mixing LPS in the wells of SDS gels with increasing amounts of the gp70 boiled in SPB for 5 min also caused the change in the LPS profile (Fig. 2A). To test whether non-specific protein interactions were responsible for the changes LPS pattern, LPS was incubated under the same conditions with BSA in place of gp70.




Figure 2. Gradual disappearance of the LPS ladder patterns by (A) increasing amounts of heat-inactivated gp70, and (B) retarded migration and smearing of gp70. (A) Silver-stained SDS gel that contained 4 M urea; (B) immunoblot of duplicate gel shown in (A) stained with monoclonal antibody 130-80-2 against *D. discoideum* gp70. LPS (4 μ g per lane, in 1 \times SPB) was loaded first into lanes 1–8 and then the following amounts of heat-inactivated gp70 in 1 \times SPB were loaded on top of the LPS: lane (1) 0 μ g, (2) 0.02 μ g, (3) 0.04 μ g, (4) 0.06 μ g, (5) 0.08 μ g, (6) 0.1 μ g, (7) 0.12 μ g, (8) 0.2 μ g, and (9) 0.1 μ g. LPS was omitted from (9). Sizes (in kDa) of prestained protein molecular markers are indicated to the left of (1). The current was turned on immediately after loading the gel.

BSA induced the same change in the electrophoretic pattern of LPS on the gel (Fig. 3). Because the modified LPS pattern was observed whether gp70 was active or denatured, and with BSA, we conclude that the loss of most of the ladder was not the result of enzymatic hydrolysis of LPS subcomponents but rather an artifact of LPS-protein interactions unrelated to the esterase activity of gp70.

The LPS-gp70 interaction caused not only the alteration of the ladder pattern of LPS but also a change in the mobility of the esterase on the gels (Fig. 2B). As shown by immunoblotting, gp70 alone migrated at 67 kDa on the blot (Fig. 2B, lane 9). Once mixed with LPS, gp70 moved as a much larger species (137 kDa; Fig. 2B, lanes 2 and 3), probably as a result of an association with LPS. With increasing amounts of gp70 mixed with LPS, the size of the LPS-gp70 complexes ranged from 67 to 200 kDa, based on the smeary signals on both the silver-stained gel and the blot (Fig. 2A and 2B, lanes 4–8).

Previous studies have shown that bacterial LPS can bind to or associate with many proteins including serum LPSbinding protein LBP [12], CD14 [13], the macrophage scavenger receptor, the beta2 integrins [14], hemoglobin [15], moth hemolin (an insect member of the Ig-superfamily) [16], OmpA of *Escherichia coli* [17], the S2 subunit of pertussis toxin [18], the 39 kDa membrane protein of *Haemophilus influenzae* type b [19], and several plasma proteins of rainbow trout [20]. More pertinent to the present study, however, were LPS-protein




Figure 3. Gradual disappearance of the LPS ladder pattern by increasing amounts of BSA. LPS (4 mg) was loaded in lanes (1)–(8) of an SDS gel containing 4 M urea, and then the following amounts of BSA in 1 \times SPB were loaded on top of LPS: (1) 0 μ g, (2) 0.02 μ g, (3) 0.04 μ g, (4) 0.06 μ g, (5) 0.08 μ g, (6) 0.1 μ g, (7) 0.12 μ g, (8) 0.2 μ g, and (9) 0.1 μ g. LPS was omitted from (9). Sizes (in kDa) of prestained protein molecular markers are indicated to the left of (1). The current was turned on immediately after loading the gel.

interactions resistant to dissociation under harsh conditions (i.e., SDS, urea, and high temperature). Similar observations were made with gp70 and Klebsiella-LPS mixtures (data not shown). LPS migration patterns from *Neisseria gonorrhoeae* [21] and Salmonelle chemotypes [5] are more compact in whole cell lysates compared to their profiles in proteinase K-digested lysates or purified LPS, implicating LPS-protein interactions in producing the altered patterns. The reduced LPS profile in whole-cell lysates was demonstrated to be due to the stable association between LPS and many membrane proteins [21].

The conclusions by Saddler et al. [4], who interpreted changes in LPS patterns on SDS-PAGE to be evidence of snail gut juice degradation of LPS, are not invalidated by our findings. In their study, however, no chemical evidence was presented for the degradation of LPS. An alternative explanation for the observed altered gel patterns would be an association of LPS molecules with (protein) components in the gut juice. This interpretation is plausible and consistent with our observations supporting the idea that nonspecific LPS-protein interactions modified the mobility of both LPS and protein on SDS gels.

4 Concluding remarks

The conclusion of this study is that nonspecific LPS-protein interactions can alter LPS ladder patterns and protein mobility on SDS gels. The observed noncatalytic LPS-protein associations may lead to the misinterpretation of experimental results if the interactions are not taken into account when analyzing LPS profiles on gels, purifying bacterial antigens as vaccines and studying enzyme effects on LPS.

We are grateful to Dr. G. Gerisch (Max-Planck Institut für Biochemie, Germany) for providing the hybridoma producing a mouse monoclonal antibody, 130-80-2, specific for gp70 (crystal protein). We also thank Dr. A. Zera (School of Biological Sciences, University of Nebraska- Lincoln) for assistance with protocols to assay esterase activity. These studies were supported by funds from the University of Nebraska Research Council (College of Arts and Sciences) and NSF Grant MCB-95133628 to C.P.C.

5 References

- [1] Malchow, D., Luderitz, O., Kickhofen, B., Westphal, O., Gerisch, G., *Eur. J. Biochem.* 1969, 7, 239-246.
- [2] Chia, C. P., Bomblies, L., Taylor, K. K., *Exp. Cell Res.* 1998, 244, 340-348.
- [3] Bomblies, L., Biegelmann, E., Doring, V., Gerisch, G., Krafft-Czepa, H., Noegel, A. A., Schleicher, M., Humbel, B. M., *J. Cell Biol.* 1990, 110, 669-679.
- [4] Saddler, J. N., Parton, R., Wardlaw, A. C., *Experientia* 1979, 35, 494-495.
- [5] Hitchcock, P. J., Brown, T. M., *J. Bacteriol.* 1983, 154, 269-277.
- [6] Watts, D. J., Ashworth, J. A., *Biochem. J.* 1970, 119, 171-174.
- [7] Harlow, E., Lane, D., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York 1988, pp. 511-552.
- [8] Laemmli, U. K., *Nature* 1970, 227, 680-685.
- [9] Tsai, C. M., Frasch, C. E., *Anal. Biochem.* 1982, 119, 115-119.
- [10] Towbin, H., Staehelin, T., Gordon, J., *Proc. Natl. Acad. Sci. USA* 1979, 76, 4350-4354.
- [11] Poole, K., Braun, V., *J. Bacteriol.* 1988, 170, 5146-5152.
- [12] Schumann, R. R., Leong, S. R., Flaggs, G. W., Gray, P. W., Wright, S. D., Mathison, J. C., Tobias, P. S., Ulevitch, R. J., *Science* 1990, 249, 1429-1431.
- [13] Wright, S. D., Ramos, R. A., Tobias, P. S., Ulevitch, R. J., Mathison, J. C., *Science* 1990, 249, 1431-1433.
- [14] Fenton, M. J., Golenbock, D. T., *J. Leuk. Biol.* 1998, 64, 25-32.
- [15] Kaca, W., Roth, R. I., Levin, J., *J. Biol. Chem.* 1994, 269, 25078-25084.
- [16] Daffre, S., Faye, I., *FEBS Lett.* 1997, 408, 127-130.
- [17] Barbas, J. A., Vazquez, D., Rodriguez-Tebar, A., *FEBS Lett.* 1985, 193, 289-293.
- [18] Lei, M. G., Morrison, D. C., *J. Biol. Chem.* 1993, 268, 1488-1493.
- [19] Gulig, P. A., Hansen, E. J., *Infect. Immun.* 1985, 49, 819-827.
- [20] Hoover, G. J., el-Mowafi, A., Simko, E., Kocal, T. E., Ferguson, H. W., Hayes, M. A., *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 1998, 120, 559-569.
- [21] Hitchcock, P. J., *Infect. Immun.* 1984, 46, 202-212.