

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

School of Natural Resources: Faculty  
Publications

Natural Resources, School of

---

3-20-1979

## A Gering Formation (Miocene) Pumice Conglomerate and Associated Beds from Broadwater, Nebraska

Robert F. Diffendal Jr.

*University of Nebraska-Lincoln*, [rfd@unl.edu](mailto:rfd@unl.edu)

Follow this and additional works at: <https://digitalcommons.unl.edu/natrespapers>



Part of the [Geomorphology Commons](#), [Natural Resources and Conservation Commons](#), [Sedimentology Commons](#), and the [Stratigraphy Commons](#)

---

Diffendal, Robert F. Jr., "A Gering Formation (Miocene) Pumice Conglomerate and Associated Beds from Broadwater, Nebraska" (1979). *School of Natural Resources: Faculty Publications*. 83.  
<https://digitalcommons.unl.edu/natrespapers/83>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in School of Natural Resources: Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Published in *Proceedings of the Nebraska Academy of Sciences and  
Affiliated Societies*, 89th Annual Meeting, April 20–21, 1979, p.  
49.

Copyright © 1979 Robert F. Diffendal.

A GERING FORMATION (MIOCENE) PUMICE CONGLOMERATE AND ASSOCIATED BEDS FROM  
BROADWATER, NEBRASKA

R. F. Diffendal, Jr., Natural Science Division, Doane College, Crete,  
Nebraska 68333

The Mitchell Pass Member of the Gering Formation was defined by Vondra, Schultz, and Stout in 1969 on the basis of exposures in the Wildcat Ridge area of western Nebraska. The basal part of the member in many areas of Wildcat Ridge is a pumice-pebble conglomerate bed.

A newly discovered pumice-pebble conglomerate locality occurs north of Broadwater, Nebraska, some twenty miles east of the easternmost previously reported exposure of the pumice conglomerate at Redington Gap on Wildcat Ridge. Pumice samples from Redington Gap and Broadwater have the same index of refraction and similar inclusions. Beds above the conglomerate and below the Monroe Creek Formation at Broadwater have yielded fossil oreodonts typical of the Gering Formation. This evidence tends to support suggestions by Stout that the exposures north of Broadwater are an eastern continuation of at least part of the Gering Formation into the Broadwater area.