

April 1985

# MIDDLE MIOCENE TO RECENT STRATIGRAPHY AND PALEOGEOGRAPHY OF WESTERN NEBRASKA

James B. Swinehart

*University of Nebraska - Lincoln*, [jswinehart1@unl.edu](mailto:jswinehart1@unl.edu)

Vernon L. Souders

*University of Nebraska - Lincoln*

H. M. DeGraw

*University of Nebraska - Lincoln*

Robert F. Diffendal

*University of Nebraska - Lincoln*, [rdiffendal1@unl.edu](mailto:rdiffendal1@unl.edu)

Follow this and additional works at: <http://digitalcommons.unl.edu/natrespapers>



Part of the [Natural Resources and Conservation Commons](#)

---

Swinehart, James B.; Souders, Vernon L.; DeGraw, H. M.; and Diffendal, Robert F., "MIDDLE MIOCENE TO RECENT STRATIGRAPHY AND PALEOGEOGRAPHY OF WESTERN NEBRASKA" (1985). *Papers in Natural Resources*. 87.

<http://digitalcommons.unl.edu/natrespapers/87>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Natural Resources by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Published in *Proceedings of the Nebraska Academy of Sciences, including the GNATS & TER-QUA Divisions, and Nine Affiliated Societies*, 95th Annual Meeting, April 12–13, 1985, p. 56.

Copyright © 1985 J. B. Swinehart, V. L. Souders, H. M. DeGraw, and R. F. Diffendal, Jr.

MIDDLE MIOCENE TO RECENT STRATIGRAPHY AND PALEO GEOGRAPHY OF WESTERN  
NEBRASKA

J.B. Swinehart, V.L. Souders, H.M. DeGraw, and R.F. Diffendal, Jr.,  
Conservation and Survey Division, IANR, University of Nebraska-Lincoln,  
Lincoln, NE 68588-0517

Two generally distinctive suites of sediments occur in western Nebraska, one dominantly volcanoclastic in origin deposited from the Oligocene through Early Miocene, and the other mainly epiclastic in origin deposited from the Middle Miocene through the Holocene. We have combined the epiclastic deposits usually placed in the Hemingford and Ogallala groups into an expanded Ogallala Group and have abandoned the member subdivisions of the Broadwater Formation.

About 19 million years ago, at the end of Arikaree deposition in western Nebraska, a major pulse of erosion (pre-Runningwater Formation, Ogallala Group), was followed by a fundamental change in depositional style and landscape evolution, characterized by a heterogeneous mixture of epiclastic valley fills. Sands and gravels from Rocky Mountain sources were first deposited in a major valley in the northern half of the area and later in valleys to the south. Episodic regional and local structural movements influenced the size and position of many Ogallala valleys. For the past 5 m.y. degradation, in response to major regional uplift, has greatly exceeded aggradation as the dominant factor affecting landscape evolution in western Nebraska. We believe that the modern Platte River system in its broad pattern began to form no earlier than late in the history of the Ash Hollow Formation of the Ogallala Group. The principal development of the North and South Platte valleys is Pliocene and younger, while other stream valleys in the region appear to have started to be formed more recently.