University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Nebraska Swine Reports

Animal Science Department

1-1-2001

Replacing Conventionally Processed Soybean Meal with Extruded/Expelled Soybean Meal in Swine Diets

Andrea M. Tucker University of Nebraska - Lincoln

Phillip S. Miller University of Nebraska - Lincoln, pmiller1@unl.edu

Austin Lewis University of Nebraska - Lincoln, alewis2@unl.edu

Duane Reese University of Nebraska - Lincoln, dreese1@unl.edu

Follow this and additional works at: http://digitalcommons.unl.edu/coopext swine



Part of the Animal Sciences Commons

Tucker, Andrea M.; Miller, Phillip S.; Lewis, Austin; and Reese, Duane, "Replacing Conventionally Processed Soybean Meal with Extruded/Expelled Soybean Meal in Swine Diets" (2001). Nebraska Swine Reports. Paper 103. http://digitalcommons.unl.edu/coopext_swine/103

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Swine Reports by an authorized administrator of Digital Commons@University of Nebraska - Lincoln.



lb/d) and the average calculated fatfree lean gain (.61 lb/d) is 15%. Therefore, if the producers were formulating their diets based on the lean gain potential of their pigs and not the actual fat-free lean gain the diets would be over-formulated for dietary crude protein. This may explain why some diets were over-formulated for dietary crude protein by as much as three percentage units. In addition, research has shown that when pigs are fed a corn-soybean meal diet with no crystalline amino acids to meet the pig's crude protein requirement the plasma urea concentration should be approximately 25 mg/dL. In many cases, the plasma urea concentrations shown in Table 1 exceed 25 mg/dL, further supporting the finding that most diets were over-formulated for crude protein.

Conclusions

Results from this on-farm study indicate that the relationship between plasma urea concentration and dietary crude protein is similar to the relation-

ship established in our research facilities. Plasma urea concentrations have the potential to be used as a means of selecting replacement boars and gilts with a high potential for lean growth. Animals would be selected for a low plasma urea concentration when fed a diet formulated to meet their dietary protein requirement. By using this technology, a producer would select animals with a more efficient utilization of dietary protein and an increase in lean muscle accretion. This technology also has the ability to help producers make management decisions on their farms. Producers could use this technology to determine if the environment, diet, facility or a management practice is limiting the lean growth potential of their pigs. These results also indicate that plasma urea concentration has the potential to be used as an indicator of the protein requirement of growing-finishing pigs. The use of plasma urea concentrations to determine the protein requirement would be less costly and time consuming than the traditional feeding and carcass analysis experiments used to identify

protein requirements for growingfinishing pigs. However, nutrient requirements for each phase of production on each operation are not the same because of differences in genetics, management, disease, and facilities between operations. Therefore, this approach may assist producers in determining the correct time point to change the nutrient density in the diet to meet the protein requirement more accurately. Also, this methodology may help producers and nutritionists clearly identify instances where excess dietary protein is provided.

A special thank-you goes out to all of the producers involved in this experiment. We appreciated your willingness to participate in this experiment, taking time to fill out the survey, and allowing us to bleed pigs on your operation. Without you this experiment would not have been possible.

¹Robert L. Fischer is a research technologist and graduate student, Phillip S. Miller is an associate professor, and Austin J. Lewis is a professor in the Department of Animal Science.

Replacing Conventionally Processed Soybean Meal with Extruded/Expelled Soybean Meal in Swine Diets

Andrea M. Tucker Phillip S. Miller Austin J. Lewis Duane E. Reese¹

Summary and Implications

The effects of extruded/expelled soybean meal (ESBM) on growth performance and carcass composition of pigs from weaning to slaughter were investigated. Two experiments were conducted. In the first experiment, weaned pigs were fed a diet containing either conventional solvent-extracted soybean meal or ESBM. Average daily gain and feed efficiency were greater

in pigs fed the control diet. The second experiment was designed to determine whether nursery diet influenced performance during the growingfinishing period. In the second experiment, half of the pigs from Experiment 1 were assigned to either a control or ESBM diet and were fed until slaughter. Average daily gain and feed efficiency of pigs fed the control diet were slightly greater than those of pigs fed the ESBM diets. Differences in performance of pigs fed the two diets were greater during the nursery phase than during the growing-finishing phase. These results support our previous research in that ESBM offers no advantage in swine growth performance

over conventional solvent-extracted soybean meal.

Introduction

Extruded/expelled soybean meal (ESBM) is produced by mechanical friction creating a high temperature for a short time period. The temperature and the time spent at a given temperature directly affects the quality and nutritional value of the product. Extruded/expelled soybean meal has the potential to be a high-quality protein and oil source if processed correctly. After extrusion, soybeans (~18% fat) are expelled (pressed) to remove

(Continued on next page)



the oil. Extruded/expelled soybean meal (the final product) has about 7% fat compared to <1% fat in conventional soybean meal. Therefore, ESBM has a greater energy content than conventional soybean meal and has the potential to be an excellent feed ingredient. Extruded/expelled soybean meal is a convenient way to include fat in swine diets for purposes such as dust reduction, in addition to its nutritional value.

In the 1998 Nebraska Swine Report, results of a study comparing conventional solvent-extracted soybean meal to ESBM fed to early-weaned pigs were presented. In that study, pigs fed ESBM grew more slowly and were less efficient than pigs fed conventional soybean meal. In contrast, recent work completed at Kansas State University (KSU) suggests that pigs fed ESBM perform similarly to pigs fed conventional solvent-extracted soybean meal. The ESBM used at KSU was from a different source than that used in the Nebraska study. An article in the 2000 Nebraska Swine Report concluded that there is considerable variation in the quality of ESBM fed to pigs and that affects its economic value.

The primary objective of this research was to investigate the effects of soybean meal type on pig performance in pigs fed either a diet containing conventional soybean meal or ESBM.

Procedures

Two experiments were conducted. Experiment 1 was a 28-day nursery trial and Experiment 2 was a 119-day growing-finishing trial.

Experiment 1

Four-hundred-eighty crossbred pigs weaned at 11 to 14 days of age (initial body weight 9.05 lb) were used. Pigs were housed in an environmentally controlled nursery with heat lamps for supplemental heat and had continuous fluorescent lighting throughout the trial. Pigs were allotted to pens with 20 pigs/pen (10 barrows and 10 gilts), and the pen was the experimental unit. Pens were allocated to either a control diet

Table 1. Composition of experiment 1 diets, % (as fed basis).

| | Phase 1 ^a | | Pha | Phase 2 ^a | |
|---------------------------------------|----------------------|-------------------|---------|----------------------|--|
| Ingredient, % | Control | ESBM ^b | Control | ESBM | |
| Corn | 39.15 | 40.40 | 50.00 | 50.65 | |
| Soybean meal (47.5% CP ^c) | 15.75 | _ | 25.75 | _ | |
| ESBM (43% CP) | _ | 16.50 | _ | 27.10 | |
| Dried whey | 27.00 | 27.00 | 14.00 | 14.00 | |
| Plasma protein | 5.00 | 5.00 | _ | _ | |
| Menhaden fishmeal | 7.00 | 7.00 | 4.00 | 4.00 | |
| Corn oil | 2.00 | _ | 2.00 | _ | |
| Dicalcium phosphate | 1.15 | 1.15 | 1.25 | 1.25 | |
| Limestone | .15 | .15 | .20 | .20 | |
| Mineral premix | .10 | .10 | .10 | .10 | |
| Vitamin premix | 1.00 | 1.00 | 1.00 | 1.00 | |
| Salt | .30 | .30 | .30 | .30 | |
| Mecadox-50 | 1.00 | 1.00 | 1.00 | 1.00 | |
| Zincoxide | .40 | .40 | .40 | .40 | |
| Calculated composition | | | | | |
| CP, % | 22.26 | 21.98 | 20.57 | 20.04 | |
| Lysine ^d ,% | 1.34 | 1.34 | 1.09 | 1.09 | |
| ME, kcal/lb ^c | 1,514 | 1,495 | 1,513 | 1,512 | |

^aPhase 1 diets fed from d 0 to 14, Phase 2 diets fed from d 14 to 28.

^dApparent digestible basis.




Figure 1. Experiment 2 pig allotment.

Table 2. Composition of experiment 2 diets, % (as fed basis).

| | Pha | se 1 ^a | Pha | ase 2 ^a | Pha | ase 3 ^a |
|---------------------------------------|---------|-------------------|---------|--------------------|---------|--------------------|
| Ingredient, % | Control | ESBM ^b | Control | ESBM | Control | ESBM |
| Corn | 68.15 | 68.65 | 76.15 | 76.90 | 80.30 | 81.59 |
| Soybean meal (47.5% CP ^c) | 27.25 | _ | 19.25 | _ | 15.25 | _ |
| ESBM (43% CP) | _ | 28.75 | _ | 20.50 | _ | 15.96 |
| Tallow | 2.00 | | 2.00 | _ | 2.00 | _ |
| Dicalcium phosphate | 1.10 | 1.10 | 1.10 | 1.10 | .95 | .95 |
| Limestone | .40 | .40 | .40 | .40 | .40 | .40 |
| Mineral premix | .10 | .10 | .10 | .10 | .10 | .10 |
| Vitamin premix | .70 | .70 | .70 | .70 | .70 | .70 |
| Salt | .30 | .30 | .30 | .30 | .30 | .30 |
| Calculated Composition | | | | | | |
| CP,% | 18.60 | 18.06 | 15.46 | 15.20 | 13.91 | 13.63 |
| Lysine ^d ,% | 1.00 | 1.00 | .78 | .78 | .67 | .67 |
| | ,547 | 1,556 | 1,549 | 1,544 | 1,552 | 1,539 |

^aPhase 1 diets fed from 27 to 110 lb (d 0 to 56), Phase 2 diets fed from 110-180 lb (d 56 to 91), Phase 3 fed diets from 180-240 lb (d 91 to 119).

^bESBM=Extruded/expelled soybean meal.


^cCP = Crude protein; ME = metabolizable energy.

^bESBM=Extruded/expelled soybean meal.


^cCP = Crude protein; ME = metabolizable energy.


dApparent digestible basis.





^aNursery Diet effect (P<.05).





^aNursery Diet effect (P<.05).

Figure 2. The response of a) average daily gain (ADG), b) average daily feed intake (ADFI), and c) ADG/ADFI to extruded/expelled soybean meal (ESBM) in Experiment 1.

(n = 12) or an experimental diet (n = 12). Pigs were given ad libitum access to feed and water throughout the 28-day feeding trial.

Diets were formulated to contain similar percentages of digestible lysine and were corn-soybean meal based containing either conventional soybean meal (Control) or ESBM (Table 1). Pigs were fed their respective treatment diet for 28 days. There were two feeding phases during Experiment 1 to meet the changing nutritional require-

ments of the pigs. Phase 1 diets were fed from day 0 to 14 (1.34% digestible lysine) and phase 2 diets were fed from day 14 to 28 (1.09% digestible lysine).

Pig and feeder weights were recorded weekly and blood samples were collected on days 0, 14, and 28 of the trial for analysis of plasma urea nitrogen.

Experiment 2

Two-hundred-forty pigs from Experiment 1 were selected immedi-

ately after the nursery trial to continue on in the grower-finisher phase of the study. We selected an equal number of pigs from each Experiment 1 diet (120 pigs fed Control and 120 pigs fed ESBM) to achieve a common initial pig body weight (27 lb) for all pens in Experiment 2. Pigs were housed in a modifiedopen-front building 10 pigs/pen (5 barrows and 5 gilts), and pen was the experimental unit. Pens were assigned either a Control diet (n = 12)or an ESBM diet (n = 12). Pigs were assigned to a pen based on the dietary treatment fed during the nursery phase such that all pigs in a pen were fed the same diet during Experiment 1. This created four possible Experiment 1-Experiment 2 diet combinations: Control-Control, Control-ESBM, ESBM-Control, and ESBM-ESBM, respectively (Figure 1).

Diets were formulated on an equal digestible lysine basis (Table 2). Experiment 2 was divided into three feeding phases. Phase 1 diets were fed from days 0 to 56 (27 to 110 lb). Phase 2 diets were fed from days 56 to 91 (110 to 180 lb), and Phase 3 diets were fed from days 91 to 119 (180 to 240 lb). The diets contained 1.00%, 0.78% and 0.67% lysine, respectively.

Pigs and feeders were weighed approximately every two weeks to make phase changes close to target weights. Blood samples were collected on days 56, 91 and 119. On day 119, backfat (BF) and longissimus muscle area (LMA) were measured using real-time ultrasound by a trained technician. Pigs were transported to a slaughter facility and TOBEC (total body electrical conductivity) measurements were recorded for each carcass.

Results

Experiment 1

Average daily gain (ADG), average daily feed intake (ADFI), and feed efficiency (ADG/ADFI) are shown in Figures 2 a, b, and c, respectively. Overall, and from days 0-14 and days 14-28, ADG was greater (P < .05) in the Control pigs than the ESBM pigs.

(Continued on next page)



There were no significant differences between the treatment groups in ADFI at any point during the nursery trial. From days 0-14, days 14-28 and days 0 to 28, ADG/ADFI was greater (P < .05) in the Control pigs than the pigs fed ESBM. On days 14 and 28, plasma urea concentrations (PUC) were greater (P < .05) for pigs fed ESBM compared to the Control group (data not shown).

Experiment 2


Average daily gain, average daily feed intake, and ADG/ADFI data are shown in Figure 3 a, b, and c, respectively. From days 0-56 and for the entire growing-finishing period (days 0 to 119), ADG was greater (P < .05) for the Control pigs than for ESBMfed pigs. From days 91-119 there was a nursery x growing-finishing diet interaction (P < .05) for ADG. Average daily feed intake was greater (P < .05) for Control pigs from days 56-91. Control pigs had a greater (P < .05) ADG/ ADFI from days 0-56. There were no significant differences in plasma urea concentration (data not shown) between the treatment groups.

Carcass data are shown in Table 3. There were no differences in BF and LMA measurements between treatments. Hot carcass weight (HCW) and total pounds of primal cuts were greater (P < .05) for Control pigs than for pigs fed ESBM.

Considering only the 120 pigs that were maintained on either the Control diet or the ESBM diet from weaning till slaughter, there was a trend (P < .10) for greater ADG and ADG/ADFI for the Control pigs vs. ESBM-fed pigs (ADG: 1.52 vs. 1.47 lb, respectively; ADG/ADFI: 1.00 vs. 0.98 , respectively).


Conclusions

During the nursery trial, pigs fed ESBM had reduced ADG and ADG/ ADFI versus pigs fed the Control diet. The greater PUC for the ESBM-fed




^aGrowing-finishing diet effect (P<.05).

^bGrowing-finishing diet x nursery diet interaction (P<.05).



^aGrowing-finishing diet effect (P<.05).

^bGrowing-finishing diet x nursery diet interaction (P<.05).



^aGrowing-finishing diet effect (P<.05).

Figure 3. The response of a) average daily gain (ADG), b) average daily feed intake (ADFI), and c) ADG/ADFI to extruded/expelled soybean meal (ESBM) in Experiment 2.

bNursery diet (P<.05).



Table 3. Effects of soybean meal type on carcass composition.

| | GF ^a Diet | | P-value |
|------------------------------------|----------------------|--------|---------|
| | Control | ESBM | GF Diet |
| HCW ^b | 178.88 | 173.81 | <.05 |
| Ham, lb | 21.52 | 20.97 | NS |
| Loin, lb | 24.65 | 24.23 | NS |
| Shoulder, lb | 25.89 | 25.32 | NS |
| Total pounds lean ^c | 90.23 | 87.40 | <.05 |
| Primal cut ^d , % | 40.28 | 40.57 | NS |
| Total lean ^e , % | 50.44 | 50.28 | NS |
| Backfat, in | .70 | .70 | NS |
| LMA ^t , in ² | 16.00 | 15.58 | NS |

 $^{{}^}aGF = {\rm growing\text{-}finishing\ diet;\ Control} = {\rm conventional\ soybean\ meal;\ ESBM} = {\rm extruded/expelled\ soybean\ meal.}$

pigs suggests that protein quality and(or) amino acid availability may be compromised in ESBM. During the growing-finishing trial, growth performance differences between the Control group

and the ESBM-fed pigs were reduced. This observation could be related to age of the pig. If the quality of the ESBM was poor (damaged protein and(or) presence of antinutritional fac-

tors), the deleterious affects would be reduced as the pig matured. Considering the whole period from weaning to finishing, there was a trend for pigs fed the Control to have a slight advantage over pigs fed ESBM.

Extruded/expelled soybean meal may be a satisfactory ingredient in swine diets when fed either during the growing-finishing period or from weaning to finishing. The variation in ESBM from different processing plants and questions about quality control and nutrient availability in ESBM need to be explored.

¹Andrea M. Tucker is a graduate student, Phillip S. Miller is an associate professor, Austin J. Lewis is a professor, and Duane E. Reese is an associate professor in the Department of Animal Science.

Economic Value of Ractopamine (PayleanTM) for Finishing Pigs

Duane E. Reese Larry L. Bitney¹

Summary and Implications

Ractopamine, a feed additive which improves feed efficiency, daily gain and several carcass characteristics recently became available to pork producers. An economic feasibility analysis on the feeding of 4.5, 9.0, and 18.0 g/ ton ractopamine to finishing pigs fed a 16% crude protein (0.82% lysine) cornsoybean meal diet from 150 to 240 lb was conducted. The analysis was performed in two stages: 1) an economic benefit for ractopamine was calculated from cost savings due to improved feed efficiency and daily gain, and 2) the amount of carcass premium needed per pig to recover the added cost of feeding ractopamine was calculated for each dietary level of ractopamine. We assumed one pound of PayleanTM,

containing 9 grams of ractopamine per pound, cost \$26. As expected, the economic benefit (considering improved feed efficiency and daily gain) of feeding ractopamine increases as corn and soybean meal prices increase. However, its use cannot be justified economically through improved feed efficiency and daily gain alone (corn =\$2.00/bu; soybean meal =\$200/ton). A producer would need to earn carcass premiums averaging \$.41, \$1.85, or \$4.97 per pig in order to recover the cost of feeding 4.5, 9.0, and 18.0 g/ton ractopamine, respectively. From the standpoint of costs and returns and assuming carcass premium is based on 10th rib backfat, it appears easier to justify feeding 9 g/ton ractopamine compared to 4.5 or 18 g/ton, because the first 9 grams of ractopamine resulted in the biggest reduction in 10th rib backfat (.09 inches), while an additional 9 g/ton (total of 18 g/ton) reduced backfat another .04 inches

only. However, if carcass premium is based on a measure of loin eye area, feeding 4.5 g/ton ractopamine may be the best choice. We conclude that a consistent carcass premium is necessary to justify feeding ractopamine economically and that producers supplement published research information on responses to feeding ractopamine with data generated on their own pigs.

Introduction

Pork producers have the opportunity to use a new feed additive, ractopamine, in finishing pig diets. Ractopamine (PayleanTM; Elanco Animal Health) belongs to a class of compounds know as beta-agonists. These compounds are similar in structure and pharmacological properties to epinephrine (adrenaline), a hormone secreted by the adrenal gland. Beta-agonists alter how nutrients that pigs (Continued on next page)

^bHCW = hot carcass weight.

^cTotal pounds lean = pounds of boneless ham, loin, shoulder, belly, and trimmings.

dPrimal cut, % = pounds of boneless ham, loin, and shoulder/HCW.

eTotal lean, % = total pounds of boneless lean/HCW.

fLMA = longissimus muscle area.

 $^{{}^{}g}NS = Not significant (P > .05).$