

2008

Myenteric neurons of the ileum that express somatostatin are a target of prion neuroinvasion in an alimentary model of sheep scrapie

David A. Schneider
U.S. Department of Agriculture

Huijun Yan
Washington State University

Lindsay M. Fry
Washington State University

Janet Alverson
U.S. Department of Agriculture

Stephen N. White
U.S. Department of Agriculture

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/zoonoticpub>

 Part of the [Veterinary Infectious Diseases Commons](#)

Schneider, David A.; Yan, Huijun; Fry, Lindsay M.; Alverson, Janet; White, Stephen N.; and O'Rourke, Katherine I., "Myenteric neurons of the ileum that express somatostatin are a target of prion neuroinvasion in an alimentary model of sheep scrapie" (2008). *Other Publications in Zoonotics and Wildlife Disease*. 102.
<http://digitalcommons.unl.edu/zoonoticpub/102>

This Article is brought to you for free and open access by the Wildlife Disease and Zoonotics at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Other Publications in Zoonotics and Wildlife Disease by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

David A. Schneider, Huijun Yan, Lindsay M. Fry, Janet Alverson, Stephen N. White, and Katherine I. O'Rourke

Myenteric neurons of the ileum that express somatostatin are a target of prion neuroinvasion in an alimentary model of sheep scrapie

David A. Schneider · Huijun Yan · Lindsay M. Fry ·
Janet Alverson · Stephen N. White ·
Katherine I. O'Rourke

Received: 26 December 2007 / Revised: 21 March 2008 / Accepted: 22 March 2008 / Published online: 22 April 2008
© Springer-Verlag 2008

Abstract Neuroinvasion of the enteric nervous system by prions is an important step in dissemination to the brain, yet very little is known about the basic process of enteric neuroinvasion. Using an alimentary model of neonatal disease transmission, neuroinvasion by scrapie prions in the ileum of lambs was detected by immunohistochemical staining for the disease-associated form of the prion protein, PrP^{Sc}. Odds ratios (OR) were determined for the frequency of PrP^{Sc} staining within enteric somata categorized by plexus location (myenteric, submucosal) and neurochemical staining (PGP 9.5, neural nitric oxide synthase, somatostatin, substance P, and vasoactive intestinal polypeptide). PrP^{Sc} was observed in $4.48 \pm 4.26\%$ of myenteric neurons and $2.57 \pm 1.82\%$ of submucosal neurons in five lambs aged 208–226 days but not in a lamb aged 138 days. The relative frequency of PrP^{Sc} within enteric somata was interdependent on plexus location and neurochemical type. Interestingly, PrP^{Sc} was observed more frequently within myenteric neurons than in submucosal neurons (PGP 9.5; OR = 1.72, 95% confidence interval = 1.21–2.44), and was observed within the myenteric plexus approximately $4\times$ (2.16–6.94) more frequently in somatostatin neurons

than in the general neural population stained by PGP 9.5. Nerve fibers stained for somatostatin were present in the mucosa and near PrP^{Sc} staining within Peyer's patches. The results suggest that somatostatin-expressing enteric neurons, with fiber projections near Peyer's patches, but with somata present in greatest proportion within the myenteric plexus, are an early target for neuroinvasion by scrapie prions and could serve an important role in neural dissemination.

Keywords Prion · Scrapie · Enteric nervous system · Somatostatin · Peyer's patches

Introduction

Prion diseases—or transmissible spongiform encephalopathies (TSEs)—are a group of slowly progressive, fatal neurologic diseases caused by unconventional agents called prions [39]. Although some debate persists regarding the exact nature of prions, disease is critically associated with the conversion of cellular host prion protein (PrP^C) to TSE-associated isoforms (PrP^{TSE}) [7, 38]. Natural prion disease occurs in humans (Creutzfeldt–Jakob disease and others) and in several types of animals significant to agriculture (scrapie of sheep and goats and bovine spongiform encephalopathy of domestic cattle) and to wildlife management (chronic wasting disease of deer and elk).

Disease transmission commonly occurs via the alimentary tract but specific knowledge about how prions disseminate from the gut to the brain is incomplete (see reviews by Mabbot and MacPherson [28] and by Beekes and McBride [5]). Neuroinvasion occurs in the periphery and results in dissemination to the brain via autonomic nerves; in particular, parasympathetic and sympathetic efferent nerves projecting to the gut are thought to be common pathways.

D. A. Schneider (✉) · S. N. White · K. I. O'Rourke
Animal Disease Research Unit, Agricultural Research Service,
US Department of Agriculture, 3003 ADBF, WSU,
PO Box 646630, Pullman, WA 99164-6630, USA
e-mail: das@vetmed.wsu.edu

H. Yan · L. M. Fry
Department of Veterinary Microbiology and Pathology,
Washington State University, Pullman, WA 99164, USA

J. Alverson
Animal and Plant Health Inspection Service,
Veterinary Services, US Department of Agriculture,
Helena, MT 59601, USA

However, the gut is most extensively innervated by a third division of the autonomic nervous system, the enteric nervous system (see reviews by Brookes and Costa [6], and Furness [15]). It is estimated in sheep that the small intestine alone contains more than 30 million myenteric and 50 million submucosal neurons, a total population that rivals that in the spinal cord [16].

The enteric nervous system is a functionally complete, ganglionated nerve network that is organized into a myenteric plexus and a submucosal plexus, out of which all layers of the gut wall are innervated. With some variation amongst species, there is a division of labor such that most neurons which control gut motility are located within the myenteric plexus, and most neurons which control mucosal blood flow and secretion are located within the submucosal plexus. Coordination is achieved through inter-plexus enteric projections, extrinsic autonomic projections, and viscerofugal projections from the enteric nervous system to prevertebral ganglia.

The enteric nervous system includes many different neuron classes, which have been classified not only by plexus location and region of the alimentary tract, but also by somato-dendritic morphology, electrophysiology, neurochemistry, fiber projection, and physiologic action on a target tissue. Major functional classes of enteric neuron include intrinsic primary afferent neurons and several types each of interneurons, smooth muscle motor neurons, secretomotor neurons, and vasomotor neurons. Though not fully characterized, even local gut immune function is modulated by the enteric nervous system, in part through peptidergic innervation of Peyer's patches [50]. The enteric nervous system is thus a complex neural system that is intimately associated with all layers of the gut, including the mucosal layer through which prions gain entry into the host, and Peyer's patches in which prion replication occurs early in disease progression.

Given that extrinsic autonomic nerves synapse with enteric neurons [15], that enteric neurons express PrP^c [40] and innervate all layers of the gut, and that scrapie-associated prion protein (PrP^{Sc}) is first detected in sheep within the enteric nervous system [3, 19, 46], it is hypothesized that neuroinvasion of the enteric nervous system is important for the dissemination of scrapie prions to the brain via autonomic nerves.

Many observations suggest that neurons projecting into sites of prion replication (Peyer's patches) should be targets for invasion by prions [5, 28], but a more diverse group of neurons (with projections into the mucosa or around vessels) may be exposed immediately following uptake of prions from the gut lumen [25]. The purpose of this study was to identify enteric neural targets of scrapie prions early in disease progression in a natural disease model in lambs. The results demonstrate that following

neonatal alimentary disease transmission, frequency of PrP^{Sc} detection within ileal enteric neurons is disproportionately greater for myenteric neurons expressing the peptide transmitter, somatostatin. The identification of enteric neural targets of scrapie prions early in disease progression contributes significantly in understanding the factors involved in peripheral neuroinvasion and neural dissemination of prions.

Materials and methods

Oral inoculation model

All animal experiments were approved by the Washington State University Institutional Animal Care and Use Committee. The study included seven Montadale or MontadaleX Suffolk lambs born to Montadale ewes. All lambs were VRQ/VRQ (see "Genotyping" section below), and on the day of birth were fed half a cotyledon from a placenta shed previously from an unrelated ARQ/VRQ ewe clinically affected with scrapie and with staining for PrP^{Sc} in sections of medulla at the level of the obex. Cotyledons were stored at -20°C until used.

Peripheral prion replication in lambs was assessed antemortem by third eyelid [31] or rectal [13, 18] biopsy obtained at 4–6 months of age. Lambs were humanely euthanized at about 7 months of age by intravenous injection of sodium pentobarbital (Fatal-Plus; Vortech Pharmaceuticals, Dearborn, MI). Tissues collected immediately postmortem included medulla, thoracolumbar spinal cord, celiacomesenteric ganglia, ileocecal lymph node, ileocecal junction, and a 30 cm length of ileum ending within ~ 3 cm of the ileocecal junction. All tissues were fixed by immersion in 10% neutral-buffered formalin; the ileum was also gently distended with formalin.

Genotyping

Approximately 100 ng DNA was used as a template for amplification of the open reading frame of PRNP using the following primer pair: 5'-GGCATTGATGCTGACACC-3' and 5'-TACAGGGCTGCAGGTAGAC-3', corresponding to nucleotide positions 22234–22252 and 23106–23125 of GenBank accession no. U67922. Reactions were performed in a final volume of 90 μL under the following conditions: 1 \times Qiagen PCR buffer, 2.5 mM MgCl₂ and 2.5 U Taq polymerase (201225, TaqPCR Core kit; Qiagen, Valencia, CA). Amplification was performed under a temperature regime of 95 $^{\circ}\text{C}$ for 5 min, 62 $^{\circ}\text{C}$ for 30 s, and 72 $^{\circ}\text{C}$ for 59 s for one cycle; 95 $^{\circ}\text{C}$ for 30 s, 62 $^{\circ}\text{C}$ for 30 s, and 72 $^{\circ}\text{C}$ for 59 s for 30 cycles; and 95 $^{\circ}\text{C}$ for 30 s, 62 $^{\circ}\text{C}$ for 30 s and 72 $^{\circ}\text{C}$ for 7 min for one cycle. Amplified products were

sequenced by standard dideoxynucleotide analysis (Amplicon Express, Pullman, WA) using primer pair 5'-CTGGGGTC AAGGTGGTAGCC-3' and 5'-GGTGGTGACTGTGTGTT GCTTGA-3', corresponding to nucleotide positions 22553–22573 and 22838–22860 of GenBank accession no. U67922. Genotypes are reported as the deduced amino acid sequences at codons 136 (alanine, A; valine, V), 154 (arginine, R; histidine, H), and 171 (glutamine, Q; arginine, R) of the diploid genotype.

PrP^{Sc} detection by ELISA

A commercial test developed for lymphoid tissue (HerdChek Chronic Wasting Disease [CWD] Antigen Test kit; IDEXX, Westbrook, Maine) was used for detection of PrP^{Sc} in unfixed cotyledons as previously described [2]. This test is used for the postmortem detection of PrP^{Sc} in white-tailed deer retropharyngeal lymph node tissue but is also suitable for the detection of PrP^{Sc} in sheep cotyledons; the kit utilizes a PrP^{Sc} ligand immobilized on the surface of the CWD-antigen capture plate and monoclonal antibodies that recognize the ovine PrP protein. Briefly, 300 mg of cotyledon was homogenized twice in a FastPrep instrument (Qbiogene, Carlsbad, CA) for 30 s each at maximum speed (6.5 m/s) in a disruption tube containing ceramic beads and 0.9 mL of distilled, deionized water. The ELISA was then conducted according to the manufacturer's protocols. Interpretation of samples analyzed by ELISA was performed by comparing the A_{450} of the cotyledon samples with that of negative- and positive-control samples supplied with the kit (reference wavelength = 620 nm). Samples were considered positive if the mean A_{450} values were greater than or equal to the cutoff value as calculated by the test manufacturer.

Immunohistochemistry (IHC)

All formalin-fixed tissues were placed in cassettes, incubated for 60 min in 98% formic acid, washed extensively in distilled water, and re-equilibrated in formalin for 24 h, prior to paraffin embedding. Thin (3–5 μ m) paraffin sections were mounted on Superfrost Plus slides, air dried overnight, and then baked at 57°C for 100 min. IHC staining was carried out using a BenchMark automated processor (Ventana Medical Systems, Tucson, AZ) with modifications to the technique previously described [31, 43]. Antigen retrieval was achieved by 60–90 min heated incubation in CC1 buffer (Ventana Medical Systems). Scrapie-associated prion protein (PrP^{Sc}) was detected using a cocktail of well-characterized monoclonal antibodies (F89/160.1.5 and F99/97.6.1; Table 1) applied for 32 min at 37°C. Evaluation of PrP^{Sc} dissemination was determined using Basic AEC Detection Kit (760–020; Ventana Medical Systems). Slides were then counterstained using Bluing Reagent and Hematoxylin kits (Ventana Medical Systems) and coverslipped in aqueous mounting media (S3025; Dako, Carpinteria, CA). In dual fluorescence labeling studies, an antibody directed toward a neurochemical of interest (Table 1) was included in the cocktail of anti-PrP^{Sc} primary antibodies. Relevant combinations of fluorescently-labeled secondary antibodies were applied for 20 min at 37°C (Table 2). Fluorescently labeled slides were coverslipped using Prolong Gold antifade reagent (P36930; Molecular Probes, Eugene, OR).

Using these techniques, PrP^{Sc} staining could be detected in the obex, lymph nodes, and ileum from a clinically affected ewe but not from an unexposed, clinically normal ewe. IHC staining for all neural and glial marker antibodies was dependent upon inclusion of the primary antibody and an appropriate fluorophore-conjugated secondary antibody.

Table 1 Primary antibodies

Target	Name	Species and isotype	Working dilution/concentration	Source (reference)
PrP ^{Sc}	F89/160.1.5	Mouse IgG ₁	5 μ g/mL	Dr. Katherine O'Rourke, USDA-ARS, Pullman, WA [30, 31]
	F99/97.6.1	Mouse IgG ₁	5 μ g/mL	Dr. Katherine O'Rourke, USDA-ARS, Pullman, WA [31]
Glial fibrillary acid protein	GFAP	Rabbit IgG	1:400	Z0334; Dako, Carpinteria, CA [24, 45]
Elav protein (human neuronal protein, Hu)	Hu	Mouse IgG _{2b}	1:200	A-21271; Molecular Probes, Eugene, OR [8]
Neural nitric oxide synthase	NOS	Rabbit IgG	1:200	SA227; Biomol Research Laboratories, Plymouth Meeting, PA [29]
Protein gene product 9.5	PGP	Rabbit IgG	1:200	Z5116; Dako
Somatostatin	SOM	Rabbit IgG	1:1,600	20067; ImmunoStar
Substance P	SP	Rat IgG _{2a}	1:50	MAB356; Chemicon, Temecula, CA
Vasoactive intestinal polypeptide	VIP	Guinea pig IgG	1:800	T-5030 (GHC7161); Peninsula Laboratories [32]

Table 2 Secondary antibodies used in fluorescence IHC

Target visualized	Antibody	Source
PrP ^{Sc} , Hu	Cy3 AffiniPure donkey anti-mouse IgG	715-165-150; Jackson ImmunoResearch Laboratories, West Grove, PA
SP	Cy3 AffiniPure donkey anti-rat IgG	712-165-150; Jackson ImmunoResearch Laboratories
PrP ^{Sc}	Alexa Fluor 488 goat anti-mouse IgG	A-11029; Molecular Probes, Eugene, OR
GFAP, NOS, PGP, SOM	Alexa Fluor 488 goat anti-rabbit IgG	A-11034; Molecular Probes, Eugene, OR
VIP	Alexa Fluor 488 goat anti-guinea pig IgG	A-11073; Molecular Probes, Eugene, OR

Microscopy

Slides were viewed and photographed using an Axio Imager.M1 microscope (Carl Zeiss Microimaging, Thornwood, NY) equipped with an LED illuminator for bright field microscopy and an X-Cite 120 FI Illuminating system (EXFO Photonic Solutions, Mississauga, Ontario, Canada) for epi-fluorescence microscopy. Cy3 and Alexa Fluor 555 fluorescence were observed using the Carl Zeiss filter set 00 (excites 530–585 nm, dichroic mirror 600, 615 nm long-pass filter), and Alexa Fluor 488 fluorescence was observed using the Carl Zeiss filter set 38HE (excites 450–490 nm, dichroic mirror 495, 500–550 nm band-pass filter). The microscope was equipped with an AxioCam MRc5 digital camera (Carl Zeiss Microimaging) connected to a computer workstation running AxioVision 4.5 imaging software (Carl Zeiss Microimaging). For presentation in Figs. 4 and 7, maximum intensity projections were produced from sequentially scanned z-stack series (0.5–1 μ m intervals) using an LSM 510 META laser scanning microscope (Carl Zeiss Microimaging) equipped with 488 and 543 nm lasers (pinholes set to 1 Airy unit).

Statistical methods

All analyses were performed using SAS for Windows (ver. 9.1.3, service pack 4; SAS Institute, Cary, NC). The data set was categorized by animal, enteric plexus location (myenteric or submucosal) and by neurochemical type (PGP, nNOS, SOM, SP, or VIP; abbreviations defined in Table 1). The total number of neurons (defined by somata staining) and the number of those colocalized with PrP^{Sc} were recorded for each neurochemical type. The total association of PrP^{Sc} colocalization in neurochemically-defined (NOS⁺, SOM⁺, SP⁺, or VIP⁺) somata compared to colocalization in the total neural population (PGP⁺) was determined for each animal by Fisher's exact testing. The null hypothesis of random detection of PrP^{Sc} amongst neurochemically-defined enteric neurons was assessed by conditional logistic regression using a generalized linear mixed model, the GLIMMIX procedure (rel. June 2006; SAS Institute), and the fixed effects of plexus location, neurochemical type, and the interaction

of these two effects. The regression was conditioned by including the random effect of subjects (lambs) and adjusted for overdispersion by including the random effect of residuals. Holm-adjusted *P* values (P_{adj}) were produced using the method of Bonferroni in logical step-down fashion [23]. Odds ratios (ORs) and corresponding upper and lower CL_{adj} s are the exponentiation of the least squares means parameter and CL_{adj} s estimates, respectively. Least squares means parameter and Holm-adjusted 95% confidence limit (CL_{adj}) estimates were produced for two "families" of post-hoc comparisons of interest; family-wise error rate, $\alpha = 0.05$.


Results

Alimentary transmission and staging disease progression

Cotyledons randomly selected from the pool of cotyledons used for alimentary inoculation had PrP^{Sc} staining by IHC and a PrP^{Sc} ELISA titer between 1:64 and 1:256. Ewes giving birth to the lambs were not a likely source of PrP^{Sc} since PrP^{Sc} staining was neither detected by IHC in antemortem peripheral lymphoid tissues (third eyelid and rectal mucosa) and postmortem obex, nor by IHC and western blot in shed cotyledons randomly sampled at the birth of these lambs.

Tissues from a lamb euthanized before 2 weeks of age due to an unrelated illness were not collected. In the remaining six lambs, PrP^{Sc} was detected antemortem in rectal mucosa-associated lymphoid tissue between 124–219 days of age, and in ileal Peyer's patches and ileocecal lymph node collected postmortem between 138–226 days of age (Fig. 1b, d, f; scrapie-negative control tissues shown in Fig. 1a, c, e). PrP^{Sc} was not detected in the central nervous system (medulla and intermediolateral column of T8–L3 spinal cord segments) or celiacomesenteric ganglia (138–226 days of age), except in one lamb (226 days of age) in which sparse punctate staining in celiacomesenteric ganglia appeared to be located mostly in satellite cells (Fig. 2a–d). PrP^{Sc} was detected in the enteric nervous system of the ileum in five lambs aged 208–226 days, but not in a lamb aged 138 days (examples shown in Fig. 3). The appearance of PrP^{Sc} staining within enteric somata was generally punctate (Fig. 3d–f, Fig. 4a–c) but sometimes

Fig. 1 PrP^{Sc} staining (red) was not observed in various lymphoid tissues from a scrapie-negative control ewe (**a, c, e**) but was readily detected in lymphoid tissues from a lamb orally inoculated with scrapie prions at birth (**b, d, f**). Lymphoid tissues shown are from: **a, b** rectal mucosa-associated lymphoid tissue (*RMALT*); **c, d** ileal Peyer's patch; **e, f** ileocecal lymph node. Bars 100 μ m



globular when near cell margins. PrP^{Sc} was also occasionally observed within enteric glia (Fig. 4d–f).

Frequency of PrP^{Sc} detection within defined populations of enteric neurons

The occurrence of PrP^{Sc} in PGP⁺ somata was tallied for a total of 10,841 myenteric neurons and 19,495 submucosal neurons. The frequency of PrP^{Sc} detection in PGP⁺ somata ranged from 1.21 to 11.67% (mean and standard deviation: $4.48 \pm 4.26\%$) in myenteric neurons, and from 1.01 to 5.68% ($2.57 \pm 1.82\%$) in submucosal neurons ($n = 5$ lambs). Frequencies of PrP^{Sc} detection in enteric neurons defined by plexus location and neurochemical type are shown for each lamb in Fig. 5. The pattern of PrP^{Sc} detection frequency amongst eight enteric neural subgroups relative to that in PGP⁺ somata varied between lambs except for a consistently increased detection frequency in SOM⁺ neurons of the myenteric plexus (Breslow–Day test for homogeneity: Chi-square = 1.7067, $P = 0.7895$).

The significance and strength of the association of PrP^{Sc} detection frequency with subpopulations of enteric neurons was measured by odds ratio estimates for somata defined by plexus location and neurochemical type and analyzed by conditional logistic regression. The odds of PrP^{Sc} detection were significantly dependent upon plexus location ($F = 4.66$, $P = 0.0376$), neurochemical type ($F = 20.59$, $P < 0.0001$), and the interaction of these main effects ($F = 6.66$, $P = 0.0004$). The detailed post-hoc analyses of this finding are summarized below and in Fig. 6.

PrP^{Sc} was detected more frequently in the myenteric plexus versus the submucosal plexus (Fig. 6a) for the general population of neurons (PGP; 1.72, 1.21–2.44) and for the subpopulation of SOM⁺ neurons (3.92, 2.09–7.35). Relative to local PGP⁺ neural populations, the frequency of PrP^{Sc} detection was significantly increased for only SOM⁺ neurons in the myenteric plexus (3.87, 2.16–6.94; Fig. 6b) and in the submucosal plexus (1.70, 1.02–2.82; Fig. 6c). Significantly lower frequencies of detection were associated with SP⁺ neurons in the myenteric plexus (0.21, 0.05–0.87;

Fig. 2 Detection of PrP^{Sc} staining (*red*) in the celiacomesenteric ganglia from a lamb aged 226 days following oral inoculation with scrapie prions at birth. **a** PrP^{Sc} detected in a satellite cell. The *box* depicts the region shown at higher magnification in **b**. **c** PrP^{Sc} at the margin of this ganglion cell may also be located in a satellite cell in which only a part of the nucleus is present in this section. The *box* depicts the region shown at higher magnification in **d**. *Bars* 50 μ m (**a**, **c**), 5 μ m (**b**, **d**)





Fig. 3 Examples of PrP^{Sc} staining (*red*) detected in the ileum of two lambs orally inoculated with scrapie prions at birth. **a–c** In a lamb aged 138 days, PrP^{Sc} staining is present within a Peyer's patch follicle (*lower half* of image) but not within adjacent submucosal neurons (*arrowheads* denote ganglion limits). **d–f** PrP^{Sc} staining in two submucosal neurons (*arrows*) of a lamb aged 216 days. PrP^{Sc} staining was also

present in a Peyer's patch follicle located out of frame *above* this ganglion. The *background* (*green*) column is included to provide tissue contrast. PrP^{Sc} staining remains *red* in the *merged* column of this single labeling experiment, whereas autofluorescence appears *bright yellow*. Images are maximum intensity projections from 6 μ m z-stacks. *Bars* 20 μ m




Fig. 4 Initial detection of PrP^{Sc} staining (red) in the enteric nervous system of the ileum from a lamb aged 216 days following oral inoculation with scrapie prions at birth. **a–c** PrP^{Sc} detected within (arrow), and outside of (arrowheads), PGP⁺ enteric neurons (green). **d–f** PrP^{Sc}

detected within (arrows) GFAP⁺ enteric glia (green). Images are maximum intensity projections from 2.5 μm (**a–c**) and 1.5 μm (**d–f**) confocal z-stacks. Abbreviations defined in Table 1




Fig. 5 Frequency of detecting PrP^{Sc} staining within subpopulations of enteric neuron in the ileum of lambs early in disease progression following oral inoculation with scrapie prions at birth. Note the consistent increase in the frequency of PrP^{Sc} detection within myenteric plexus SOM⁺ neurons as compared to that in the general myenteric neuron population (-PGP-). Each bar color represents data from a lamb (lambs 1–5) at approximately 7 months of age. Frequency of PrP^{Sc} detection is given as the percent (%), vertical axis within each type of neuron per lamb. Abbreviations are defined in Table 1. Arrows indicate the direction of significant differences (Fischer’s exact test; $P < 0.05$)

Fig. 6b) and VIP⁺ neurons in the submucosal plexus (0.23, 0.07–0.76; Fig. 6c).

SOM⁺ nerve fibers in the ileum

SOM⁺ nerve fibers were frequently observed projecting between mucosal epithelium and Peyer’s patches with

PrP^{Sc} staining (Fig. 7). Similar to a previous description [49], staining for SOM was also occasionally observed in “open-type” epithelial cells of the ileal mucosa (not shown).

Discussion

The process of neuroinvasion of the enteric nervous system by prions was studied in the ileum of sheep, the first peripheral region in which neuroinvasion occurs in naturally acquired disease [3, 12, 47]. We determined that the relative frequency of PrP^{Sc} staining, used as a surrogate marker [1, 26] for scrapie prions within enteric neurons, is dependent upon that neuron’s plexus location and neurochemical type. In particular, SOM⁺ myenteric neurons were identified as a consistent target for neuroinvasion early in disease progression. From the evidence, we postulate that a significant subgroup of SOM⁺ neurons innervate compartments of Peyer’s patches where efficient neuroinvasion is thought to occur, and represent an efficient pathway for neural dissemination within the small intestine. We expect that prion invasion of neurons that do not convert PrP^c into PrP^{Sc} that is readily detectable by immunohistochemistry [4], should they exist within the enteric nervous system, would be underrepresented in the present study.




Fig. 6 The relative frequency of detecting PrP^{Sc} staining within ileal enteric neurons of lambs is interdependent on plexus location and neurochemical type early in disease progression. **a** The frequency of PrP^{Sc} detection was greater in the myenteric versus submucosal plexus for the general (PGP) neural population and for the subpopulation of SOM⁺ neurons. **b, c** SOM⁺ neurons were the only subpopulation studied that had an increased frequency of PrP^{Sc} detection in either plexus (**b** myenteric, **c** submucosal), relative to that in the respective general neural population. The difference between an odds ratio (OR) and 1 (dotted vertical lines) indicates the relative strength of the association with the frequency of PrP^{Sc} detection. ORs significantly greater or less than 1 (depicted in red; $P_{adj} < 0.05$) indicate associations with increased or decreased frequency, respectively, of PrP^{Sc} detection. CL_{adj} Holm-adjusted confidence limits, P_{adj} Holm-adjusted P values, PrP^{Sc} the number of neurons in which PrP^{Sc} was detected, MYP myenteric plexus, SMP submucosal plexus, Total the total number of neurons counted. See Table 1 for abbreviations of neurochemical types. y -axis First column, x -axis \log_{10}

SOM⁺ enteric neurons are targets of peripheral neuroinvasion by scrapie prions

There is consensus that transport of prions across gut epithelium and exposure of mucosal nerve fibers are processes relevant to peripheral neuroinvasion and neural dissemination [5, 26, 28, 36, 47]. Considering that the greatest proportion of mucosa-projecting neurons typically occurs in the submucosal plexus of large mammalian species [44], it

is surprising that PrP^{Sc} detection early in disease was most frequent in neurons located in the myenteric plexus (Fig. 6a, PGP), and was relatively infrequent in a major type of submucosal neuron (Fig. 6c, VIP). Further, PrP^{Sc} detection in SOM⁺ neurons of the myenteric plexus was very consistent and occurred at nearly 4× greater frequency (Fig. 6b).

SOM⁺ enteric neurons of the small intestine generally belong to one of two classes that differ in function and projection: secretomotor neurons that project to the mucosal layer, and descending interneurons that synapse with other enteric neurons [6]. Determining the projections of SOM⁺ enteric nerve fibers in sheep is thus particularly germane to locating a compartment in which efficient neuroinvasion occurs. Given the present data, it is conceivable that SOM⁺ neurons targeted by scrapie prions are distributed in greatest proportion within myenteric ganglia but which, nevertheless, project to mucosal compartments in which efficient neuroinvasion occurs early in disease progression.

Implications of SOM⁺ secretomotor neurons targeted by prions

Jeffrey et al. [25] observed, soon after the injection of infectious material into the lumen of the small intestine of lambs, that PrP^{Sc} staining was widespread within mucosal villi. Mucosal villi are highly innervated by enteric neurons, including fibers of secretomotor neurons, which travel close to the epithelium to affect its function [21, 22, 41]. Fibers of numerous mucosa-projecting neurons, especially secretomotor neurons, are thus potentially exposed upon mucosal transportation of prion and prior to replication of prion within Peyer's patches.

In sheep, mucosal nerve fibers include those that express VIP, SP, NOS [8, 27, 51], and also SOM (Fig. 7). Furthermore, the secretomotor action of VIP, SP, and SOM is typically preserved within large mammalian species [44]. Therefore, it is not surprising that PrP^{Sc} was detected in each of these neurochemical types of submucosal neuron early in disease progression following alimentary transmission in sheep. That PrP^{Sc} was detected as much as 4× more frequently in SOM⁺ neurons, however, may indicate that efficient enteric neuroinvasion occurs in a mucosal compartment uniquely innervated by SOM⁺ neurons. Evidence implicates such a compartment is associated with Peyer's patches.

Neuroinvasion occurring via nerve fiber projections to Peyer's patches is consistent with the typical detection of PrP^{Sc} first within Peyer's patches and then nearby ganglia, and with close associations between fibers and PrP^{Sc}-laden cells [3, 20]. Recent evidence in mice further demonstrates that efficient neuroinvasion following alimentary transmission is dependent upon follicular dendritic cells [17], a cell type




Fig. 7 Nerve fibers stained for SOM in the mucosal layer of the ileum from a lamb aged 208 days and orally inoculated with scrapie prions at birth. **a–c, d–f** Two examples of SOM⁺ (green) nerve fibers projecting between the mucosal epithelium (EPI) and a Peyer's patch (PP) with

PrP^{Sc}-staining (red). Note the relatively close association of SOM⁺ fibers with PrP^{Sc} staining (boxes). Images are maximum intensity projections from 6 μm (**a–c**) and 2 μm (**d–f**) confocal z-stacks. SOM somatostatin

associated with nerve fibers in the interfollicular zone and dome of Peyer's patches [11]. Although SOM⁺ Peyer's patch fibers are yet to be confirmed in mice [10, 48], these have been observed in cats [14], and now also in sheep (Fig. 7). It is thus conceivable that in sheep, efficient neuroinvasion near Peyer's patches is dependent upon proximity of SOM⁺ nerve fibers to PrP^{Sc}-laden immune cells or exosomes [35, 37].

Implications of SOM⁺ descending interneurons targeted by prions

A subgroup of descending myenteric interneurons of the small intestine expresses SOM [6]. In the present study in sheep, prion targeting of SOM⁺ descending interneurons would imply neuroinvasion early in disease occurred within ganglia, in contrast to that occurring outside of ganglia via nerve fibers. Nerve endings of extrinsic nerves are unlikely to have been a significant source of prion within enteric ganglia since PrP^{Sc} was first detected in enteric neurons. Prions might alternatively enter ganglia from a non-neural

source, such as blood. However, if blood-borne prion were relevant to enteric neuroinvasion early in disease, then it would be reasonable to expect relatively higher frequency PrP^{Sc} detection in vasomotor neurons.

Vasomotor neurons have not been extensively studied in large mammalian species but probably include subgroups of VIP⁺ and SP⁺ neurons [44, 51]. The relative frequency of PrP^{Sc} detection in VIP⁺ and SP⁺ neurons, however, was not increased in either plexus (Fig. 6b, c). Therefore, simple prion entry from a non-neural source such as blood does not likely account for the higher frequency of PrP^{Sc} detection within SOM⁺ neurons. Regardless, detection of PrP^{Sc} in SOM⁺ descending interneurons early in disease progression has implications for neural dissemination.

SOM⁺ descending interneurons form a synaptic network that may descend the length of the small intestine—from duodenum to ileum, at least in guinea pigs [9, 33, 34, 42]. If this also occurs in sheep, then prion dissemination through SOM⁺ descending interneurons might account for the observed oral spread of prion from the ileum to duodenum [3, 47].

In summary, following neonatal alimentary transmission in sheep, immunohistochemical detection of PrP^{Sc} (as a surrogate marker for neuroinvasion by scrapie prions) in ileal enteric neurons was interdependent on the neuron's plexus location and neurochemical type. Specifically, SOM⁺ neurons, especially those located in the myenteric plexus, were found to be targets of neuroinvasion early in disease progression. These results are consistent with the hypothesis that efficient neuroinvasion occurs near Peyer's patches in compartments innervated by SOM⁺ neurons. In addition, neuroinvasion of SOM⁺ enteric neurons early in disease progression could explain the neural dissemination observed within the small intestine of sheep with naturally acquired disease.

Acknowledgments We thank Tom Truscott for histological assistance; Duane Chandler, Alicia Ewing, Jim Reynolds, Caitlin Rizzo, and Ira Mickelsen for assistance with animal care and necropsy; and Bruce Mackey (USDA, Albany, CA) for statistical consultation. This work was paid for by CRIS 5348-32000-019-00D from the Agricultural Research Service, US Department of Agriculture. Lindsay Fry was supported by a summer research stipend from the Research Scholars Program, College of Veterinary Medicine, Washington State University. Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the US Department of Agriculture.

References

- Aguzzi A, Heikenwalder M, Miele G (2004) Progress and problems in the biology, diagnostics, and therapeutics of prion diseases. *J Clin Invest* 114:153–160
- Alverson J, O'Rourke KI, Baszler TV (2006) PrP^{Sc} accumulation in fetal cotyledons of scrapie-resistant lambs is influenced by fetus location in the uterus. *J Gen Virol* 87:1035–1041
- Andreoletti O, Berthon P, Marc D, Sarradin P, Grosclaude J, van Keulen L, Schelcher F, Elsen JM, Lantier F (2000) Early accumulation of PrP(Sc) in gut-associated lymphoid and nervous tissues of susceptible sheep from a Romanov flock with natural scrapie. *J Gen Virol* 81:3115–3126
- Barron RM, Campbell SL, King D, Bellon A, Chapman KE, Williamson RA, Manson JC (2007) High titers of transmissible spongiform encephalopathy infectivity associated with extremely low levels of PrP^{Sc} in vivo. *J Biol Chem* 282:35878–35886
- Beekes M, McBride PA (2007) The spread of prions through the body in naturally acquired transmissible spongiform encephalopathies. *FEBS J* 274:588–605
- Brookes SJH, Costa M (2002) Cellular organization of the mammalian enteric nervous system. In: Brookes SJH, Costa M (eds) *Innervation of the gastrointestinal tract*. Taylor and Francis, New York, pp 393–468
- Bueler H, Aguzzi A, Sailer A, Greiner RA, Autenried P, Aguet M, Weissmann C (1993) Mice devoid of PrP are resistant to scrapie. *Cell* 73:1339–1347
- Chiocchetti R, Grandis A, Bombardi C, Lucchi ML, Dal Lago DT, Bortolami R, Furness JB (2006) Extrinsic and intrinsic sources of calcitonin gene-related peptide immunoreactivity in the lamb ileum: a morphometric and neurochemical investigation. *Cell Tissue Res* 323:183–196
- Costa M, Furness JB, Smith IJ, Davies B, Oliver J (1980) An immunohistochemical study of the projections of somatostatin-containing neurons in the guinea-pig intestine. *Neuroscience* 5:841–852
- De Jonge F, Van Nassauw L, De Man JG, De Winter BY, Van Meir F, Depoortere I, Peeters TL, Pelckmans PA, Van Marck E, Timmermans JP (2003) Effects of *Schistosoma mansoni* infection on somatostatin and somatostatin receptor 2A expression in mouse ileum. *Neurogastroenterol Motil* 15:149–159
- Defaweux V, Dorban G, Demonceau C, Piret J, Jolios O, Thellin O, Thielen C, Heinen E, Antoine N (2005) Interfaces between dendritic cells, other immune cells, and nerve fibres in mouse Peyer's patches: potential sites for neuroinvasion in prion diseases. *Microsc Res Tech* 66:1–9
- Ersdal C, Ulvund MJ, Espenes A, Benestad SL, Sarradin P, Landsverk T (2005) Mapping PrP^{Sc} propagation in experimental and natural scrapie in sheep with different PrP genotypes. *Vet Pathol* 42:258–274
- Espenes A, Press CM, Landsverk T, Tranulis MA, Aleksandersen M, Gunnes G, Benestad SL, Fuglesteit R, Ulvund MJ (2006) Detection of PrP(Sc) in rectal biopsy and necropsy samples from sheep with experimental scrapie. *J Comp Pathol* 134:115–125
- Feher E, Fodor M, Burnstock G (1992) Distribution of somatostatin-immunoreactive nerve fibres in Peyer's patches. *Gut* 33:1195–1198
- Furness JB (2006) Structure of the enteric nervous system. In: Furness JB (ed) *The enteric nervous system*. Blackwell, Malden, pp 1–28
- Gabella G (1987) The number of neurons in the small intestine of mice, guinea-pigs and sheep. *Neuroscience* 22:737–752
- Glaysheer BR, Mabbott NA (2007) Role of the GALT in scrapie agent neuroinvasion from the intestine. *J Immunol* 178:3757–3766
- Gonzalez L, Jeffrey M, Siso S, Martin S, Bellworthy SJ, Stack MJ, Chaplin MJ, Davis L, Dagleish MP, Reid HW (2005) Diagnosis of preclinical scrapie in samples of rectal mucosa. *Vet Rec* 156:846–847
- Heggebo R, Press CM, Gunnes G, Lie KI, Tranulis MA, Ulvund M, Groschup MH, Landsverk T (2000) Distribution of prion protein in the ileal Peyer's patch of scrapie-free lambs and lambs naturally and experimentally exposed to the scrapie agent. *J Gen Virol* 81:2327–2337
- Heggebo R, Gonzalez L, Press CM, Gunnes G, Espenes A, Jeffrey M (2003) Disease-associated PrP in the enteric nervous system of scrapie-affected Suffolk sheep. *J Gen Virol* 84:1327–1338
- Hens J, Schrodli F, Brehmer A, Adriaensens D, Neuhuber W, Scheuermann DW, Schemann M, Timmermans JP (2000) Mucosal projections of enteric neurons in the porcine small intestine. *J Comp Neurol* 421:429–436
- Hens J, Vanderwinden JM, De Laet MH, Scheuermann DW, Timmermans JP (2001) Morphological and neurochemical identification of enteric neurones with mucosal projections in the human small intestine. *J Neurochem* 76:464–471
- Holm S (1979) A simple sequentially rejective multiple test procedure. *Scand J Stat* 6:65–70
- Hortells P, Monzon M, Monleon E, Acin C, Vargas A, Bolea R, Lujan L, Badiola JJ (2006) Pathological findings in retina and visual pathways associated to natural Scrapie in sheep. *Brain Res* 1108:188–194
- Jeffrey M, Gonzalez L, Espenes A, Press CM, Martin S, Chaplin M, Davis L, Landsverk T, MacAldowie C, Eaton S, McGovern G (2006) Transportation of prion protein across the intestinal mucosa of scrapie-susceptible and scrapie-resistant sheep. *J Pathol* 209:4–14
- Jeffrey M, Gonzalez L (2007) Classical sheep transmissible spongiform encephalopathies: pathogenesis, pathological phenotypes and clinical disease. *Neuropathol Appl Neurobiol* 33:373–394
- Lalatta-Costerbosa G, Mazzoni M, Clavenzani P, Di Guardo G, Mazzuoli G, Marruchella G, De Grossi L, Agrimi U, Chiocchetti

- R (2007) Nitric oxide synthase immunoreactivity and NADPH-d histochemistry in the enteric nervous system of Sarda breed sheep with different PrP genotypes in whole-mount and cryostat preparations. *J Histochem Cytochem* 55:387–401
28. Mabbott NA, MacPherson GG (2006) Prions and their lethal journey to the brain. *Nat Rev Microbiol* 4:201–211
 29. Mbaku EM, Zhang L, Pearce WJ, Duckles SP, Buchholz J (2003) Chronic hypoxia alters the function of NOS nerves in cerebral arteries of near-term fetal and adult sheep. *J Appl Physiol* 94:724–732
 30. O'Rourke KI, Baszler TV, Miller JM, Spraker TR, Sadler-Rigglerman I, Knowles DP (1998) Monoclonal antibody F89/160.1.5 defines a conserved epitope on the ruminant prion protein. *J Clin Microbiol* 36:1750–1755
 31. O'Rourke KI, Baszler TV, Besser TE, Miller JM, Cutlip RC, Wells GA, Ryder SJ, Parish SM, Hamir AN, Cockett NE, Jenny A, Knowles DP (2000) Preclinical diagnosis of scrapie by immunohistochemistry of third eyelid lymphoid tissue. *J Clin Microbiol* 38:3254–3259
 32. Pfannkuche H, Schemann M, Gabel G (2002) Ruminal muscle of sheep is innervated by non-polarized pathways of cholinergic and nitrergic myenteric neurones. *Cell Tissue Res* 309:347–354
 33. Pompolo S, Furness JB (1998) Quantitative analysis of inputs to somatostatin-immunoreactive descending interneurons in the myenteric plexus of the guinea-pig small intestine. *Cell Tissue Res* 294:219–226
 34. Portbury AL, Pompolo S, Furness JB, Stebbing MJ, Kunze WA, Bornstein JC, Hughes S (1995) Cholinergic, somatostatin-immunoreactive interneurons in the guinea pig intestine: morphology, ultrastructure, connections and projections. *J Anat* 187:303–321
 35. Porto-Carreiro I, Fevrier B, Paquet S, Vilette D, Raposo G (2005) Prions and exosomes: from PrPc trafficking to PrPsc propagation. *Blood Cells Mol Dis* 35:143–148
 36. Press CM, Heggebo R, Espenes A (2004) Involvement of gut-associated lymphoid tissue of ruminants in the spread of transmissible spongiform encephalopathies. *Adv Drug Deliv Rev* 56:885–899
 37. Prinz M, Heikenwalder M, Junt T, Schwarz P, Glatzel M, Heppner FL, Fu YX, Lipp M, Aguzzi A (2003) Positioning of follicular dendritic cells within the spleen controls prion neuroinvasion. *Nature* 425:957–962
 38. Prusiner SB, Groth D, Serban A, Koehler R, Foster D, Torchia M, Burton D, Yang SL, DeArmond SJ (1993) Ablation of the prion protein (PrP) gene in mice prevents scrapie and facilitates production of anti-PrP antibodies. *Proc Natl Acad Sci USA* 90:10608–10612
 39. Prusiner SB (1998) Prions. *Proc Natl Acad Sci USA* 95:13363–13383
 40. Shmakov AN, McLennan NF, McBride P, Farquhar CF, Bode J, Rennison KA, Ghosh S (2000) Cellular prion protein is expressed in the human enteric nervous system. *Nat Med* 6:840–841
 41. Song ZM, Brookes SJ, Steele PA, Costa M (1992) Projections and pathways of submucous neurons to the mucosa of the guinea-pig small intestine. *Cell Tissue Res* 269:87–98
 42. Song ZM, Brookes SJ, Ramsay GA, Costa M (1997) Characterization of myenteric interneurons with somatostatin immunoreactivity in the guinea-pig small intestine. *Neuroscience* 80:907–923
 43. Spraker TR, O'Rourke KI, Balachandran A, Zink RR, Cummings BA, Miller MW, Powers BE (2002) Validation of monoclonal antibody F99/97.6.1 for immunohistochemical staining of brain and tonsil in mule deer (*Odocoileus hemionus*) with chronic wasting disease. *J Vet Diagn Invest* 14:3–7
 44. Timmermans JP, Hens J, Adriaensen D (2001) Outer submucous plexus: an intrinsic nerve network involved in both secretory and motility processes in the intestine of large mammals and humans. *Anat Rec* 262:71–78
 45. Tolcos M, Harding R, Loeliger M, Breen S, Cock M, Duncan J, Rees S (2003) The fetal brainstem is relatively spared from injury following intrauterine hypoxemia. *Brain Res Dev Brain Res* 143:73–81
 46. van Keulen LJ, Schreuder BE, Vromans ME, Langeveld JP, Smits MA (1999) Scrapie-associated prion protein in the gastrointestinal tract of sheep with natural scrapie. *J Comp Pathol* 121:55–63
 47. van Keulen LJ, Vromans ME, van Zijderveld FG (2002) Early and late pathogenesis of natural scrapie infection in sheep. *APMIS* 110:23–32
 48. Van Op den Bosch J, van Nassauw L, Lantermann K, van Marck E, Timmermans JP (2007) Effect of intestinal inflammation on the cell-specific expression of somatostatin receptor subtypes in the murine ileum. *Neurogastroenterol Motil* 19:596–606
 49. Vergara-Esteras P, Harrison FA, Brown D (1990) The localization of somatostatin-like immunoreactivity in the alimentary tract of the sheep with observations on the effect of an infection with the parasite *Haemonchus contortus*. *Exp Physiol* 75:779–789
 50. Vulchanova L, Casey MA, Crabb GW, Kennedy WR, Brown DR (2007) Anatomical evidence for enteric neuroimmune interactions in Peyer's patches. *J Neuroimmunol* 185:64–74
 51. Wathuta EM (1986) The distribution of vasoactive intestinal polypeptide-like, substance P-like and bombesin-like immunoreactivity in the digestive system of the sheep. *Q J Exp Physiol* 71:615–631