University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Transactions of the Nebraska Academy of Sciences and Affiliated Societies

Nebraska Academy of Sciences

1-1-1994

Genetic Variation and Differentiation of North American Waterfowl (Anatidae)

David W. Oates Nebraska Game and Parks Commission, doates@ngpc.state.ne.us

Joann D. Principato Nebraska Game and Parks Commission

Follow this and additional works at: http://digitalcommons.unl.edu/tnas



Part of the <u>Life Sciences Commons</u>

Oates, David W. and Principato, Joann D., "Genetic Variation and Differentiation of North American Waterfowl (Anatidae)" (1994). Transactions of the Nebraska Academy of Sciences and Affiliated Societies. Paper 108. http://digitalcommons.unl.edu/tnas/108

This Article is brought to you for free and open access by the Nebraska Academy of Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Transactions of the Nebraska Academy of Sciences and Affiliated Societies by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

GENETIC VARIATION AND DIFFERENTIATION OF NORTH AMERICAN WATERFOWL (ANATIDAE)

David W. Oates and Joann D. Principato

Nebraska Game and Parks Commission 2200 North 33rd Street Lincoln, Nebraska 68503

ABSTRACT

This study examines the genetic variation in 45 taxa of all tribes and most species of North American waterfowl (Anatidae) with a starch-gel electrophoretic survey of protein variation at 25 loci. Relationships were estimated using the resulting data from the patterns of allozyme variation and summarized in both phenetic and cladistic branching diagrams. The branching diagrams (phylogenetic trees) are employed to help compare and contrast phylogenetic relationships relative to other hypothesis. Although results of this study generally concur with classic phylogenetic trees and the taxonomic designations of the current American Ornithologists' Union (AOU) Check-list, exceptions are noted. Genetic data strongly contradict inclusion of Chen canagica (emperor goose) within the genus Chen. Clangula hyemalis (oldsquaw) and Melanitta nigra (black scoter) do not cluster on the branching diagrams as would be predicted from classical analysis. It is possible that they form a divergent group within the Tribe MERGINI.

† † †

The classification of waterfowl (Anatidae), while having been recently revised (American Ornithologists' Union [AOU] 1983, 1991), has remained controversial for the past several decades (Bellrose, 1980). Recognizing that avian taxa appeared to be oversplit relative to other taxa, Delacour and Mayr (1945) made a strong case for more inclusive genera and pooled many monotypic groups in the Anatidae. This taxonomic reasoning was followed in a number of subsequent studies (e.g. Brush, 1976; Delacour, 1954–1964; Johnsgard, 1961).

Early taxonomic studies of avian species typically involved analysis of morphology, plumage, and behavior. Delacour and Mayr (1945) employed behavior patterns, anatomy, and plumage in the study of Anatidae, while Johnsgard (1961) relied primarily on behavior and Livezey (1986) on morphology. Molecular analyses

were recognized as being of value to avian taxonomy, beginning with Sibley's (1960) electrophoretic evaluation of avian egg-white proteins.

While Sibley's avian studies (e.g. Sibley, 1968, 1970; Sibley and Ahlquist, 1972; Sibley et al., 1969) were focused at the generic and familial levels, later studies largely involved the attempt to identify avian species and their relationships. These studies involved the electrophoretic analysis of egg white, blood, and feather proteins (Brown et al., 1970; Brush, 1976; Ford et al., 1974; Morgan et al., 1976; Shaughnessy, 1970). They succeeded to varying degrees, but were limited primarily by the uncertain homologies and genetic basis of the few protein phenotypes detected (Brush, 1979).

Current electrophoretic avian studies use more refined histochemical staining techniques. Identification of up to ten alleles each from 20-30 biochemically detectable loci (Aquadro and Avise, 1982; Avise et al., 1980a; Johnston, 1983; Mindell and Sites, 1987; Seeb et al., 1986; Yang and Patton, 1981) is typical, making electrophoresis a valuable tool for measuring genetic distances and estimating phylogenetic relationships (Ankney et al., 1986; Baker, 1990; Barrett and Vyse, 1982; Barrowclough, 1983; Barrowclough et al., 1981; Browne et al., 1993; Gutierrez et al., 1983; N. Johnson et al., 1988; Morgan et al., 1976; Patton and Avise, 1986; Sherman, 1981; Smith and Zimmerman, 1976; Zink, 1982). Protein-electrophoretic analyses have been developed for the enforcement of wildlife laws and are employed when identification of a species is otherwise impossible (e.g. Harvey, 1990; Oates et al., 1983; Seeb et al., 1990; Utter, 1991).

Electrophoretic analysis of protein variation was employed in this study to examine phylogenetic relationships among individuals representing all of the tribes and most of the species of waterfowl found in

Table 1. Scientific classification of the species of the family Anatidae (swans, geese and ducks) analyzed in this study [after AOU (1983) and Bellrose (1980)]. Common names follow Bellrose (1980).

Subfamily	Tribe	Sci	entific name	Common name
Anserinae	Dendrocygnini	1.	Dendrocygna bicolor helva	Fulvous whistling-duck
	Cygnini	2.	Cygnus columbianus	Tundra swan
	Anserini	3.	Anser albifrons frontalis	Greater white-fronted goose
		4.	Anser albifrons gambelii	Tule goose
		5.	Chen caerulescens caerulescens	Snow goose
		6.	Chen caerulescens caerulescens	Blue goose
		7.	Chen caerulescens atlantica	Greater snow goose
		8.	Chen rossii	Ross' goose
		9.	Chen canagica	Emperor goose A
			Chen canagica	Emperor goose B
		11.	Branta bernicla nigricans	Black brant
		12.	$Branta\ bernicla\ hrota$	American brant
		13.	Branta canadensis canadensis	Canada goose
		14.	Branta canadensis minima	Cackling Canada goose
Anatinae	Cairinini	15.	Aix sponsa	Wood duck
	Anatini	16.	Anas crecca	Green-winged teal
		17.	Anas rubripes	American black duck
		18.	Anas fulvigula	Mottled duck
			Anas platyrhynchos	Mallard
			Anas acuta	Northern pintail
		21.	Anas discors	Blue-winged teal
		22.	Anas cyanoptera	Cinnamon teal
		23.	Anas clypeata	Northern shoveler
		24.	Anas strepera	Gadwall
		25 .	Anas americana	American widgeon
	${f A}$ YTHYINI	26.	Aythya valisneria	Canvasback
		27.	Aythya americana	Redhead
		28.	$Aythya\ collar is$	Ring-necked duck
		29.	Aythya marila	Greater scaup
		30.	Aythya affinis	Lesser scaup
	${f M}$ ergini	31.	Somateria mollissima	Common eider
		32.	Somateria spectabilis	King eider
		33.	Polysticta stelleri	Steller's eider
		34.	Histrionicus histrionicus	Harlequin duck
		35.	Clangula hyemalis	Oldsquaw
		36.	Melanitta nigra	Black scoter
		37.	Melanitta perspicillata	Surf scoter
		38.	Melanitta fusca	White-winged scoter
			Bucephala clangula	Common goldeneye
			Bucephala islandica	Barrow's goldeneye
		41.	-	Bufflehead
			Lophodytes cucullatus	Hooded merganser
			Mergus merganser	Common merganser
		44.	-	Red-breasted merganser
	Oxyurini		Oxyura jamaicensis	Ruddy duck

North America. The phylogenetic relationships were examined using both phenetic and cladistic approaches, with branching diagrams constructed for each to summarize the electrophoretic data.

MATERIALS AND METHODS

A total of 429 individuals, representing 40 species and 5 subspecies, was collected (Table 1) from the U.S. Fish and Wildlife Service in the states of Alaska, California, Colorado, Nebraska, and Washington. As soon as possible after collection, samples were frozen, shipped on dry ice to the laboratory, and subsequently stored at -20° C. In most cases a complete wing was obtained for the study, while in other cases a sample consisted of a one-gram piece of muscle. In preparation for electro-

phoresis, small pieces of muscle were extracted and placed in 12×72 mm culture tubes to which equal volumes of water were added.

Electrophoresis followed procedures outlined in May et al. (1979) and Utter et al. (1974). Three buffer systems were employed: 1) MF-tris-boric acid EDTA gel with tray buffer (pH 8.5) (Boyer et al. 1963); 2) RW-tris-citric acid gel buffer (pH 8.5) with lithium hydroxide-boric acid tray buffer (pH 8.5) (Ridgway et al. 1970); and 3) AC+-modification of the amine-citrate buffer of Clayton and Tretiak (1972). In the AC+ buffer system, the tray remained unchanged (pH 6.1), while the gel buffer was raised to pH 6.4 with N-(3-aminopropyl) morpholine. Staining for enzyme activity followed methods outlined in Allendorf et al. (1977), Harris and

Table 2. Proteins: their locus abbreviations, buffer systems and Enzyme Commission numbers. Buffer systems are abbreviated as described in the text. Loci found to be monomorphic throughout a family are marked with an asterisk.

Protein	Locus Abbreviation	Buffer System	Enzyme Commission Number
Adenosine deaminase	ADA	MF	3.5.4.4
Adenylate kinase	*AK	AC+	2.7.4.3
Albumen	ALB	RW	_
Aspartate aminotransferase	$^{AAT ext{-}1}_{*AAT ext{-}2}$	AC+	2.6.1.1
Creatine kinase	* <i>CK</i>	RW	2.7.3.2
Glucose phosphate isomerase	GPI	RW	5.3.1.9
Glyceraldehyde-3-phosphate dehydrogenase	*gЗpdh	AC+	1.2.1.12
Guanine deaminase	GDA	AC+	3.5.4.3
Hemoglobin	HEM	AC+	_
Iscocitrate dehydrogenase	IDH	AC+	1.1.1.42
Lactate dehydrogenase	*LDH-1 LDH-2	AC+	1.1.1.27
Malate dehydrogenase	мдн-1 мдн-2	AC+	1.1.1.37
Nucleoside phosphorylase	NP	AC+	2.4.2.1
Peptidase	*PEP-1 PEP-2 PEP-3 PEP-4	AC+	3.4.1.1
Phosophoglucomutase	PGM	RW	5.4.2.2
Phosphomannose isomerase	<i>РМІ-1</i> <i>РМІ-2</i>	AC+	5.3.1.8
Superoxide dismutase	SOD-1 SOD-2	\mathbf{MF}	1.15.1.1

Hopkinson (1976), Selander et al. (1971), and Shaw and Prasad (1970). A list of the 25 enzymes resolved is given in Table 2.

Locus designation followed the system of nomenclature suggested by Allendorf and Utter (1976) and Shaklee (1990). For each locus, the mobility of the most common allele in mallards (*Anas platyrhynchos*) was used as a standard and designated as 100 with the mobility of all other alleles calculated relative to this allele. A mallard sample was included on every gel for reference. It was often necessary to compare each allele to several others, in addition to that of the standard mallard, when numerous alleles occurred at some loci. After electrophoresis, each sample was scored for its observed genotype. Allelic frequencies at each locus were calculated for each species.

The use of electrophoretic data to estimate phylogenetic trees has been addressed by Avise (1994), Farris (1981), Felsenstein (1981), Hartl and Clark (1989), Mickevich and Mitter (1981), and Straney (1981). Branching diagrams (phylogenetic trees) for both phenetic and cladistic approaches were constructed from a matrix of genetic distances. Distance analyses, based on the estimation of pair-wise genetic distances between taxa (Avise, 1994), avoid both the necessity of a transition series of alleles and the need for assignment of ancestral or derived states.

In this study, the phenetic approach utilizes the mutation-drift model of Nei (1972). It is assumed that time and genetic divergence are correlated, with many genes evolving at a constant rate (Forey et al., 1992; Mayr, 1991; Wilson, 1976) and genetic distances increasing with time (Avise, 1994). Unbiased distance values (D) were estimated between every pair of species from the allelic frequencies, using Nei's (1978) formula. The phenogram (Fig. 1) was produced from a matrix of D-values using the unweighted pair-group method with arithmetic averages (UPGMA) (Sneath and Sokal, 1973). Extant clusters were "right-justified" along the genetic distance axis (Avise, 1994).

A cladistic pattern among extant taxa and hypothetical ancestors, based on the most parsimonious solution instead of rates of evolution, is presented using a Wagner distance analysis (Farris, 1972). Modified Rogers' distance values (D_T) (Wright, 1978) were calculated from the allelic frequencies and subjected to the distance Wagner procedure of Swofford (1981) to generate an unrooted network. The network was rooted at the midpoint of the longest pathway between extant taxa (Avise, 1994; Farris, 1972), creating a rooted distance Wagner diagram (Fig. 2).

Both the phenogram (Fig. 1) and distance Wagner

diagram (Fig. 2) are compared to a classical phylogenetic estimate (Fig. 3) derived from Bellrose (1980) and based on Johnsgard (1978). The generic and species designations are those of the AOU *Check-list of North American Birds* (1957, 1983) and AOU supplements to the check-list (1985, 1987, 1989, 1991).

BIOSYS-1 (Swofford and Selander 1981) was used to analyze the electrophoretic data and estimate heterozygosities (H).

ELECTROPHORESIS

Gene products from approximately 60 presumptive loci were initially examined, with 25 loci resolved adequately enough for routine scoring. Since no genetic variation was found at the AK, AAT-2, CK, G3PDH, LDH-1, and PEP-1 loci, they were considered to be monomorphic throughout the family. The mean heterozygosity (weighted by sample size) was found to be consistent with published avian values (Aquadro and Avise, 1982; Avise, 1994; Patton and Avise, 1986).

Resulting electrophoretic patterns of most loci agreed with previously published results (e.g. Ankney et al., 1986; Avise et al., 1980a; Browne, 1993; Patton and Avise, 1986). Loci of special interest or those which had results previously unpublished are reported in this section. Allelic frequencies of all loci considered to be polymorphic can be found in Table 3.

Adenosine deaminase

One ADA locus was resolved in the family Anatidae. A high degree of polymorphism was exhibited both within and between a majority of the species. Heterozygotes exhibited a two-banded pattern at the ADA locus, consistent with a monomeric structure.

Peptidase

Peptidase zones of activity (PEP-1, PEP-2, PEP-3) appeared in all species of the family Anatidae. No genetic variation was found at the PEP-1 locus (resolved with a DL-leucylglycylglycine substrate). PEP-2,3 were resolved with DL-leucyl-DL-alanine. An additional zone of activity, PEP-4 (also resolved with leucylalanine), appeared only in Chen canagica (emperor goose) and was scored on a presence or absence basis.

Mannose-6-phosphate isomerase

MPI-1, a highly polymorphic monomer, was clearly expressed in all species. MPI-2, another zone of activity, generally migrated at a uniform distance from MPI-1. Dendrocygna b. helva (fulvous whistling-duck) expressed MPI-2(100) locus mobility (identical to most of the others in the subfamily Anserinae) and was fixed for a faster MPI-1(152) allele. MPI-2 was not expressed in all species examined, analogous to that of CK-1,2 in North Ameri-




Figure 1. Phenogram constructed using the unweighted pair-group method with arithmetic averages (UPGMA) from a matrix of D-values. Phenetic relationships of the family Anatidae based upon Nel's (D) (Nel, 1978).




Figure 2. Distance Wagner Diagram for the family Anatidae constructed using modified Rogers' distance measure (D_{τ}) and distance Wagner procedure of Swofford (1981).

can thrushes (Avise et al., 1980a). Although MPI-1,2 can be interpreted as being the products of a single structural locus, consideration was also given to the identification by Finnerty and Johnson (1979) of the heritable, non-allelic variation found for xanthine dehydrogenase and aldehyde oxidase in Drosophila melanogaster (see also Finnerty et al., 1979, Johnson et al., 1981). It was concluded in this study, as in Avise et al. (1980a), that MPI-1,2 were present as two sets of taxonomically useful characters. It could not be unequivocally determined, however, whether or not the two zones of activity ultimately represented the products of a single structural locus.

PATTERNS OF GENETIC DIFFERENTIATION

The subfamily Anserinae

The subfamily Anserinae includes the whistling-ducks, swans, and geese: the tribes Dendrocygnini, Cygnini and Anserini, respectively. Those in the subfamily were not diagnostic for all species from the other Anatidae at the GDA, NP, PEP-2, and MPI-1,2 loci. The ADA, GDA, GPI, NP, PEP-2,3, PGM, and MPI-1,2 loci were useful for identification within the subfamily.

DENDROCYGNINI (Whistling-ducks): Dendrocygna. Dendrocygna b. helva (fulvous whistling-duck) was the only species available from this predominantly South American tribe. Dendrocygna was distinguished from the other subfamily members by its electrophoretic migration patterns of alleles at the PEP-2 loci and by large frequency differences at the PGM locus. Based on the phenetic and cladistic analyses, Dendrocygna represents the most divergent genus of the Anserinae examined. It joins the rest of the subfamily at a D of 0.33 in the phenogram (Fig. 1). Since Oxyura and Dendrocygna are sister taxa on the distance Wagner diagram (Fig. 2), and Oxyura is observed as being the most divergent member of the subfamily Anatinae in both the phenogram (Fig. 1) and classical taxonomy (Fig. 3), the sister group relation of the *Dendrocygna* to the remainder of the subfamily Anserinae could not be resolved in the analyses. Sibley and Monroe (1990) classify this tribe as the family Dendrocygnidae, based on the DNA-DNA hybridization studies of Sibley and Ahlquist (1983, 1987, 1990).

Cygnini (Swans): Cygnus. Only specimens from one member of this tribe, Cygnus columbianus (tundra swan, formerly whistling swan) were available for analysis. While Cygnus columbianus could be distinguished from the genus Anser and some members of the genus Chen at GPI(100), it differed from all of the other genera of the subfamily Anserinae at the ADA, NP, and PMI-1,2 loci. These findings seem to coincide with the classification of the tribe as the subfamily Cygninae by Sibley and Monroe (1990) who base their classification on the




Figure 3. Phylogeny (classification) of the family Anatidae based upon structure and behavior [after Bellrose (1980) based on Johnsgard (1961, 1978)].

DNA-DNA hybridization studies of Sibley and Ahlquist (1983, 1987, 1990). Although a phenogram and a rooted Wagner network are not necessarily expected to form concordant branching, the diagrams do exhibit identical branching in the case of *Cygnus*. In the phenetic analysis, *Cygnus* has a D of 0.29 from the members of the genera *Anser*, *Branta*, and *Chen*.

Anserini (Geese): Anser, Branta, and Chen. The electrophoretic data separated the Anserini (geese) into three heterogenous aggregations: the species Chen canagica (emperor goose), the genus Branta, and a group made up of members of Anser and the remaining Chen. Both the phenogram (Fig. 1) and the distance Wagner diagram (Fig. 2) imply that Cygnus may be a sister taxon to these geese.

Chen canagica (emperor goose) was the most divergent taxon examined, exhibiting an additional peptidase locus (PEP-4), and differing from other members of the subfamily Anserinae by fixation for the ADA(110) allele. At the MPI-1 locus, two of the five specimens, designated "Emperor A," were fixed for the MPI-1(110) allele, while the other three specimens, designated "Emperor B," were fixed for the MPI-1(93) allele. The probability is less than 0.05 that these samples represent one population which is polymorphic at the MPI-1 locus.

Considering the Anser/Chen complex, Chen c. caerulescens (snow goose and blue goose), Chen c.

Table 3. Allele frequency estimates for the family Anatidae. An explanation of the numbers is given on page 130. Full names for species are given in Table 1. Sample number (N) is in parentheses following the name unless it differs at a particular locus, in which case it is given as a superscript.

TAXONOMY ADA **ALB													$**_A$	LB		
Locus:	100	107	95	110	113	98	103	79	120	105	100	98		101	97	104
Demonarchini																
DENDROCYGNINI 1. Dendrocygna b. h. (6)				0.25						0.75			1.00			
Cygnini ——————————————————————————————————				0.25						0.75			1.00			
2. Cygnus c. (2)									1.00				1.00			
Anserini —————					_				1.00				1.00			
3. Anser a. f. (8)	0.38									0.62			1.00			
4. Anser a. g. (20)	1.00									0.02			1.00			
5. Chen cae. c. (20)	0.10	0.28	0.05	0.57									1.00			
6. Chen cae. c. (10)	0.20	0.50	0.05	0.30		0.15							1.00			
7. Chen cae. a. (10)	0.15	0.35	0.05	0.45		0.20							1.00			
8. Chen r. (10)	0.15	0.35	0.00	0.45		0.05							1.00			
9. Chen can. (2)	0.20	0.00		1.00		0.00							1.00			
10. Chen can. (3)				1.00									1.00			
11. Branta b. n. (11)	0.04	0.64	0.23	0.09									1.00			
12. Branta b. h. (2)		0.75	0.25										1.00			
13. Branta c. c. (10)	0.10	0.75	0.15										1.00			
14. Branta c. m. (17)	0.03	0.94	0.03										1.00			
Cairinini —																
15. Aix s. (10)			0.10				0.90							1.00		
Anatini ———————																
16. Anas c. (10)	0.50	0.22	0.07		0.14		0.07				1.007	7				
17. Anas r. (10)	0.75^{7}	0.15^{7}					0.05^{7}	,			1.00					
18. Anas f. (10)	0.55	0.45									1.00					
19. Anas p. (31)	0.82	0.18									1.00^{2}	29				
20. Anas a. (10)	0.30	0.70										1.00				
21. Anas d. (10)		0.11	0.28	0.11			0.28	0.22				0.05		0.95		
22. Anas c. (10)	0.05		0.70				0.15							1.00		
23. Anas c. (10)			0.56				0.22	0.22				1.00				
24. Anas s. (10)			0.06				0.94								1.00	
25. Anas a. (10)			0.15				0.80	0.05					1.00			
Aythyini ——————																
26. Aythya v. (10)		0.95					0.05						1.00			
27. Aythya a. (20)		0.80					0.20						1.00			
28. Aythya c. (10)			0.35	0.10			0.55						1.00			
29. Aythya m. (5)		1.00											1.00			
30. Aythya a. (10)		0.95				0.05							1.00			
Mergini —————																
31. Somateria m. (10)					1.00								1.00			
32. Somateria s. (3)					1.00								1.00			
33. Polysticta s. (7)							1.00						1.00^{1}			
34. Histrionicus h . (2)					1.00								1.00			
35. Clangula h. (10)						0.44	0.56						1.00			
36. <i>Melanitta n.</i> (10)		0.06			0.94								1.00			
37. <i>Melanitta p.</i> (10)					1.00								1.00			
38. Melanitta f. (10)					0.25				0.75				1.00			
39. Bucephala c. (10)					1.00								1.00			
40. Bucephala i. (10)					1.00						0.05		0.95			
41. Bucephala a. (10)					1.00								1.00			
42. <i>Lophodytes c.</i> (10)		1.00														1.00
43. Mergus m. (10)		-			1.00											1.00
44. Mergus s. (10)					0.95		0.05									1.00
Oxyurini														_		
45. Oxyura j. (10)								1.00					1.00			

 $^{{\}bf **Loci} \ at \ which \ electrophoretic \ patterns \ agree \ with \ previously \ published \ results.$

Table 3. Continued.

	*	*AAT-					**GPI					**GDA					
	100		-700	100	160	-66		-150	385	180	100	94	108	89	81		
					_							_					
1.	1.00			1.00							1.00						
$-{2}$.	1.00			1.00								1.00					
				1.00								1.00					
3.	1.00					0.88				0.12				0.88	0.12		
4.	1.00					1.00								1.00			
5.	1.00			0.02		0.98								1.00			
6.	1.00					0.95	0.05							1.00			
7.	1.00					0.90	0.10							1.00			
8.	1.00 1.00			1.00		1.00								1.00			
9.	1.00			1.00										1.00 1.00			
10. 11.	0.95		0.05	1.00 1.00										1.00			
11. 12.	1.00		0.05	1.00										0.50	0.50		
12. 13.	1.00			1.00										0.95	0.05		
14.	1.00			1.00										1.00	0.00		
														1.00			
15.	1.00			1.00							1.00						
16.	1.00			1.00							0.50		0.50				
17.	1.00			1.00							1.00						
18.	1.00			1.00							1.00						
19.	0.98^{29}	0.02^{29}	9	0.95	0.02				0.03		1.00						
20.	1.00			1.00							1.00						
21.	1.00			1.00							0.25	0.75					
22.	1.00			1.00								1.00					
23.	1.00			1.00								1.00					
24.	1.00			1.00								1.00					
25.	1.00			1.00							1.00						
26.	1.00			1.00							1.00						
27.	1.00			1.00							1.00						
28.	1.00			1.00							1.00						
29.	1.00			0.90		0.10					1.00						
30.	1.00			1.00							0.75	0.05	0.20				
31.	1.00			1.00							1.00						
32.	1.00			1.00							1.00						
33.	1.00			1.00							No Da	ta					
34.	1.00			1.00							1.00						
35.	1.00			1.00								1.00					
36.	1.00			1.00		0.05	0.05				1.00						
37.	1.00					0.95	0.05				1.00						
38.	1.00 1.00			1.00		1.00					1.00						
39. 40.	1.00			1.00 1.00							1.00 1.00						
40. 41.	1.00			1.00							1.00						
41. 42.	1.00			1.00							1.00						
42. 43.	1.00			1.00							1.00						
44.	1.00			1.00							1.00						
<u>45.</u>		1.00		0.05			0.05	0.00			1.00						
4 0.		1.00		บ.บอ			บ.บอ	0.90			1.00						

 $[\]ensuremath{^{**}\text{Loci}}$ at which electrophoretic patterns agree with previously published results.

136 D. W. Oates and J. D. Principato

Table 3. Continued.

		**HE	M			**	IDH				4	** <i>MD</i>	** _М DH-2					
	-100	-109 -	118	-94	-100		-95	-81	100	155	242	31	-100 -	162	-31	-25	100 -	136
1.	No Dat	ta			1.00				1.00				1.00				1.00	
2.	1.00				1.00				1.00				1.00				1.00	
3.	1.00				1.00				1.00				1.00				1.00	
4.	1.00				1.00				1.00				1.00				1.00	
5.	0.80			0.20	1.00				1.00				1.00		=		1.00	
6.	0.90			0.10	1.00				1.00				0.95		0.05		1.00	
7. 8.	0.80 1.00			0.20	$1.00 \\ 1.00$				$1.00 \\ 1.00$				1.00 1.00				1.00 1.00	
8. 9.	1.00				1.00				1.00 1.00								1.00	
9. 10.	1.00				1.00				1.00 1.00				1.00 0.67			0.33	1.00	
11.	1.00				1.00				1.00				1.00			0.55	1.00	
11. 12.	1.00				1.00				1.00				1.00				1.00	
13.	1.00^{21}	l			0.90^{2}	1		0.10^{21}	1.00^{21}	l			1.00				1.00	
14.	1.00				0.97			0.10	0.97			0.03	1.00				1.00	
								0.00				0.00						
15. ——	1.00				1.00				1.00				1.00				1.00	
16.	1.00				1.00				1.00				1.00				1.00	
17.	1.00				1.00				1.00				1.00				1.00	
18.	1.00				1.00				1.00				1.00				1.00	
19.	0.97	0.02	0.01		1.00				1.00				1.00				1.00	
20.	1.00				1.00				1.00				1.00				1.00	
21.	1.00				1.00				1.00				1.00				0.95	0.05
22.		0.15		0.85	1.00				1.00				1.00				1.00	
23.				1.00	1.00				1.00				1.00				1.00	
24.	1.00				0.95	0.95	5		1.00				1.00				1.00	
25.	1.00				1.00				1.00				1.00				1.00	
26.	1.00				1.00				0.78	0.22			1.00				1.00	
27.	1.00				1.00				1.00				1.00				1.00	
28.	1.00				1.00				1.00				1.00				1.00	
29.	1.00				1.00				1.00				1.00				1.00	
30.	1.00				1.00				1.00				1.00				1.00	
31.			1.00		1.00				1.00				1.00				1.00	
32.	0.17		0.83		1.00				1.00				1.00				1.00	
33.	1.00				1.00				1.00				1.00				1.00	
34.		1.00			1.00				1.00				1.00				1.00	
35.		1.00			1.00				1.00		0.05		1.00				1.00	
36.	1.00	1.00			1.00				0.95		0.05		1.00				1.00	
37.	1.00		1.00		1.00				1.00				1.00				1.00	
38.	1.00		1.00		$1.00 \\ 1.00$				1.00 1.00				1.00 1.00				1.00 1.00	
39. 40.	1.00				1.00				1.00				0.85	0.15			1.00	
40. 41.	1.00				1.00				1.00				1.00	0.10			1.00	
41. 42.	1.00				1.00				1.00				1.00				1.00	
42. 43.	1.00				1.00				1.00				1.00				1.00	
44.	1.00				1.00		1.00		1.00				1.00				1.00	
 45.	1.00				1.00				1.00				1.00				1.00	

 $[\]ensuremath{^{**}\text{Loci}}$ at which electrophoretic patterns agree with previously published results.

						^k NP								PEP-2			
	100	87	113	<u>79</u>	93	73	40	107	123	55	100	82	107	94_	114	75	88
1.		0.08		0.08		0.84											
2.			1.00									1.00					
3.							******	1.00				1.00					
4.								1.00				1.00					
5.								1.00				1.00					
6.								1.00				1.00					
7.								1.00				1.00					
8.								1.00				1.00					
9.								1.00				1.00					
0. 1.								$1.00 \\ 1.00$			0.09	$1.00 \\ 0.09$		0.82			
2.								1.00			0.09	0.09		1.00			
2. 3.								1.00^{2}	1			0.79	0.02	0.17		0.02	
4.								1.00			0.06	$0.73 \\ 0.24$	0.02	0.70		0.02	
5.		0.90				0.10					0.94		0.06				
6.		0.94^{9}				0.06^{9}					0.21		0.07		0.72		
7.	0.75	0.20		0.05							0.95	0.05					
8.	0.70	0.25	0.05								0.60	0.25	0.10			0.05	
9.	0.77	0.19	0.02			0.02					0.92	0.02					
0.	0.55	0.35	0.05	0.05							0.67		0.22	0.06	0.05		
1.		0.06^{8}									0.05		0.10	0.80	0.05		
2.	0.72		0.28											1.00			
3.	1.00^{6}													1.00			
4.	0.05			0.95							0.07		0.93				
5.	0.10		0.05	0.75	0.05	0.05							1.00				
6.				1.00 0.89		0.11					0.15		1.00				
7. 8.				0.89 0.93^7		0.11 0.07^7					$0.15 \\ 1.00$		0.85				
9.				0.90	0.10	0.07					1.00						
0.			0.05	0.90	0.10	0.05					1.00						
1.	1.00											0.86					
2.	1.00										0.11	1.00					
3.	0.08	0.75			0.17						No Da						
4.							1.00				1.00						
5.	0.05				0.90			0.05					0.14	0.86			
6.			0.95					0.05			0.38		0.06	0.56			
7.					1.00												
8.					1.00									1.00			
9.			0.95						0.05		0.94						0.0
0.			1.00								0.71						0.2
1.									1.00		0.86		0.07		0.07		
2.				0.50	0.50					1.00	0.43		0.57				
3. 4.				0.64	0.36					1.00	$1.00 \\ 0.92$	0.08					
													0.17				
15.						1.00					0.83		0.17				

^{**}Loci at which electrophoretic patterns agree with previously published results.

138 D. W. Oates and J. D. Principato

Table 3. Continued.

	PEP-2	(cont.)					PEP-4	**PGM							
	79	66	100	86	95	108	89	80	104	75	92	91	No Loc. 100	100	140
1.		1.00		0.75	0.25								1.00	1.00	
2.					1.00								1.00		
3.				1.00									1.00		
4.				0.75				0.25					1.00		
5.							0.94	0.06					1.00		0.03
6.							1.00						1.00		
7.							0.83	0.17					1.00	0.25	
8.							0.60				0.40		1.00		
9.							1.00						1.00		
LO.							1.00						1.00	0.17	
l1.					0.05		0.18				0.77		1.00		
l2.							0.75				0.25		1.00		
l3.			0.35		0.60		0.05						1.00	0.05	
L4.				0.03	0.85		0.06				0.06		1.00	0.03	
L5.			0.06	0.22	0.72								1.00		
l6.									1.00				1.00	0.79	0.21^{7}
L7.			0.95				0.05		1.00				1.00	1.00	0.21
l8.			0.75		0.15		0.00		0.10				1.00	1.00	
L9.	0.06		0.95		0.02		0.02		0.01				1.00	1.00	
20.	0,00		0.85		0.05		0.02		0.10				1.00	0.95	0.05
21.			0.17	0.33	0.33			0.11					1.00		0.06^{8}
22.				0.29	0.57		0.14						1.00	0.90	
23.				0.50	0.50								1.00	1.00	
24.			0.85	0.15									1.00	1.00	
25.			1.00										1.00	1.00	
26.			0.25		0.70							0.05	1.00		
27.			0.88		0.12							0.00	1.00	0.13	
28.			1.00		0.1_								1.00	0.30	
29.			0.90		0.10								1.00	0.50	
30.			0.72	0.06	0.22								1.00	0.20	
31.			1.00		_								1.00		
32.			1.00										1.00		
33.			1.00										1.00	0.50	
34.					1.00								1.00	1.00	
35.			0.05		0.05	0.90							1.00		
36.								0.05					1.00	0.05	
37.	1.00						0.78		0.22				1.00		
38.			1.00										1.00		
39.						0.05			0.95				1.00	0.90	
1 0.									1.00				1.00	1.00	5
41.					0.10				0.90				1.00		
12 .					1.00								1.00	1.00	
1 3.				0.05									1.00		
14.				0.15	0.60			0.15		0.10			1.00		
1 5.			0.20						0.80				1.00		1.00^{8}

 $[\]ensuremath{^{**}\text{Loci}}$ at which electrophoretic patterns agree with previously published results.

Table 3. Continued.

**	PGM (cont.)					M	PI-1					MPI	-2	**S	DD-1	**S01)-2
	55	72	100	120	133	55	67	110	75	93	152	142	No Loc	. 100_	-100	-33	100	250
1.											1.00			1.00	1.00			1.00
2.	1.00		1.00	_									1.00		1.00			1.00
3.	1.00									0.88		0.12		1.00	1.00		0.12	0.88
4.	1.00									1.00				1.00	1.00			1.00
5.	0.97									0.95	0.05			1.00	1.00		0.03	0.97
6.	1.00									0.95	0.05			1.00	1.00			1.00
7. 8.	$0.75 \\ 1.00$									1.00				1.00	1.00 1.00			1.00 1.00
9.	1.00							1.00		1.00				1.00 1.00	1.00			1.00
10.	0.83							1.00		1.00				1.00	1.00			1.00
10. 11.	1.00							0.09		0.91				1.00	1.00			1.00
12.	1.00							0.00		1.00				1.00	1.00			1.00
13.	0.95							0.14^{23}	l	0.86^{2}	1			1.00^{21}				1.00
14.	0.97									1.00				1.00	1.00			1.00
15.	0.95	0.05	1.00										1.00		1.00	_	1.00	
16.			0.11				0.72			0.17	_		1.00		1.00		1.00	
17 .			0.65	0.35									1.00		0.95	0.05	1.00	
1 8.			0.80	0.10	0.05			0.05					1.00		1.00		1.00	
19.			0.84	0.11	0.03							0.02	1.00		1.00		1.00	
20.			0.85				0.15						1.00		1.00		1.00	
21.			0.40			0.15	0.45						1.00		1.00		1.00	
22. 23.			0.30			0.35	$0.70 \\ 0.35$		0.05				1.00 1.00		1.00		1.00 1.00	
23. 24.			$0.25 \\ 0.55$	0.10		0.33	0.35		0.05				1.00		1.00		1.00	
2 4 . 25.			0.55	0.10			0.95			0.05			1.00		1.00		1.00	
							0.00											
26.	1.00		0.95	0.05									1.00		1.00		1.00	
27.	0.87		0.75	0.10	0.05		0.15						1.00		1.00		1.00	
28. 29.	$0.70 \\ 0.50$		$0.95 \\ 0.90$		0.05							0.10	1.00 1.00		1.00 1.00		1.00 1.00	
29. 30.	0.80		1.00									0.10	1.00		1.00		1.00	
31.	1.00								1.00				1.00		1.00			1.00
$\frac{31.}{32.}$	1.00								1.00				1.00		1.00			1.00
33.	0.50		0.90				0.10		1.00				1.00		1.00		No D	ata
34.							1.00						1.00		1.00			1.00
35.	1.00			0.10		0.80	0.10						1.00		1.00		1.00	
36.	0.95								0.95	0.05			1.00		1.00		1.00	
37.	1.00										0.33		1.00		1.00			1.00
38.	1.00		0.00						0.50	0.00		1.00	1.00		1.00			1.00
39.	0.10		0.20						0.50	0.30			1.00		1.00			1.00
40.	1.00		0.06					0.06	1.00 0.83	0.05			1.00 1.00		1.00 1.00			1.00 1.00
41. 42.	1.00		0.06					0.06	1.00	0.05			1.00		1.00			1.00
42. 43.	1.00		0.20						0.80				1.00		1.00			1.00
44.	1.00		0.20	0.05					0.00	0.95			1.00		1.00			1.00
4 5.									_	0.10	0.05	0.85	1.00		1.00			1.00

^{**}Loci at which electrophoretic patterns agree with previously published results.

atlantica (greater snow goose), and Chen rossi (Ross' goose)—members of the Ross'/snow goose cluster—were found to be identical, differing from the greater white-fronted/tule goose representatives of Anser by the large frequency differences at the ADA and PEP-3 loci. Anser a. frontalis (greater white-fronted goose) and Anser a. gambelli (tule goose) appeared to have distinguishing loci, most notably ADA. While the tule geese were fixed for the ADA(100) allele, all of the greater white-fronted geese possessed the ADA(100) allele at a frequency of only 0.38.

Branta, most similar to the Anser/Chen complex, excluding C. canagica, differed from the group at the ADA, PGI, and PEP-2 loci. Genetic differences were exhibited within Branta at the PGM and PEP-2,3 loci. Although no fixed differences occurred, allelic frequency differences found at these loci may allow discrimination between Branta b. nigricans (black-brant) and Branta b. hrota (American brant) at the PEP-3 locus, and between Branta c. canadensis (Canada goose) and Branta c. minima (cackling Canada goose) at the PEP-2 locus.

The subfamily Anatinae

Anatinae, the largest and most complex subfamily, contains the "typical" ducks. With some exceptions, the relationships based on the phenogram parallel the generic relationships, and the major clusterings on the Wagner diagram reflect the divisions into tribes. The three tribes (Anatini, Aythyini, Cairinini) are clustered with the Anserinae in the distance Wagner diagram (Fig. 2), in contrast with their placement on the phenogram (Fig. 1) and in classical taxonomy (Fig. 3). However, the common root is extremely short, 0.049, and its placement is subject to interpretation.

CAIRININI (Muscovy ducks and allies): Aix. The genus Aix is represented by one species in North America, Aix sponsa (wood duck), which is the only North American member of the tribe. It differed at the ADA, ALB, NP, PEP-2,3, and PGM loci from the majority of both Aythya and Anas. Concurring with existing literature, Aix is closest to Anas on the distance Wagner diagram (Fig. 2), but is closest to Aythya on the phenogram (Fig. 1).

Anatini (Dabbling ducks): Anas. The genus Anas, a large polytypic aggregation, was examined. Large frequency differences were found at the ADA, ALB, NP, and PEP-2,3 loci for the ten species analyzed. Anas platyrhynchos (mallard), Anas fulvigula (mottled duck), and Anas rubripes (American black duck) form a genetically close knit group. Clangula hyemalis (oldsquaw), one of the most dissimilar Mergini, is polyphyletic with Anas in the phenogram (Fig. 1), but does not cluster with the genus in the distance Wagner diagram (Fig. 2) nor in classical taxonomy (Fig. 3).

AYTHYINI (Pochards and allies): Aythya. Aythya affinis (lesser scaup) and Aythya marila (greater scaup) were electrophoretically identical to a great degree. With small N sampled, Aythya americana (redhead) and Aythya valisineria (canvasback) differed only for allelic frequencies at the PEP-3 locus. Aythya collaris (ring-necked duck) differed from the rest of Aythya the ADA and PEP-3 loci. The five species of Aythya constitute one of the least differentiated genera studied.

MERGINI (Eiders, Scoters, Mergansers and Allies): Somateria, Polysticta, Histrionicus, Clangula, Melanitta, Bucephala, Lophodytes, and Mergus. Patton and Avise (1986) identified relationships within Mergini and between Mergini and Aythyini as potential areas of disagreement between their genetic phylogeny and classical phylogenies. Because the phylogenetic relationships suggested by our analyses of fourteen species in eight Mergini genera are very similar to the classical phylogenies, only those of interest or which disagree are discussed.

The genetic relationships among the *Bucephala* (common goldeneye, Barrow's goldeneye, and bufflehead), *Histrionicus* (harlequin duck), *Somateria* (common eider and king eider) and two of the *Melanitta* (surf scoter and white-winged scoter) conform to accepted taxonomy. However, since *Melanitta nigra* (black scoter) and *Clangula hyemalis* (oldsquaw) are nearest relatives in the distance Wagner diagram (Fig. 2), a common ancestral divergence from the rest of the Mergini may be indicated.

While relationships in the phenogram (Fig. 1) among Lophodytes cucullatus (hooded merganser), Mergus merganser (common merganser), and Mergus serrator (red-breasted merganser) are in agreement with the current generic designations, this agrees in the distance Wagner diagram (Fig. 2) if Histrionicus histrionicus (harlequin duck) is included.

The *Polysticta stelleri* (Steller's eider) samples, shipped from Alaska, were unfortunately permitted to warm during transit, resulting in the loss of enzyme activity from several of the more labile enzymes. As a consequence of the partial loss of enzymes, Steller's eider was not included in the cluster analyses and, in turn, not placed on the distance diagrams (Figs. 1, 2 and 4). The species was included in Table 3 because it differed from the *Somateria* (common eider and king eider) by a fixation for alternate alleles at the *ADA*, *HEM*, and *MPI-1* loci.

OXYURINI (Stiff-tailed Ducks): Oxyura. Oxyura jamacensis (ruddy duck) was determined for alleles at




Figure 4. Means and ranges of genetic distances (Nel, 1978) at various taxonomic levels within the family Anatidae. Numbers of paired comparisons of species are given in parentheses.

the ADA, AAT-1, NP, GPI and PGM loci. It is observed as being the most divergent member of the Anatinae in the phenogram (Fig. 1), agreeing with classical taxonomy (Fig. 3) and the phylogenetic assignments of Johnsgard (1978) and Woolfenden (1961). Sibley and Monroe (1990) classify this tribe as the subfamily Oxyurinae, based on the DNA-DNA hybridization studies of Sibley and Ahlquist (1983, 1987, 1990). Oxyura joins the other members of the subfamily at a D of approximately 0.40 on the phenogram (Fig. 1). Oxyura shares a common root with Dendrocygna on the distance Wagner diagram (Fig. 2).

COMPARATIVE GENETIC DISTANCES

A summary of the means and ranges of genetic distances (distances (distances (distances)) at various taxonomic levels (Fig. 4) reveals considerable heterogeneity in interspecific genetic distances within genera (congeneric). Although genetic variation is approximately the same order of magnitude within bird species as that of other vertebrates, the degree of genetic differentiation among conspecific populations of birds is shifted to small values when compared to other vertebrates such as mammals and amphibians (Barrowclough, 1983)

DISCUSSION

Still pertinent (Bellrose, 1980) may be the early observations of Delacour and Mayr (1945) that "an over-evaluation of a few primary functional characters has led to ... confusion in the taxonomy of the Anatidae." With recent classifications attempting to reflect actual evolutionary relationships (Johnsgard, 1978; Livezey, 1986; Sibley and Ahlquist, 1982; Sibley and Monroe, 1990), the information gained in this study from the electrophoretic examination of allelic states will provide a sound basis with which to estimate phylogenetic relationships.

Comparative studies using electrophoretic data have shown that avian congeneric species are as close genetically to one another as are conspecific populations of other vertebrates (Avise et al., 1980a, 1980b, 1980c; Barrowclough and Corbin, 1978). Aquadro and Avise (1982) reported that, while the median D between species within 12 avian genera was 0.04, the same value was 0.40 for 32 non-avian genera. In this study of waterfowl, the average genetic distance between congeneric species, 0.08, supports the studies cited above (see also Patton and Avise, 1986).

Patton and Avise (1986) determined, after their analyses of the Anatidae, that close genetic similarity among members of the family was probably not due to recent divergence but rather to a deceleration in the rate of protein evolution relative to non-avian vertebrates (see also Barrowclough, 1983). Observed differences may become increasingly significant as a consequence of a decelerated rate of protein evolution.

The slowing of divergence by genetic drift, coupled with high effective population sizes, may contribute to the observation of reduced levels of avian genetic differentiation (Zink, 1986). Since geographic differentiation may be the cause of observed genetic distances which do not correlate with phenotypic similarities or differences (Barrowclough and Johnson, 1988), Avise et al. (1992) recommend that both the newer genetic and classical behavioral perspectives be considered in order to appreciate fully the geographic population structure of an avian species.

Consideration should also be given to the possibility of taxonomic over-splitting or excessive lumping (polyphyletic classification), which can often be identified by comparison of D values (Sibley and Ahlquist, 1982). Reduced values of mean D would be observed from taxonomic over-splitting, and increased values of mean D would stem from excessive lumping.

The multi-locus study of waterfowl by Patton and Avise (1986) examined electrophoretic variation at 19 loci in 26 species, with results supporting the relationships proposed by Delacour and Mayr (1945), Johnsgard (1968), and Morony et al. (1975). However, ambiguities were identified in the relationships among and between the tribes Mergini and Aythyini, leading to the suggestion of further analysis. In our study, based on electrophoretic variation at 25 loci in 40 species and 5 subspecies of waterfowl, we find that the genetic relationships among North American waterfowl agree, for the most part, with classical phylogenies (Johnsgard 1978) and taxonomic relationships (AOU, 1957, 1983, 1985, 1987, 1989, 1991) previously proposed.

The particular agreement of the phenogram (Fig. 1) to the classical phylogeny (Fig. 3) may reflect the phenetic nature of previous interpretations. In previous taxonomic studies, Brush (1976), Delacour and Mayr (1945), and Johnsgard (1961) based their conclusions upon observed character state similarities, interpreting their data empirically. Close concordance observed in the diverse approaches is not completely surprising. With waterfowl being subject to several recent classical taxonomic studies, relationships within the Anatidae will become more firmly established as the information base increases.

While the phenogram is observed to agree with classical phylogeny, it exhibits the greatest similarity at the subfamily and tribal levels. The distance Wagner clustering, however, exhibits greater similarity to the classical phylogeny at the species and generic levels. Meeting both criteria is the observed divergence of Chen canagica (emperor goose) from other members of the genus Chen. Chen canagica has been placed in three different genera: Philacte (AOU, 1957), Anser (Delacour and Mayr, 1945; Sibley and Monroe, 1990), and Chen (AOU, 1983). The results of our study do not support the placement of Chen canagica in either Anser or Chen.

Genetic differences were found among the emperor geese examined. The specimens were collected from Izembeck Lagoon, a region on the northern side of the Alaska peninsula, which serves as a staging area for emperor geese migrating north to nest. Emperor geese nest in only two areas of the world: northwestern Alaska and eastern Siberia. Those staging on the Alaska peninsula may represent a mixture of geese bound for both nesting areas (Eisenhauer and Kirkpatrick, 1978). Since our data indicate that two discrete breeding groups exist which are more divergent than many waterfowl species, the study of additional emperor geese is warranted.

A major discrepancy common to both branching diagrams is the inconsistent placement of *Clangula hyemalis* (oldsquaw) and *Melanitta nigra* (black sco-

ter). In the phenogram (Fig. 1), Clangula hyemalis does not cluster with the Mergini and Melanitta nigra does not cluster with the other Melanitta species. They jointly form a divergent Mergini branch in the distance Wagner diagram (Fig. 2). Johnsgard (1978) suspected a close affinity of the two species, and based these suspicions on comparisons of structure and behavior.

Considerable controversy exists as to the "best" method of deriving phylogenies from electrophoretic data. Although the phenogram and distance Wagner diagram are not entirely concordant, other branching diagrams may have been developed which might provide an even better fit to the existing data.

ACKNOWLEDGMENTS

This study was funded by the Federal Aid to Wildlife Restoration Act P-R Project W-38-R and the Nebraska Game and Parks Commission. Samples were obtained from the U.S. Fish and Wildlife Service in the states of Alaska, California, Colorado, Nebraska, and Washington. Analyses were performed at the Pacific Fisheries Research Laboratory in Washington with Drs. Jim and Lisa Seeb supervising activities and Paul Abersold assisting in electrophoretic analysis. Margo Ems, Ruth Wusk, Liz Huff, and Jan Bouc typed and prepared the text and tables for this study. Barbara Voeltz conducted the reference search.

LITERATURE CITED

- Allendorf, F. W. and F. M. Utter. 1976. Gene duplication in the family Salmonidae. II. Linkage between two loci coding for aspartate aminotransferase in cutthroat trout (Salmo clarki). Hereditas 82: 19–24.
 - ——, N. Mitchell, N. Ryman, and G. Stahl. 1977. Isozyme loci in brown trout (*Salmo trutta L.*): detection and interpretation from population data. *Hereditas* 86: 179–190.
- American Ornithologists' Union. 1957. Check-list of North American birds, Fifth edition. Baltimore, Maryland. American Ornithologists' Union.
- ——. 1983. Check-list of North American birds, Sixth edition. Baltimore, Maryland. American Ornithologists' Union.
- ——. 1985. Thirty-fifth supplement to the check-list of North American birds. The Auk 102: 680–686.
- ——. 1987. Thirty-sixth supplement to the check-list of North American birds. The Auk 104: 591–596.
- ——. 1989. Thirty-seventh supplement to the check-list of North American birds. The Auk 106: 532-538.
- ——. 1991. Thirty-eighth supplement to the check-list of North American birds. *The Auk* 108: 750–754.
- Ankney, C. D., D. G. Dennis, L. N. Wishard, and J. E. Seeb. 1986. Low genic variation between black ducks and mallards. The Auk 103: 701-709.
- Aquadro, C. F., and J. C. Avise. 1982. Evolutionary genetics of birds. VI. A reexamination of protein divergence using varied electrophoretic conditions. *Evolution* 36: 1003–

1019.

- Avise, J. C. 1994. Molecular markers, natural history, and evolution. New York, Chapman and Hall: 511pp.
- ——, R. T. Alisauskas, W. S. Nelson, and C. D. Ankney. 1992. Matriarchal population genetic structure in an avian species with female natal philopatry. *Evolution* 46: 1084–1096.
- ———, J. C. Patton, and C. F. Aquadro. 1980a. Evolutionary genetics of birds. I. Relationships among North American thrushes and allies. *The Auk* 97: 135–147.
- ———, and ———. 1980c. Evolutionary genetics of birds. Comparative molecular evolution in New World warblers and rodents. *Journal of Heredity* 71: 303–310.
- Baker, A. J. 1990. Population differentiation in colonizing species of birds. In B. D. Bell, R. O. Cossee, J. E. C. Flux, B. D. Heather, R. A. Hitchmough, C. J. R. Robertson, M. J. Williams (eds.), 20th International Ornithological Congress, Volume I. New Zealand, Ornithological Congress Trust Board: 504–512.
- Barrett, V. A., and E. R. Vyse. 1982. Comparative genetics of three trumpeter swan populations. *The Auk* 99: 103– 108.
- Barrowclough, G. F. 1983. Biochemical studies of microevolutionary processes. *In A. H. Brush*, and G. A. Clark, Jr. (eds.), *Perspectives in Ornithology*. New York, Cambridge University Press: 223–261.
- ———, and K. W. Corbin. 1978. Genetic variation and differentiation in the Parulidae. The Auk 95: 691–702.
- ——, and R. M. Zink. 1981. Genetic differentiation in the Procellariiformes. *Comparative Biochemical Physiology* 69B: 629–632.
- ——, and N. K. Johnson. 1988. Genetic structure of North American birds. In H. Ouellet (ed.), 19th International Ornithological Congress, Volume II. Ottawa, Canada, University of Ottawa Press: 1630–1638.
- Bellrose, F. C. 1980. Ducks, geese, and swans of North America, Third edition. Harrisburg, Pennsylvania, Stackpole Books: 540 pp.
- Boyer, S.H., D.C. Fainer, and M.A. Naughton. 1963. Myoglobin: inherited structural variation in man. *Science* 140: 1228–1231.
- Brown, I. R. F., W. H. Bannister, and C. DeLucca. 1970. A comparison of Maltese and Sicilian sparrow haemoglobins. *Comparative Biochemical Physiology* 34: 557–562.
- Browne, R. A., C. R. Griffin, P. R. Chang, M. Hubley, and A. E. Martin. 1993. Genetic divergence among populations of the Hawaiian duck, Laysan duck and mallard. *The Auk* 110: 49–56.
- Brush, A. H. 1976. Waterfowl feather proteins: analysis of use in taxonomic studies. *Journal of Zoology* 179: 467–498.
- ——. 1979. Comparison of egg-white proteins: effect of electrophoretic conditions. *Biochemical Systematic Ecology* 7: 155–165.
- Clayton, J. W. and D. N. Tretiak. 1972. Amine citrate buffers for pH control in starch gel electrophoresis. *Journal of Fisheries Research Board of Canada* 29: 1169–1172.
- Delacour, J. 1954–1964. The waterfowl of the world. Volumes 1–4. New York, Arco Publishing Company, Inc.

- ——, and E. Mayr. 1945. The family Anatidae. Wilson Bulletin 57: 3–55.
- Eisenhauer, D. I., and C. M. Kirkpatrick. 1978. Ecology of the emperor goose in Alaska. Wildlife Monographs 57: 62 pp.
- Farris, J. S. 1972. Estimating phylogenetic trees from distance matrices. *American Naturalist* 106: 645–668.
- ——. 1981. Distance data in phylogenetic analysis. In V. A. Funk and D. R. Brooks (eds.), Advances in cladistics. Proceedings of the Willi Hennig Society. Bronx, New York, New York Botanical Garden: 3–24.
- Felsenstein, J. 1981. Evolutionary trees from gene frequencies and qualitative characters: finding maximum likelihood estimates. *Evolution* 35: 1229–1242.
- Finnerty, V. and G. B. Johnson. 1979. Post-translational modification as a potential explanation of high levels of enzyme polymorphism: xanthine dehydrogenase and aldehyde oxidase in *Drosophila melanogaster*. *Genetics* 91: 695–722.
- ——, M. McCarron, and G. B. Johnson. 1979. Gene expression in *Drosophila melanogaster*: post-translational modification of aldehyde oxidase and xanthine dehydrogenase. *Molecular and General Genetics* 172: 37–43.
- Ford, H. A., A. W. Ewing, and D. T. Parkin. 1974. Blood proteins in Darwin's finches. Comparative Biochemical Physiology 47B: 369–375.
- Forey, P. L., C. J. Humphries, I. J. Kitching, R. W. Scotland, D. J. Siebert, and D. M. Williams. 1992. Cladistics: a practical course in systematics. Oxford, Oxford University Press: 191 pp.
- Gutierrez, R. J., R. M. Zink, and S. Y. Yang. 1983. Genic variation, systematic and biogeographic relationships of some galliform birds. The Auk 100: 33-47.
- Harris, H., and D. A. Hopkinson. 1976. Enzyme electrophoresis in human genetics. Amsterdam, North Holland Publishing Company.
- Hartl, D. L., and A. G. Clark. 1989. Principles of population genetics. Sunderland, Massachusetts, Sinauer Associates, Inc.: 682 pp.
- Harvey, W. D. 1990. Electrophoretic techniques in forensics and law enforcement. In D. H. Whitmore (ed.), Electrophoretic and isoelectric focusing techniques in fisheries management. Boca Raton, Florida, CRC Press: 313-312.
- Johnsgard, P. A. 1961. The taxonomy of the Anatidae—a behavioural analysis. *Ibis* 103a: 71–85.
- ——. 1968. Waterfowl. Lincoln, University of Nebraska Press: 138 pp.
- 1978. Ducks, geese, and swans of the world. Lincoln, University of Nebraska Press: 404 pp.
- Johnson, G., V. Finnerty, and D. Hartl. 1981. Post translational modification of xanthine dehydrogenase in a natural population of *Drosophila melanogaster*. Genetic Variation 98: 817–831.
- Johnson, N. K., R. M. Zink, and J. A. Marten. 1988. Genetic evidence for relationships in the avian family Vireonidae. *The Condor* 90: 428–445.
- Johnston, R. F., (ed.) 1983. Current ornithology. New York, Plenum Press: 425 pp.
- Livezey, B. 1986. A phylogenetic analysis of Recent anseriform genera using morphological characters. The Auk 103: 737–754.
- May, B., J. Wright, and M. Stoneking. 1979. Joint segregation of biochemical loci in Salmonidae: Results from ex-

- periments with Salvelinus and review of the literature on other species. Journal of Fisheries Research Board of Canada 36: 1114–1128.
- Mayr, E. 1991. *Principles of systematic zoology*. New York, McGraw-Hill, Inc.: 475 pp.
- Mickevich, M. R., and C. Mitter. 1981. Treating polymorphic characteristics in systematics: A phylogenetic treatment of electrophoretic data. *In* V. A Funk and D. R. Brooks (eds.), *Advances in cladistics*. Proceedings of the Willi Hennig Society. Bronx, New York, New York Botanical Garden: 45–60.
- Mindell, D. P., and J. W. Sites, Jr. 1987. Tissue expression patterns of avian isozymes: a preliminary study of phylogenetic applications. *Systematic Zoology* 36: 137–152.
- Morgan, R. P., L. A. Noe, and C. J. Henny. 1976. Biochemical identification of the mallard, *Anas platyrhynchos*, and black duck, *A. rubripes*. *Comparative Biochemical Physiology* 53B: 499–503.
- Morony, J. J., W. J. Bock, and J. Farrand, Jr. 1975. Reference list of birds of the world. New York, American Museum of Natural History.
- Nei, M. 1972. Genetic distance between populations. American Naturalist 106: 283–292.
- ——. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89: 583–590.
- Oates, D. W., J. E. Seeb, L. N. Wishard, and P. Abersold. 1983. *Biochemical identification of North American waterfowl*. Nebraska Technical Series, Number 13. Lincoln, Nebraska Game and Parks Commission.
- Patton, J. C. and J. C. Avise. 1986. Evolutionary genetics of birds. IV. Rates of protein divergence in waterfowl (Anatidae). Genetica 68: 129-143.
- Ridgway, G. J., S. W. Sherburne, and R. D. Lewis. 1970. Polymorphism in the esterases of Atlantic herring. Transactions of the American Fisheries Society 99: 147-151.
- Seeb, J. E., G. H. Kruse, L. W. Seeb, and R. G. Weck. 1990. Genetic structure of red king crab populations in Alaska facilitates enforcement of fishing regulations. Alaska Sea Grant College Program Report 90-04: 91-502.
- ——, L. W. Seeb, and F. M. Utter. 1986. Use of genetic marks to access stock dynamics and management programs for chum salmon. *Transactions of the American Fisheries Society* 115: 448–454.
- Selander, R. K., M. H. Smith, S. Y. Yang, W. E. Johnson, and J. B. Gentry. 1971. Biochemical polymorphism and systematics in the genus *Peromyscus*. II. Genic heterozygosity and genetic similarity among populations of the old-field mouse (*Peromyscus polionotus*). Studies in Genetics VI, University of Texas Publications 7103: 49–90.
- Shaklee, J.B., F.W. Allendorf, D.C. Morizot and G.S. Whitt. 1990. Gene nomenclature for protein-coding loci in fish. Transactions of the American Fisheries Society 119(1): 2-15
- Shaughnessy, P. D. 1970. Serum proteins of two sibling species of giant petrel (*Macronectes* spp.). Comparative Biochemical Physiology 33: 721–723.
- Shaw, C. R., and R. Prasad. 1970. Starch gel electrophoresis of enzymes: a compilation of recipes. *Biochemical Genetics* 4: 297–320.
- Sherman, P. W. 1981. Electrophoresis and avian genealogical analyses. *The Auk* 98: 419–422.
- Sibley, C. G. 1960. The electrophoretic patterns of avian egg-

- white proteins as taxonomic characters. *Ibis* 102: 215–284.
- ——. 1968. The relationships of the "wren thrush" *Zeledonia* coronata. Ridgeway. *Postilla* 125: 1–12.
- ——. 1970. A comparative study of the egg-white proteins of passerine birds. *Peabody Museum of Natural History Bulletin* 32: 1–131.
- ——, and J. E. Ahlquist. 1972. A comparative study of the egg-white proteins of non-passerine birds. *Peabody Museum of Natural History Bulletin* 39: 1–276.
- ———, and ———. 1982. The relationships of the yellow-breasted chat (*Icteria virens*) and the alleged slowdown in rate of macromolecular evolution in birds. *Postilla* 187: 1–18.
- ——, and ——. 1983. The phylogeny and classification of birds based on the data of DNA-DNA hybridization. *In* R. F. Johnston (ed.), *Current Ornithology*. New York, Plenum Press, 1: 245–292.
- ——, and ———. 1987. Avian phylogeny reconstructed from comparisons of the genetic material, DNA. *In C. Patterson (ed.)*, *Molecules and morphology in evolution: conflict or compromise.* New York, Cambridge University Press: 95–121pp.
- ———, and ———. 1990. Phylogeny and classification of birds of the world. New Haven, Yale University Press: 976 pp.
- ———, K. W. Corbin, and J. H. Haavic. 1969. The relationships of the flamingos as indicated by the egg-white proteins and hemoglobins. *Condor* 71: 155–179.
- ——, and B. L. Monroe, Jr. 1990. *Distribution and tax-onomy of birds of the world*. New Haven, Yale University Press: 1,111pp.
- Smith, J. K., and E. G. Zimmerman. 1976. Biochemical genetics and evolution of North American blackbirds, family Icteridae. *Comparative Biochemical Physiology* 53B: 319–324.
- Sneath, P. H. A., and R. R. Sokal. 1973. Numerical taxonomy. San Francisco, W. H. Freeman and Company: 573 pp.

- Straney, D. O. 1981. The stream of heredity: Genetics in the study of phylogeny. *In M. H. Smith and J. Soule (eds.)*, *Mammalian Population Genetics*. Athens, Georgia, University of Georgia Press: 100–138.
- Swofford, D. L. 1981. On the utility of the distance Wagner procedure. In V. A. Funk and D. R. Brooks (eds.), Advances in cladistics. Proceedings of the Willi Hennig Society. Bronx, New York, New York Botanical Gardens: 25-44.
- ——, and R. B. Selander. 1981. BIOSYS-1: A FORTRAN program for the comprehensive analysis of electrophoretic data in population genetics and systematics. *Journal of Heredity* 72: 281–283.
- Utter, F. W. 1991. Biochemical genetics and fishery management: an historical perspective. *Journal of Fish Biology* 39A: 1–20.
- ——, H. O. Hodgins, and F. W. Allendorf. 1974. Biochemical genetic studies of fishes: potentialities and limitations. In D. C. Malins and J. R. Sargent (eds.), Biochemical and biophysical perspectives in marine biology. London, Academic Press: 213–238
- Wilson, A. C. 1976. Gene regulation in evolution. *In F. J. Ayla (ed.), Molecular evolution*. Sunderland, Massachusetts, Sinauer Association, Inc.: 225–231.
- Woolfenden, G. E. 1961. Postcranial osteology of the waterfowl. Bulletin of the Florida State Museum of Biological Sciences 6: 1–129.
- Wright, S. 1978. Evolution and the genetics of populations. Volume IV. Variability within and among natural populations. Chicago, University of Chicago Press: 580 pp.
- Yang, S. Y. and J. L. Patton. 1981. Genic variability and differentiation in the Galapagos finches. The Auk 98: 230-242.
- Zink, R. M. 1982. Patterns of genic and morphologic variation among sparrows in the genera *Zonotrichia, Melospiza, Junco*, and *Passerella*. The Auk 99: 632–649.
- ——. 1986. Summary. In H. Ouellet (ed.), 19th International Ornithological Congress, Volume II. Ottawa, Canada, University of Ottawa Press: 1665–1668.