

1-1-1999

Development of Intervention Strategies to Extend the Shelf- Life of Fresh Ground Pork

David M. Gaebler

University of Nebraska - Lincoln

Roger W. Mandigo

University of Nebraska - Lincoln, rmandigo1@unl.edu

Follow this and additional works at: http://digitalcommons.unl.edu/coopext_swine



Part of the [Animal Sciences Commons](#)

Gaebler, David M. and Mandigo, Roger W., "Development of Intervention Strategies to Extend the Shelf- Life of Fresh Ground Pork" (1999). *Nebraska Swine Reports*. Paper 130.

http://digitalcommons.unl.edu/coopext_swine/130

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Swine Reports by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



DAY 2

- 7 a.m. Breakfast
- 7:30 a.m. Pork carcass evaluation and review of eight carcasses for fabrication
- 8 a.m. Pork slaughter demonstration
HACCP and microbial interventions
pH and other quality measurements
Measures of carcass composition
Hot boning demonstration
- 10:30 a.m. Pork carcass fabrication
- 12 p.m. Lunch
- 1 p.m. Value-added product demonstration including bacon, low-fat ham and fresh pork sausage
Demonstration of PSE and DFD pork processing
Marinated pork products

DAY 3

- 7 a.m. Breakfast
- 7:30 a.m. Pork Quality Assurance
Review of quality and consistency on carcasses
Taste panel evaluation of hog used in the demonstrations
Assessment of carcass value
Evaluation of cured products made the previous day
Carcass grading demonstration
- 12 p.m. Lunch
Adjourn

Course Evaluation

Past evaluations by participants indicate the course is successful and many participants had very positive remarks about it. When asked to identify things they liked about the course, the most popular answers related to “the hands-on nature of the course” and the “evaluation of the market hog from live to the meat products.”

¹Dennis E. Burson is an associate professor and Dana J. Hanson is a graduate student in the Department of Animal Science.

Development of Intervention Strategies to Extend the Shelf-Life of Fresh Ground Pork

David M. Gaebler
Roger W. Mandigo¹

Summary and Implications

The effects of storage time, packaging atmosphere and raw material source on shelf-life of fresh ground pork were studied. Fresh ground pork (18 percent fat) was packaged in an atmosphere of 80:20 percent O₂:CO₂ or 100 percent CO₂ and placed in unlighted refrigerated storage (34°F) for a period of two or eight days to simulate distribution time of the product from manufacturer to retail merchandiser. Products were then placed under lighted storage for eight additional days (100 foot candles, 34°F) to simulate retail display conditions. Ground sirloin had higher percent surface metmyoglobin (darkness and brown color) than ground pork shoulder after eight days of lighted storage. Lipid oxidation (rancidity) was higher in ground pork shoulder than ground pork sirloin. Pork shoulder had higher a (redness) values than pork sirloin in both atmospheres. Microbial loads (aerobic microorganisms) were higher in product stored eight days versus two days; however, total aerobic microbial loads did not exceed 10⁶ (the level commonly used to indicate microbial spoilage) for product stored in either atmosphere. Carbon dioxide successfully extends product shelf-life up to eight days under lighted storage conditions.*

Introduction

The preparation of meat products for retail display has changed dramatically over the last 20 years. Large supermarket chains have reduced or eliminated in-store preparation of fresh

red meat products to reduce labor and capital equipment costs. Today, meat products are prepared, packaged and labeled at large processing facilities, and shipped in refrigerated trucks to centralized distribution warehouses which then distribute products to individual stores. A consequence of this change is the necessity of longer shelf-life for products to reach the consumer. Fresh red meats packaged in oxygen permeable film have an expected shelf-life of two to three days under retail display conditions. One method for increasing shelf-life of refrigerated meats is to modify the atmosphere within the package. The shelf-life of fresh meat can be increased to six to 10 days with modified atmosphere packaging in a high-oxygen environment, and up to 21 days in a low-oxygen environment.

Modified Atmosphere Packaging (MAP) is one of several methods used by processors to control microbial spoilage of food products. Normal atmosphere contains 20.9 percent oxygen and 0.1 percent carbon dioxide. By increasing the carbon dioxide levels in the package, growth of aerobic spoilage organisms can be delayed, thereby extending the shelf-life of the meat product.

The two most frequently used atmospheres in MAP products are an 80:20 percent mixture of oxygen and carbon dioxide and 100 percent carbon dioxide. These atmospheres use different strategies to achieve the same result. The 80:20 percent O₂:CO₂ mixture uses higher-than-normal carbon dioxide levels to reduce aerobic microorganisms in combination with higher-than-normal oxygen levels to help maintain red meat color normally associated with freshness. The atmosphere is sealed into individual packages which

(Continued on next page)



are boxed and shipped to the retail store, ready for display.

The 100 percent CO₂ product is prepared in a traditional oxygen permeable packaging, placed into a master container which is filled with carbon dioxide and then sealed. Packages are removed from the master pack, allowed to bloom to a bright red color and placed on the shelf for sale.

The objectives of this research were to determine differences in characteristics of ground pork placed in lighted storage due to: effects of atmosphere (80:20 percent O₂:CO₂ or 100 percent CO₂), length of storage (two days versus eight days unlighted storage) and raw material source (pork sirloin with pork loin fat trim or pork shoulder meat).

Materials and Methods

Fresh pork sirloin, pork loin fat trim and pork shoulder meat were purchased from a commercial processor five days after slaughter. The meat was analyzed for fat content and fresh sirloin meat and pork loin fat trim were formulated to a fat content equal to the shoulder meat (18 percent fat). The pork was ground through a 1/8 inch plate, chilled and made into half-pound portions. Ground pork was placed into packages and filled with either 100 percent carbon dioxide or an 80:20 percent mixture of oxygen and carbon dioxide. Packages of ground pork were placed in unlighted storage (34°F) for either two or eight days to simulate the range of time prepackaged meat would spend in route to retail outlets. Packaged products were then placed under light (1076 lux light, 34°F) for eight days to simulate lighting conditions found in retail display cases. The products were kept in their original atmospheric environment throughout the study. Individual packages were opened immediately prior to chemical and physical analysis. Product was evaluated every two days (zero to eight days of lighted storage) for total bacteria (APC), coliforms, psychrotropic bacteria, exterior color L* (lightness), a* (redness) and b* (yellowness) values, surface metmyoglobin and surface




Figure 1. Surface metmyoglobin of ground pork in 80:20 percent oxygen:carbon dioxide modified atmosphere packaging.




Figure 2. Surface metmyoglobin of ground pork in 100 percent carbon dioxide for product held either two or eight days in unlighted storage prior to lighted retail display.

metmyoglobin reducing ability (MRA), pH and oxidation reduction potential. Lipid oxidation was also measured and reported as thiobarbituric acid reactive substances (TBARS). Surface metmyoglobin formation was measured by obtaining reflectance readings through eight days of lighted storage.

Results and Discussion

In a high-oxygen environment (80:20 percent oxygen:carbon dioxide) surface metmyoglobin was higher for ground pork from sirloin than for ground pork shoulder meat (Figure 1) in product held two or eight days (unlighted storage) prior to lighted retail display. Surface metmyoglobin levels remained below 50 percent

throughout eight days of lighted retail storage (50 percent metmyoglobin is the value commonly used as the level above which consumers reject meat for purchase based on color perception). Surface metmyoglobin increased during lighted storage from 27.5 percent to 31 percent (sirloin) and 31 percent to 37.5 percent (shoulder) after two days of unlighted storage and from 31 percent to 33 percent (sirloin) and 28 percent to 37.5 percent (shoulder) after eight days of unlighted storage. Surface metmyoglobin of products in an atmosphere of 100 percent CO₂ decreased over time from 30 percent to 23 percent (after two days of unlighted storage) due to enzymatic reduction of the meat system, but remained relatively constant at 23 percent after eight




Figure 3. Metmyoglobin reducing ability of ground pork in modified atmosphere packaging.




Figure 4. Changes in surface redness due to different meat source and atmosphere.

Table 1. Fatty acid profile of ground pork.

Fatty Acid	Ground Pork Sirloin		Ground Pork Shoulder	
	Mean (n=8)	SEM ^a	Mean (n=8)	SEM ^a
Mystiric Acid (14:0)	1.40	0.14	1.40	0.06
Palmitic Acid (16:0)	23.23	0.56	22.93	0.72
Palmitoleic (16:1)	3.30	0.22	3.94	0.19
Stearic (18:0)	12.65	0.52	11.67	0.30
Oleic (18:1)	43.48	0.84	45.16	0.55
Linoleic (18:2)	15.48	0.44	14.45	0.76
Arachidonic (18:3)	0.46	0.09	0.55	0.13
Total	100.00		100.00	

^aStandard error of the mean.

days of unlighted storage (Figure 2). Metmyoglobin reducing ability of ground pork packaged in 100 percent carbon dioxide was higher than ground pork packaged in 80:20 percent oxygen:carbon dioxide (Figure 3). In an anaerobic environment, enzymes will reduce metmyoglobin to deoxymyoglobin, reducing the level of surface metmyoglobin (Figure 2) over time, as long as reducing equivalents are not depleted.

Lipid oxidation of ground pork is measured as thiobarbituric acid reactive substances (TBARS). A TBARS value of 1.0 is considered the threshold value for consumers to detect rancidity in fresh ground pork. Ground shoulder meat was higher in unsaturated fatty acids (palmitoleic, oleic and arachidonic fatty acids) than ground sirloin (Table 1). Unsaturated fatty acids contribute to higher lipid oxidation and correspondingly higher TBARS values. The TBARS values for ground pork in 100 percent carbon dioxide remained below 0.3 mg malonaldehyde/kg meat throughout lighted storage and there were no significant differences between meat sources (data not shown). Surface a* values (redness) were higher for pork shoulder than for pork sirloin (Figure 4) for both atmospheres.

Total aerobic plate counts, coliforms and psychotropic bacteria counts are given in Tables 2 through 4. In an 80:20 percent O₂:CO₂ atmosphere, products held for eight days in unlighted storage had higher bacterial levels than products held for two days, although levels remained below 10⁶ throughout eight days of lighted storage. Aerobic plate counts were lower for products stored in 100 percent CO₂ and remained low throughout lighted storage (Table 2). Carbon dioxide has an inhibitory effect on the growth of microorganisms by both extending the lag phase of bacteria prior to the growth phase and decreasing pH due to increased solubility of carbon dioxide in meat at low temperatures. Psychotropic plate counts and coliforms remained low throughout lighted storage in both atmospheres (Tables 3 and 4).

(Continued on next page)



Table 2. Mean aerobic plate counts for ground pork in modified atmosphere packaging.

Treatment		Aerobic plate counts (log ₁₀ cf.u./gram)					SEM	
		Day 0	Day 2	Day 4	Day 6	Day 8		
80:20 O ₂ :CO ₂	2-day unlighted storage	3.27 ^a	3.25 ^a	3.41 ^{ab}	3.81 ^{bc}	3.94 ^c	0.12	
	8-day unlighted storage	3.95 ^c	4.14 ^d	4.65 ^e	5.22 ^f	5.51 ^g	0.12	
	Pork sirloin	3.38 ^a	3.59 ^b	3.96 ^c	4.36 ^d	4.63 ^e	0.09	
	Pork shoulder	3.85 ^c	3.80 ^{cd}	4.11 ^{cd}	4.67 ^e	4.82 ^e	0.09	
100% CO ₂	Pork sirloin	2-day unlighted storage	3.19 ^w	3.05 ^w	3.14 ^w	3.17 ^w	3.13 ^w	0.06
		8-day unlighted storage	3.15 ^w	3.21 ^w	3.26 ^w	3.63 ^y	4.00 ^z	0.06
	Pork shoulder	2-day unlighted storage	3.37 ^x	3.34 ^{wx}	3.41 ^x	3.51 ^x	3.47 ^x	0.06
		8-day unlighted storage	3.49 ^{xy}	3.33 ^{wx}	3.58 ^y	3.49 ^y	3.56 ^y	0.06

^{a-g,w-z} Means within an atmosphere treatment with the same subscript are not significantly different (P<0.05).

Table 3. Mean psychotropic plate counts for ground pork stored in modified atmosphere packaging.

Treatment		Psychotropic plate counts (log ₁₀ cf.u./gram)					SEM	
		Day 0	Day 2	Day 4	Day 6	Day 8		
80:20 O ₂ :CO ₂	2-day unlighted storage	2.79 ^a	3.33 ^b	3.42 ^b	13.68 ^c	4.31 ^{de}	0.08	
	8-day unlighted storage	4.09 ^d	4.39 ^e	5.11 ^f	5.61 ^g	5.87 ^b	0.08	
100% CO ₂	Sirloin	2-day unlighted storage	2.39 ^w	2.62 ^{wx}	2.52 ^w	2.42 ^w	2.68 ^{wx}	0.14
		8-day unlighted storage	2.06 ^{wx}	2.52 ^w	3.23 ^y	3.49 ^y	4.11 ^z	0.14
	Shoulder	2-day unlighted storage	2.78 ^{wx}	3.32 ^y	3.10 ^y	2.59 ^{wx}	2.85 ^{xy}	0.14
		8-day unlighted storage	2.91 ^{xy}	2.52 ^w	3.27 ^y	3.33 ^y	3.61 ^y	0.14

^{a-d,w-z} Means within an atmosphere treatment with the same letter are not significantly different (P<0.05).

Table 4. Mean coliform plate counts for ground pork stored in modified atmosphere packaging.

Treatment		Coliforms (c.f.u./gram)					SEM
		Day 0	Day 2	Day 4	Day 6	Day 8	
80:20 O ₂ :CO ₂	2-day unlighted storage	15 ^a	14 ^a	20 ^a	16 ^a	12 ^a	1.15
	8-day unlighted storage	12 ^a	51 ^b	19 ^a	20 ^a	18 ^a	1.15
	Pork Sirloin	13 ^a	52 ^d	38 ^{cd}	35 ^c	22 ^b	1.15
	Pork Shoulder	14 ^a	13 ^a	10 ^a	<10 ^a	<10 ^a	1.15
100% CO ₂	2-day unlighted storage	14 ^w	11 ^x	<10 ^x	<10 ^x	<10 ^x	1.11
	8-day unlighted storage	<10 ^x	<10 ^x	<10 ^x	<10 ^x	<10 ^x	1.11

^{a-b,w-x} Means within an atmosphere treatment with the same letter are not significantly different (P<0.05).

Conclusions

The generally expected shelf-life of red meat packaged in oxygen permeable films is two to three days. Modified atmosphere packaging was successful in extending shelf-life of ground pork from two to three days of lighted storage to at least six days for color and up to eight days for microbial

spoilage. Ground pork shoulder meat had greater redness than ground pork sirloin meat. Microbial spoilage of ground pork was maintained below spoilage levels for up to eight days of lighted display. Ground pork packaged in an atmosphere of 80:20 percent O₂:CO₂ can be held in unlighted storage for up to eight days and achieve an additional six to eight days of lighted

storage shelf-life, provided ground pork is produced from freshly slaughtered meats and processed in a clean, sanitary environment with good temperature control.

¹David M. Gaebler is a graduate student and Roger W. Mandigo is a professor in the Department of Animal Science.