University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Range Beef Cow Symposium

Animal Science Department

December 1999

Can Cow Adaptability and Carcass Acceptability Both Be Achieved?

R. D. Green

Colorado State University

T. G. Field

Colorado State University

N. S. Hammett

Colorado State University

B. M. Ripley

Colorado State University

S. P. Doyle

Colorado State University

Follow this and additional works at: <https://digitalcommons.unl.edu/rangebeefcowsymp>

Part of the [Animal Sciences Commons](#)

Green, R. D.; Field, T. G.; Hammett, N. S.; Ripley, B. M.; and Doyle, S. P., "Can Cow Adaptability and Carcass Acceptability Both Be Achieved?" (1999). *Range Beef Cow Symposium*. 130.

<https://digitalcommons.unl.edu/rangebeefcowsymp/130>

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Range Beef Cow Symposium by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

**CAN COW ADAPTABILITY AND CARCASS ACCEPTABILITY
BOTH BE ACHIEVED?**

R.D. Green, T.G. Field, N.S. Hammett, B.M. Ripley, and S.P. Doyle
Department of Animal Sciences
Colorado State University

The Beef Industry Is Lost in its Own Fog?

Over the past couple of decades, the beef cattle industry has become a confusing place to exist. Messages have been conveyed to producers at a fast and furious pace. This would not be a problem if these messages were consistent and if they were compatible with each other, yet this is far from the real situation. Daryl Tatum has been known to occasionally coin the term to describe confusion as “someone being lost in his/her own fog”. Unfortunately, this verbage very accurately describes the beef cattle production environment of the 1990s.

One could compose a lengthy list of dichotomies in the current beef cattle industry. A partial list might include:

- 1) A forage-based production system (low input) versus a concentrate-based feedlot system (high input).*
- 2) Fierce pride in producer individuality and independence versus strategic alliances and cooperative relationships.*
- 3) Segmentation and resulting inefficiencies versus vertical coordination and/or integration.*
- 4) Traditional purebred cattle-focused seedstock production versus commercially-oriented specification seedstock production.*
- 5) “Show cattle” versus “performance cattle”.*
- 6) Commodity-based marketing versus value-based marketing.*
- 7) Totally “vested” beef cattle producers versus “less-vested” small land-owners.*
- 8) Public lands versus private land use for beef production.*
- 9) “Artificially selected” cattle versus “naturally selected” cattle.*
- 10) “Animal welfare” versus “animal rights”.*
- 11) Systematic crossbreeding versus mongrelization.*
- 12) Purebred breeding versus composite breeding.*
- 13) Increased quality and consistency versus increased genetic variation.*
- 14) High tech production versus low cost production.*
- 15) Matching the cow to the production environment versus matching the calf to the marketing environment (i.e. cow adaptability versus carcass acceptability).*

The collective concerns and issues listed above, along with a number of others we could further list, have contributed to “the fog” for beef cattle producers. Given how spontaneously and explosively these issues can appear (or increase in importance), what is a cow-calf producer to do? The objectives of this presentation are to: 1) Provide an overview of how to match a

producer and production system to a specific industry target; 2) Discuss current and future tools needed for proper genetic decision-making, and 3) Provide some perspective on how the beef cattle industry can go about increasing “quality and consistency” while maintaining balances in cow herd efficiency and profitability.

The Beef Industry Definitely Has a #1 Leverage POINT

The U.S. National Beef Quality Audit (NBQA), conducted by Colorado State and Texas A&M Universities in 1991, was a “wake-up” call to the beef cattle industry. The results of the audit of the slaughter cattle population, conducted in 28 plants across the U.S., indicated a total of some \$280 in inefficiencies for each fed steer and heifer produced in the beef business. Furthermore, when these inefficiencies were categorized, it became apparent that the majority of losses occurred due to excess fat production with lower consistency in taste than desired. These results indicated that the beef cattle industry needed to achieve change at the carcass level, by implementing a combination of changes in feeding and management practices coupled with genetic improvement.

The executive summary of the 1991 NBQA stated:

“Some cattle are born to be misfits. Examples include cattle that are light-muscled, cattle that must be fed to the fattest end of yield grade 3 to grade Select, and cattle that must be fed to 1,500 lb live weights to reach desirable muscle:bone ratios. The beef industry would be well-served to identify such cattle and eliminate their parents from the breeding herd.”

In addition, one of the strategies for improving the quality of beef identified by the consensus panel of the 1991 NBQA strategy workshop was to “eliminate biological types (not breeds per se) of cattle that fail to conform”. This statement and philosophy brought the term “genetic non-conformance” into vogue. Since that time, considerable discussion has taken place regarding how to genetically better hit the industry targets for end-product performance. Everything from eliminating breeds (or the more politically correct vernacular of “eliminating specific biological types”), to combining breeds into composites, to making a new industry standard of 50% British: 50% Continental slaughter cattle has been argued.

The NBQA was repeated again in 1995 and guess what was found? A few management effects were different, cattle were less finished due to market conditions, and consequently quality grades had declined. This resulted in a shift in recommendations away from decreasing fat (i.e. 1991) to increasing quality and consistency (i.e. 1995). As a result, the production segment of the industry responded by saying....how are we supposed to shoot for a target when the bull’s eye keeps changing?

There are several reasons why little genetic change occurred in the cattle population over those five years, principally: 1) time -- five years does not quite constitute a single generation interval of beef cattle, 2) most of the necessary genetic tools for identifying specific seedstock animals for specific roles (in relation to end-product performance) were still under development,

3) these genotypes, in most cases, were filling a perceived useful niche in the cow-calf sector, 4) genetics which were being inappropriately used in 1991 were still being inappropriately used in 1995, 5) there has been a long-standing perception that production of cattle with “value-based genetics” will not receive “value-based rewards”, and 6) there has been concern relative to the effects of genetic change in carcass performance on raising rather than lowering production costs.

What about Lowering Costs of Production?

All changes in a commercial cow-calf operation must be evaluated in terms of their effect on profitability of the whole enterprise. Given the problem that profitability is often, in the short-term, very affected by external market conditions, Dickerson (1984) advocated that these changes be evaluated on the basis of economic efficiency measured as the ratio of input costs per unit of output product value. When one operates under this philosophy, cost of production becomes very important relative to desired increases in product value mentioned above as industry goals. Furthermore, it is imperative to remember that many of these desired ends are often antagonistically related, meaning that we must be careful to keep the “big picture” in perspective.

For example, traditionally we have thought that in relative economic terms, reproductive efficiency is roughly twice as important as growth performance which is approximately five times as important as carcass merit (Melton et al., 1979). A few years ago, a reanalysis of the importance of these three types of traits under a more current, value-based type of marketing system was completed. Under this more current marketing system, the former 10 reproduction: 5 growth: 1 product ratio was now closer to 2 reproduction: 1 growth: 1 product (Melton, 1995). A more recent evaluation of these economic weights has been presented from the American Gelbvieh Association’s Gelbvieh Alliance marketing program (Figure 1). After some 110,000 feedlot cattle had gone through their program, the estimated relative importance of these three trait categories was approximately 4:2:1 (Schiefelbein, 1998). There are several things about these relative economic values that are very important. First, under the general assumptions used in their derivation, these results indicate that while we have paid a lot of attention to growth of calves in the past, that will not suffice in the future. In most cases, the problems are in the other two categories: reproductive efficiency because it has been so difficult to genetically change, and carcass merit because we simply have not paid much attention to this area.

Figure 1. Relative importance of trait categories (Schiefelbein, 1998).

Secondly, one should not fall prone to the common error of assuming that these economic weights are universally true. They are applicable to one particular system and environment but may be quite different if the system is changed. One of the universal strengths that makes the beef industry unique is that it uses God-given resources from the land which cannot be more efficiently utilized by other production systems. Cattle harvest energy from sunlight, soil, and water that is then converted to a higher quality form of protein. They do this from a set of natural resources that cannot be “farmed” any other way. The problem is that those resources exist under such a wide array of ecosystems that we are challenged to come up with one management system that will work for all environments (Hohenboken, 1988). Herein also lies our genetic dilemma when we try to build the best beast to harvest and harness that energy from the environment.

Thirdly, we also often tend to over-generalize in the beef industry when talking about “THE TARGET”. As Dell Allen of Excel, Inc. has stated, there are several different target markets in our industry (Allen, 1987). The first question that a commercial producer must ask before addressing anything else genetically, is “Which target am I going to aim my production resources toward?”. As marketing of cattle in alliance and grid programs has escalated over the past 24 months, it has become clear that there are major targets in “lean beef”, “high-quality beef”, and “export beef” trade. There are certainly other smaller specialty markets as well. The market may change over time in relation to premiums and discounts for “leanness” versus “quality”. However, a given producer must decide *before the genetic decisions are made* on a well-defined target that is comfortable. Given the current plethora of alliance marketing programs, one must become educated on where he/she fits and then set their target based on that marketing program. *Only then can one truly go about determining the relative importance of these traits.*

What Should Be in a Commercial Producer’s Want Ad?

In 1987, the beef cattle symposium program at the annual meeting of the American Society of Animal Science was entitled “Bovine Nirvana”. In that program, Rick Bourdon and Bill Hohenboken discussed different perspectives on how one might describe the “ideal” cow (Bourdon, 1988; Hohenboken, 1988). As they both stated in their remarks, this beast does not really exist, primarily due to the reasons already described earlier in this paper. However, we do know that it is possible to provide some general guidelines for the specifications we would look for in performance criteria in the beef cattle production system.

Bob Taylor, our recently deceased colleague and friend, had great foresight in realizing the need to look at “balanced” performance of cattle long before it was popular. A number of years ago, he developed a simple analogy to illustrate the importance of this philosophy to his beef production students. He said that what commercial producers should do is develop a “want ad” for the type of bulls and females they use in their system. This want ad should then be what is used by the seedstock industry to develop “specification seedstock” to address the needs of the commercial production sector of the industry. While this is a very simple approach, in concept, one is left to wonder just how often it has been applied. Taylor’s generalized want ad, shown in Table 1, provides an excellent overview of the challenge a breeder has to mount in order to “hit the overall” target.

Table 1. Production and marketing specifications for beef cattle.

Trait	Optimum Range ^a	Industry Target ^b
<u>Reproduction</u>		
Age at Puberty (mos)	12-16	14
Scrotal Circumference (cm)	32-40	36
Reproductive Tract Score (14 mos)	4-5	5
Weight at Puberty (kg)		
Heifers	270-360	320
Bulls	400-500	450
Age at First Calving (mos)	23-25	24
Birth Weight		
Calves from Cows (kg)	35-45	39
Calves from Heifers (kg)	27-36	32
Body Condition Score (BCS, 1-9)	4-6	5
Postpartum Interval (d)	55-95	75
Calving Interval (d)	365-390	365
Calving Season (d)	45-90	65
Calf Crop Weaned (% cows exposed)	80-95	85
Cow Longevity (yr of age)	9-15	12
<u>Growth</u>		
Mature Cow Weight (kg at BCS 5)	400-600	500
Weaning Weight (kg; steer @ 7 mos)	200-275	240
Yearling Weight (kg; steer @ 365 d)		
Grazed and/or backgrounded	275-365	320
Weaning to Feedlot	400-500	450
Feedlot Gain (kg/d)	1.1-1.6	1.4
Feedlot Feed Efficiency (steer)	5-7 ^c	6 ^c
Days on feed (high energy ration)	60-120	90
<u>Carcass</u>		
Carcass Weight (kg)	275-365	320
Quality Grade	Select ⁺ to Choice ⁺	Choice ⁻
USDA Yield Grade	1.5-3.5	2.5
Fat Thickness (mm)	2.5-15.2	7.5
Ribeye Area (cm ²)	71-97	84
Palatability (% fat in retail cuts)	3-7	5
Warner-Bratzler Shear Force (kg)	Below 3.65	Below 3.65
Muscle to Bone (kg muscle to kg bone)	3.5-4.5	4.0
Lean Yield/Day Age (kg)		
Weaned Steer to Feedlot	0.35-0.45	0.40
Grazed Yearling Steer to Lot	0.20-0.30	0.25
Frame Score		
Steers	4-6	5
Cows	4-6	5
Bulls - Maternal	4-6	5
Terminal	5-7	6

(Adapted from Taylor and Field, 1999).

^aRange will include most commercial beef operations where an optimum combination of productivity and profitability is desired.

^bTarget gives central focus applicable to many commercial beef operations. Deviation from this target and optimum range is dependent on market, economic, and environmental conditions in specific commercial beef operations.

^cHigh-energy ration, kg feed per kg gain.

Is it Possible to Genetically Improve Cow Adaptability and Carcass Acceptability?

Within Population Selection. Fortunately, collective research results over the past 50 years have clearly shown that genetic variation exists both between and within breeds for many of the important measures of performance in beef cattle production. Table 2 provides a summary of the average levels of heritability for a variety of reproductive, growth and carcass traits as provided in an exhaustive analysis of the research literature by Koots et al. (1994a). In general, selection within breed populations is quite effective for carcass traits, moderately effective for growth related traits, and much slower for reproductive efficiency related traits.

Until recently, we have believed that there was limited opportunity to genetically improve fertility via direct selection within breeds. While indicator traits of fertility and age at puberty, such as scrotal circumference, have proven to be quite useful and heritable (Brinks et al., 1996), they have not been shown to be highly genetically correlated to fertility measured as pregnancy success. Because fertility measures are binary traits (i.e. they are observed as either pregnant or not pregnant), it is quite difficult to use phenotypic information to determine genetic differences (eg. two females may both get pregnant but may differ widely in their true genetic potential for fertility). This results in traditional analytical methods not being adequate to separate these genetic differences and thus, we have always stated that the heritability of these traits is quite low (see Table 2). More appropriate statistical methodology called “threshold modeling” allows appropriate analysis of these types of traits on an underlying continuous probability scale. One of the first applications this approach was to define a new trait called “stayability” that has been adopted by the Red Angus Association of America and is now in the process of being implemented by several other breeds (Snelling et al., 1995). This estimated breeding value is a genetic prediction of the probability of females still being in the herd at a breakeven age of six years given that they were selected as replacements. This measure combines performance differences in fertility, growth, and survivability/adaptability of these females.

In the direct fertility area, an analysis of heifer pregnancy records from the Hereford herd at the Bell Ranch in New Mexico has recently been completed (Evans et al., 1996). In that study, the researchers determined that heifer pregnancy was indeed more heritable than previously thought (14%). Furthermore, when the relationship of heifer pregnancy with yearling bull scrotal circumference was estimated, a non-linear relationship (i.e. the bulls with low SC EPD had low HP EPD, moderate SC EPD had the highest HP EPD and highest SC EPD had lower HP EPD) was revealed. A second study conducted a similar analysis using historical data from the Colorado State University Beef Improvement Center Angus population at Saratoga, WY (Doyle et al., 1996). These researchers reported a heritability level for heifer pregnancy of 19%, corroborating the result of Evans et al. (1996). These two studies indicate that it is feasible to produce genetic predictions to enable direct genetic improvement in reproductive rate. The only obstacle is getting breed association national cattle evaluation performance databases to adopt a “whole-herd reporting” format that is necessary to allow computation of these types of EPD (Golden et al., 1996). While this is only a start on the whole reproductive efficiency complex, it is a 200% improvement over current genetic capabilities in this important area.

Table 2. Levels of heritability (h^2) of beef cattle performance traits.

Trait	Number of Studies ^a	Weighted Mean h^2 ^b
<u>Reproduction</u>		
Age at First Calving (Direct)	7	6
Age at First Calving (Maternal)	1	19
Calving Date	7	8
Calving Interval (Cows)	3	1
Calving Interval (Heifers)	7	6
Calving Ease (Direct)	19	10
Calving Ease (Maternal)	11	9
Calving Rate	9	17
Scrotal Circumference	25	48
Heifer Conception Rate (Direct)	9	5
Heifer Conception Rate (Maternal)	1	2
Cow Conception Rate (Direct)	21	17
Cow Conception Rate (Maternal)	1	2
<u>Growth</u>		
Birth Weight (Direct)	167	31
Birth Weight (Maternal)	34	14
Weaning Weight (Direct)	234	24
Weaning Weight (Maternal)	38	13
Yearling Weight (Direct)	147	33
Yearling Weight (Maternal)	6	6
Mature Cow Weight	24	50
Feed Efficiency	25	32
Feed Intake	21	34
Relative Growth Rate	12	22
<u>Carcass</u>		
Backfat Thickness	26	44
Ribeye Area	16	42
Slaughter Weight	52	41
Carcass Weight	19	23
Dressing Percentage	13	39
Cutability	12	47
Lean:Bone Ratio	4	63
Marbling Score ^c	12	38
Warner-Bratzler Shear Force	12	29
Sensory Panel Tenderness	3	13
Yearling Frame Score	27	61

(Adpated from Koots et al., 1994a and Green, 1999).

^aNumber of research studies represented.

^bAverage heritability of trait, weighted by number of observations in studies. Expressed as a percentage.

^cRecent review of Marston et al. (1999) reported average of 43% heritability for marbling.

^dAll traits are expressed on an age constant basis where applicable.

Between Population Selection. Larry Cundiff and co-workers at the U.S. Meat Animal Research Center have conducted the most extensive genetic evaluation of breeds in the world over the past 30 years in the Germ Plasm Evaluation (GPE) program at the U.S. Meat Animal Research Center. The design for this project (Table 3) has allowed for the evaluation of a widely diverse set of breeds, as shown grouped by biological type in Table 4 (Cundiff and Gregory, 1999). From the collective results of this effort, they have reported that the magnitude of genetic variability between breeds is roughly equivalent to that within breeds (Table 5) for most performance traits. While this infers that genetic improvement is possible through proper breed selection implemented in designed crossbreeding programs (i.e. breed complementarity), it also points out that no one breed excels in all characteristics simultaneously, along with a great degree of overlap between various breeds.

Table 3. Sire breeds used in the germ plasm evaluation program (Cundiff and Gregory, 1999).

Cycle I (1970-72)	Cycle II (1973-74)	Cycle III (1975-76)	Cycle IV (1986-90)	Cycle V (1992-94)
<u>F₁ crosses from Hereford or Angus dams (Phase 2)^a</u>				
Hereford	Hereford	Hereford	Hereford	Hereford
Angus	Angus	Angus	Angus	Angus
Jersey	Red Poll	Brahman	Longhorn	Tuli
S. Devon	Braunvieh	Sahiwal	Salers	Boran
Limousin	Gelbvieh	Pinzgauer	Galloway	Belgian Blue
Simmental	Maine Anjou	Tarentaise	Nellore	Brahman
Charolais	Chianina		Shorthorn	Piedmontese
			Piedmontese	
			Charolais	
			Gelbvieh	
			Pinzgauer	
<u>3-way crosses out of F₁ dams (Phase 3)</u>				
Hereford	Hereford			
Angus	Angus			
Brahman	Brangus			
Devon	Santa Gertrudis			
Holstein				

^aIn Cycle V, composite MARC III (1/4 Angus, 1/4 Hereford, 1/4 Pinzgauer and 1/4 Red Poll) cows are also included.

^bHereford and Angus sires used in Cycle IV included 10 Hereford sires born from 1963 to 1969 and 14 Angus sires born from 1968 to 1970 used as reference sires in Cycles I, II, III and IV to produce reciprocal cross Hereford x Angus (HA_o) progeny, and 32 Hereford sires born from 1982 to 1985 and 28 Angus sires born from 1983 to 1985 used to produce reciprocal cross Hereford x Angus by a current sample of sires (HA_c) in Cycles IV, and as reference sires in Cycle V.

Table 4. Breeds evaluated in USDA-ARS germ plasm evaluation program grouped into biological type (Cundiff and Gregory, 1999).

Breed Group	Growth Rate/ Mature Size	Lean to Fat Ratio	Age of Puberty	Milk Production
Jersey (J)	X	X	X	XXXXXX
Longhorn (Lh)	X	XXX	XXX	XX
Hereford-Angus (HAX)	XXX	XX	XXX	XX
Red Poll (R)	XX	XX	XX	XXX
Devon (D)	XX	XX	XXX	XX
Shorthorn (Sh)	XXX	XX	XXX	XXX
Galloway (Gw)	XX	XXX	XXX	XX
South Devon (Sd)	XXX	XXX	XX	XXX
Tarentaise (T)	XXX	XXX	XX	XXX
Pinzgauer (P)	XXX	XXX	XX	XXX
Brangus (Bg)	XXX	XX	XXXX	XX
Santa Gert. (Sg)	XXX	XX	XXXX	XX
Sahiwal (Sw)	XX	XXX	XXXXX	XXX
Brahman (Bm)	XXXX	XXX	XXXXX	XXX
Nellore (N)	XXXX	XXX	XXXXX	XXX
Braunvieh (B)	XXXX	XXXX	XX	XXXX
Gelbvieh (G)	XXXX	XXXX	XX	XXXX
Holstein (Ho)	XXXX	XXXX	XX	XXXXX
Simmental (S)	XXXXX	XXXX	XXX	XXXX
Maine Anjou (M)	XXXXX	XXXX	XXX	XXX
Sales (Sa)	XXXXX	XXXX	XXX	XXX
Piedmontese (Pm)	XXX	XXXXXX	XX	XX
Limousin (L)	XXX	XXXXX	XXXX	X
Charolais (C)	XXXXX	XXXXX	XXXX	X
Chianina (Ci)	XXXXX	XXXXX	XXXX	X

^aIncreasing number of X's indicate relatively higher values.

Table 5. Relativity of variation within and between breeds for various performance criteria.

Trait	Number of Additive Genetic Standard Deviations
Age at Puberty (d)	8.5
Slaughter Weight (450 d)	8.0
Retail Product Weight (450 d)	8.2
Retail Product % (450 d)	6.6
Marbling Score (450 d)	6.1
Warner-Bratzler Shear Force (kg)	5.1

Adapted from Cundiff and Gregory (1999).

^aAssumption is made here that within a breed approximately six genetic standard deviations of variation exist in any trait.

The GPE program, along with other studies, has also shown that many genetic antagonisms exist in beef production systems. Koots et al. (1994b) summarized published estimates of genetic and phenotypic correlations between a number of traits of interest (Table 6). These estimates clearly reveal general genetic antagonisms between growth rate and calving ease, growth rate and mature cow size, maternal characteristics and cutability, and carcass quality and cutability. Additionally, the review of these authors pointed out how many genetic relationships between traits of economic importance are poorly understood. A prime example of the sparseness of information is the lack of any understanding of the relationship between measures of tenderness and other performance criteria.

The most troubling genetic antagonism we must consider when attempting to genetically improve product quality and consistency concerns the relationship between carcass attributes and measures of reproductive efficiency. There is generally a lack of this type of information in the research literature. The best existing data relating actual carcass measures to reproductive traits comes from a study by MacNeil et al. (1984) at the U.S. Meat Animal Research Center. Table 7 provides a summary of that information and indicates antagonistic relationships between selection to increase retail product weight and age at puberty, services required to settle a cow and mature size. When one considers these estimates in concert with the experiences of the swine industry with pale, soft, and exudative pork (PSE), a definite red flag is raised.

Unfortunately, even though there have been numerous attempts to make one believe otherwise, these antagonisms leave no doubt that no one breed allows breeders to have their cake and eat it, too! Bourdon (1994) used the analogy of “sensible beef stew” to describe the effectiveness of utilizing designed mating systems to “mix and match” strengths and weaknesses of breeds to meet specifications for balanced performance. This fact has been further supported in the analysis of the American Gelbvieh Alliance results where a ratio of 50% British to 50% Continental European breeding appears optimal to hit market targets (Schiefelbein, 1998). Cundiff et al. (1994) additionally pointed out the need for alteration of breed inputs in sub-tropical environments to include either some *Bos indicus* or heat tolerant *Bos taurus* germ plasm.

Table 6. Weighted mean literature estimates of genetic correlations between various performance traits.^a

Traits ^b	Phenotypic Correlation	Genetic Correlation
Calving Ease/Birth Weight	-0.28	-0.74
Birth Wt/Feed Efficiency	-0.12	-0.46
Yearling Wt/Feed Efficiency	-0.46	-0.60
Feed Intake/Feed Efficiency	--	0.71
Wean Maternal/Feed Intake	--	0.80
Scrotal Circumference/Feed Efficiency	0.12	0.61
Birth Wt/Weaning Wt	0.46	0.50
Birth Wt/Yearling Wt	0.38	0.55
Weaning Wt/Yearling Wt	0.71	0.81
Weaning Wt/Mature Wt	0.45	0.57
Weaning Wt/Slaughter Wt	0.65	0.79
Yearling Wt/Slaughter Wt	0.65	0.56
Yearling Wt/Scrotal Circumference	0.36	0.39
Backfat/Feed Intake	0.29	0.44
Backfat/Scrotal Circumference	0.27	0.78
Carcass Wt/Birth Wt	0.41	0.60
Carcass Wt/Yearling Wt	0.85	0.91
Cutability/Yearling Wt	0.85	0.87
Marbling/Yearling Wt	0.14	-0.33
Marbling/Feed Intake	0.24	0.90
Marbling/Cutability	-0.25	0.35
Ribeye Area/Weaning Wt	0.23	0.49
Ribeye Area/Yearling Wt	0.35	0.51
Ribeye Area/Slaughter Wt	0.33	0.43
Ribeye Area/Cutability	0.33	0.45
Ribeye Area/Marbling	0.06	-0.21
Tenderness/Marbling	????	????
Tenderness/Cutability	????	????

^aEstimates shown are taken from Koots et al. (1994b) and represent the weighted mean of available literature estimates.

^bTraits represented are expressed on an age constant basis where appropriate and represent direct genetic effects.

Table 7. Genetic correlations between measures of carcass merit and reproductive efficiency (MacNeil et al., 1984).

Female Trait	Postweaning Gain (kg)	Carcass Weight (kg)	Fat Trim (kg)	Retail Product (kg)
Age at Puberty (d)	.16	.17	-.29	.30
Wt at Puberty (kg)	.07	.07	-.31	.08
Services/conception	1.33	.61	.21	.28
Gestation Length (d)	-.10	.03	-.07	.13
Calving Difficulty	-.60	-.31	-.31	-.02
Birth Weight (kg)	.34	.37	-.07	.30
Mature Weight (kg)	.07	.21	-.09	.25

Heterosis....The Final Piece of the Puzzle

Fortunately, nature has provided a significant amount of heterosis observed in the reproductive efficiency and maternal trait complex to allow breeders to overcome the obstacles of direct selection for fertility and cow adaptability mentioned earlier. Heterosis levels of 20 to 25% are achievable in pounds of calf weaned per cow exposed to breeding using systems which exploit a terminal sire breed mated to crossbred females of unrelated breeds (Table 8, Cundiff and Gregory, 1999). This amount varies according to the breeds used in the crossing system because heterosis is directly proportional to the difference in gene frequencies affecting the traits between the breeds used in the cross. This is the basis for the success of the *Bos indicus* x *Bos taurus* crosses in the sub-tropical zones where these females express phenomenal heterosis in maternal and reproductive performance.

Unfortunately, in the chase to utilize this “free-lunch” heterosis gift, as has too often been the case in animal breeding, there has been too much emphasis on “maximize” and not enough emphasis on “optimize”. When we recall what was mentioned before about evaluating the effects of change on cost per unit of output product value, there is an optimum amount of everything we do, even reproductive performance. Beyond that optimum it costs more to achieve than benefits received in return. This is an important concept to keep in check.

So, How Do We Genetically Manage to Simultaneously Improve End-product Performance and Lower Cost of Production?

Given that there are literally hundreds (thousands may be even more appropriate) of feed resource and climatic environments used in cattle production, yet end-product performance must fit within specification targets, what do we do? Animal breeders have unanimously stated over the past several generations of cattle production that we must achieve this balance by using breed complementarity and heterosis in very carefully designed crossbreeding programs. This must be a several step process to work successfully. First, the proper breeds must be chosen for matching maternal performance of the cow herd to a given production environment. Secondly, the proper

lines from within those breeds must be selected to properly hit those environmental targets while also meeting minimum acceptable performance in end-product characteristics. Then a terminal sire breed must be selected to bring necessary performance for growth and end-product performance to the system. Furthermore, the sires selected from within the terminal breed (or breeds) chosen, must have documented performance for growth and carcass traits (i.e. EPD) in addition to the sires selected for maternal replacements having documented EPD for reproductive and functional soundness.

Table 8. Heterosis effects in crosses of *Bos taurus* x *Bos taurus* breeds and in crosses of *Bos indicus* x *Bos taurus* breeds from diallel crossing experiments.^a

Trait	<i>Bos taurus</i> x <i>Bos taurus</i>			<i>Bos indicus</i> x <i>Bos taurus</i>		
	N	Units	%	N	Units	%
Crossbred calves (individual heterosis)						
Calving rate, %	11	3.2	4.4			
Survival to weaning, %	16	1.4	1.9			
Birth weight, kg	16	0.	2.4	4	3.3	11.1
Weaning weight, kg	16	7.4	3.9	10	21.7	12.6
Postweaning ADG, kg/d	19	.034	2.6	6	.116	16.2
Yearling weight, kg	27	13.2	3.8			
Cutability, %	24	-.3	-.6			
Quality grade, 1/3 grade	24	.12	--	6	.3	--
Crossbred cows (maternal heterosis)						
Calving rate, %	13	3.5	3.7	7	9.9	13.4
Survival to weaning	13	.8	1.5	7	4.7	5.1
Birth weight, kg	13	.07	1.8	6	1.9	5.8
Weaning weight, kg	13	8.2	3.9	12	31.1	16.0
Longevity, yrs	3	1.36	16.2			
Lifetime production						
No. calves	3	.97	17.0			
Cumulative weaning weight	3	272	25.3			

^aEstimates are from experiments contributing to North Central Regional Project NC-1 (Iowa, Indiana, Missouri, Ohio, USDA-ARS and Nebraska), Southern Regional Project S-10 (Virginia, Florida, Louisiana, Texas, USDA-ARS and Louisiana, USDA-ARS and Florida) as reported by Cundiff and Gregory (1999).

There are several different types of crossbreeding programs available to producers. These have been discussed in detail in the past (Bourdon, 1994; Kress, 1994; Cundiff and Gregory, 1999). There are certainly advantages and disadvantages to each of them. Unfortunately, a number of the product inconsistency problems our industry is experiencing today are from

misuse and abuse of these systems. It has not usually been the choice of the particular crossbreeding program that has gotten breeders into trouble as much as the inability to properly design, implement and then stay the course in a crossbreeding program. Many programs have been doomed from the start because they were not properly thought out, while yet others have failed because a new breed has come along that tempts the curiosity too much. Furthermore, there are still many breed and tradition loyalties which run rampant which often get in the way of breeding program objectivity. These facts, coupled with the wild chase for extra growth and extra heterosis have resulted in what some have called the “mongrelization” of the U.S. beef cow herd.

Is There Any Way to Reduce Crossbreeding Variation?

Cundiff and Gregory (1999) presented an excellent summary of the effectiveness of various crossbreeding systems in terms of heterosis utilization, use of breed complementarity, and consistency of production in 1994. In that presentation, the most effective system at doing all three things simultaneously, along with being the easiest to manage effectively, was composite breeding. The theory behind composites has been amply proven by the Germ Plasm Utilization Project at the U.S. Meat Animal Research Center under the leadership and guidance of Keith Gregory. The published summary (Gregory et al., 1995) of that work proves that composite breeding offers a usable solution to many of the problems we are discussing here. Heterosis utilization is high, breed percentages are fixed and do not vary between generations, and breed differences can be utilized to match breed strengths and weaknesses to the production and marketing environment. The ability to overcome genetic antagonisms and still retain high levels heterosis in maternal performance is unmatched by any of the other designed systems. Furthermore, once the composite is formed, the breeding system is much simpler to manage than any of the others.

Detractors of the composite approach have argued that composite mating systems will increase rather than decrease variability of production due to increased levels of heterozygosity. USDA-ARS work has shown that there is not a significant increase in the variability observed in the composite lines as compared to the purebreds (Table 9). Furthermore, compared to other mating systems such as rotational crosses and rota-terminal systems, the inter-generational variation is eliminated (Figure 2). These same detractors of composites have argued that we cannot afford to give up the consistency that purebreds have worked so hard to develop through their history. They do forget, however, that those purebreds with their consistency have to be the foundation for the composite lines. Just like there is no one breed that offers everything, the beef cattle industry will not be able to develop only one maternal line composite. While that may work better for the poultry and swine industries, it will not work for the beef industry. Therefore, the challenge is for the purebred breeds to find where they will fit into various composite lines as they develop.

Reactions to the idea of composite breeding have been very interesting to watch. There have been some purebred breeds which have realized that they need to find where they can provide useful germ plasm for the formation of these composites. There have been other breeds who have denounced the very idea of such an approach, rather than asking the question of where

they, too, can fit. There seems to be the mentality that the purebred breeds will disappear from existence because of composites. How can this be true? If one really analyzes the situation, the most likely scenario is for MATERNAL line composites to be developed by geo-climatic zone which will then be terminal sire mated to produce market offspring. Many of the breeders who are embracing and implementing the composite idea are doing so because they intend to produce replacement females and sire seedstock for the maternal side while at the same time selling purebred or F_1 terminal sires.

Table 9. Genetic standard deviations (S_g) and phenotypic coefficients of variation (CV) for purebreds and composites (castrate males).

Trait	Purebreds		Composites	
	S_g	CV	S_g	CV
200-d weight, kg	13.3	.10	14.2	.11
Slaughter weight, kg	21.7	.08	28.7	.08
Carcass weight, kg	12.4	.08	17.9	.09
12th rib fat, mm	1.3	.48	2.0	.44
Retail product, %	2.2	.04	2.3	.06
Carcass lean weight, kg	8.1	.08	10.7	.09
Carcass fat weight, kg	8.6	.18	6.3	.19
Carcass bone weight, kg	2.8	.08	2.1	.10
Longissimus muscle fat, %	.6	.27	1.0	.29
Shear force, kg	.18	.22	0.59	.21

(Cundiff and Gregory, 1999 and Gregory et al., 1995).

Figure 2. Variation resulting from various mating systems. (Cundiff and Gregory, 1994)

There are, however, also a few negatives to the composite approach. One is that in order to develop a composite line, it needs to be done from a relatively large base to avoid inbreeding. The typically used number is to have 400 to 500 females in the breeding population. Then there has to be intentional avoidance of inbreeding practiced to maintain the heterosis level in the composite line. A second negative is that genetic evaluation is much more difficult, i.e. EPD are not readily available, nor as accurate, for most composites as compared to purebreeds. We are just beginning to see some movement in the U.S. to address this problem through such efforts as the American Simmental Association's multi-breed EPD program (Pollak and Quaas, 1998). Another negative is that composite breeding still cannot overcome poor breeding decisions. A composite made from the wrong breeds and the wrong lines within those breeds still is a bad product. It has to be carefully and meticulously done.

How Are We Coming with Carcass EPD?

So, even if many problems can be remedied with designed breeding programs, breeders still must be able to accurately select the best animals. If we do not have the information for end-product breeding value, then how do we select the right terminal sire? For example, there is little doubt that the Brahman crossbred female is hard to outperform in the Gulf Coast region of the U.S.. However, we also know that we need to find a way to make sure that the Brahman sire lines used in that cross do not present the wrong type of end-product specs (particularly for tenderness; Sherbeck et al., 1995a,b; Crouse et al., 1989). If we do not have genetic predictions available for these carcass traits, we are doing no better than shooting in the dark.

If there is such a need for carcass EPD and the genetic bases of these traits is relatively high, why are they not widely available? Even though there have been several factors which have contributed to this problem, fortunately we are finally in the midst of seeing them resolved. The largest hindrance to collecting carcass information has been that until recently we have had to solely rely on progeny data. This type of information requires time, expense and labor to collect and also requires cooperation in the packing plant for accurate individual identification of carcasses. The combination of these factors has resulted in somewhat limited amounts of progeny data being placed into breed performance databases in the past. In the U.S., the American Angus Association has had the most concerted effort in designed progeny testing of sires. Approximately 50% of their currently published sires have carcass information (2,772 of 5,527 with published EPD (Angus, 1998)). While this proves the difficulty of obtaining progeny data for carcass traits, it also emphasizes that useful carcass information can be obtained for a meaningful percentage of the breed. Several other breed programs are attempting to build databases (Table 10).

The second hindrance has been the lack of ability to determine true carcass value differences on live, yearling seedstock cattle to circumvent the need for progeny data. Real-time ultrasound imaging technology has been pursued over the past ten years as the primary means to obtain these live animal measures and now appears to be entering the adoption mode. A national consortium of U.S. universities worked together during the early 1990s in a project which had as one of its three objectives "to determine the efficacy of using real-time ultrasound imaging to measure body composition and carcass merit traits in beef cattle" (Bertrand et al., 1994; Green et

al., 1994; Wilson et al., 1994). The conclusions drawn from a compilation of this and other research indicate: 1) assessment of retail yield amount or percentage on the basis of 12th rib fat thickness (FT) and 12th rib ribeye area (REA) is slightly less effective using ultrasonic measures on the live slaughter animal as compared to direct measures on the carcass postmortem (Hamlin et al., 1995; Herring et al., 1994; Perkins et al., 1992b); 2) FT is a better predictor of cutability than is REA in the current cattle population (Hamlin et al., 1995; Herring et al., 1994), although not so of retail product weight, 3) ultrasonic measures of these retail yield indicators appear to be under a moderate degree of genetic control (weighted average h^2 of .37 for FT and .26 for REA (Hassen et al., 1999; Shepard et al., 1996; Evans et al., 1995; Robinson et al., 1993; Johnson et al., 1993; Duello et al., 1993; Arnold et al., 1991; Turner et al., 1990; Lamb et al., 1990; deRose et al., 1988), 4) genetic correlation estimates between ultrasonic predictors of carcass merit and other economically important traits are sparse but indicate some antagonism between REA and mature size (Shepard et al., 1996; Johnson et al., 1993), 5) prediction of intramuscular fatness and palatability traits is more difficult using ultrasound, although high and acceptable levels of accuracy have been achieved in the past few years (Brethour, 1998; Crouch, personal communication; Wilson et al., 1995), and 6) data to estimate relationships between ultrasonic measures in yearling bulls and slaughter steer carcass retail yield and palatability have been more limiting (Crouch, personal communication; Kriese 1996; Dues et al., 1996a,b; Wilson et al., 1995; Evans et al., 1995; Steinkamp, 1995; Schalles et al., 1992).

Table 10. Current U.S. national cattle evaluation programs – carcass merit (as of 1 Jan 99).

Breed	Total Sires	Total Published Sires	Sires with Carcass Data	Sires with Carcass EBV	Traits Evaluated ^a	Ultrasound Accepted?
Angus	95,995	5,527	1,944	2,772	1,2,3,4,5	Yes
Beefmaster	10,756	401	180	Not released	2,3,4	Yes
Brangus	8,999	982	150	Not released	2,3,4	Yes
Charolais	21,453	1,650	27	0	2,3,4,5	Yes
Gelbvieh	5,173	1,800	363	219	1,2,3,4	Evaluating
Hereford	94,221	4,261	4,986	1,010	2,3,4	Yes
Maine-Anjou	1,240	348	55	0	1,2,3,4	Evaluating
Red Angus	16,910	1,145	829	293	2,3,4	No
Salers	10,827	657	N/A	85	1,2,3,4	No
Shorthorn	11,788	862	565	115	1,2,3,4	No
Simmental	80,804	2,804		372	1,4,5	Yes

^aTraits: 1=Carcass Weight, 2=Ribeye Area, 3=Fat Thickness, 4=Marbling, 5=%Retail Cuts.

This last issue has been the hardest one to resolve in recommending the adoption of ultrasound-generated carcass data for breed improvement programs. As data addressing this issue have been accumulated over the past five years, the conclusions of various researchers have not all agreed. However, as more data have been analyzed in some larger breed databases, the conclusions have become more clearly in favor of the use of real-time ultrasound. Data from the Brangus (Kriese, 1996) and Angus (Crouch, personal communication) breeds have indicated high

correlations between ultrasound and actual progeny carcass data for sires where both types of information have been collected. These conclusions have led to the recent adoption of policy to accept ultrasound data by several breed associations, including Angus, Hereford, Simmental, Brangus, and Gelbvieh (Table 10), with more associations to follow suit in the next few years. Coupled with actual carcass progeny data, use of real-time ultrasound data should allow great acceleration to occur in the percentage of active sires with carcass EPD for most breeds. For example, the American Angus Association amassed enough ultrasound data in the first 9 months after adoption to increase the size of its carcass record database by almost 50% (Crouch, personal communication).

Do We Have All of the Necessary Information to Genetically Address Carcass Acceptability?

The only area that may be a little tough (no pun intended) is genetic evaluation of overall meat quality, particularly tenderness. The reason that this is a major issue for the beef industry to confront is that we have estimates that one in five of the steaks produced in the current industry are tougher than desired (Morgan et al., 1991). No industry can afford this kind of defect rate!

There have been numerous debates in the U.S. over the last few years regarding how marbling can, or cannot, be used to address the meat quality and tenderness issue. This same discussion has also been occurring in Australia as they have begun to implement eating quality assurance grading standards (George, 1999). The collective U.S. experience indicates that while it would be nice to rely on marbling and USDA Quality Grade to be the “insurance policy” for palatability, it is simply not good enough. As shown graphically in Figure 3, while the probability of getting an unpalatable steak does significantly reduce when going from Standard up through Choice and Prime grades, there is so much overlap in palatability amongst the grades that today it is possible to have steaks from carcasses of Prime and Standard grades that will be equally palatable (Smith et al., 1987).

Figure 3. Relationship between palatability and USDA quality grade. (Smith et al., 1987)

Increasing emphasis is being placed on marbling in breeding programs at the current time, largely due to the marketing success of the Certified Angus Beef program (Marston et al., 1999). Increasing selection intensity for marbling appears to be short-sighted, however, for several reasons: 1) marbling is, at best, an insurance policy for eating satisfaction of beef; 2) marbling only explains 10 to 15% of the variation in overall palatability of cooked beef product; and 3) just as in any other trait, there are genetic antagonisms with marbling which must be carefully managed (recall the discussion of cutability and marbling earlier in this paper). However, without a more direct, accurate system for assessing true palatability differences, breeders are responding to increased consumer demand for quality and consistency using marbling as their selection criterion since it is the only tool available to them. As long as this selection occurs in a balanced, multiple trait configuration it should net small, yet positive, gains over time (Marston et al., 1999).

What About Tenderness?

It seems like there has been more discussion about beef tenderness in the past five years than in all of the previous century. As reviewed by Tatum (1999), beef is perceived to currently have a toughness problem, particularly in relation to cattle of *Bos indicus* descent (O'Connor et al., 1998; Sherbeck et al., 1995). There are two ways to handle this problem; tenderize the product post-mortem and/or genetically fix it. We know that postmortem aging, electrical stimulation, and calcium chloride injection post-mortem can be used to reduce toughness problems (Tatum et al., 1997). We also know that tenderness, assessed as Warner-Bratzler shear force of loin or rib steaks at a 14 d aging endpoint is heritable ($h^2 = .38$) and variable (Wulf et al., 1996). Although calpastatin, a primary inhibitor of muscle proteolysis post-mortem, appeared to offer a useful selection criterion in early research targeting genetic improvement in tenderness (Wulf et al., 1996; Koohmaraie et al., 1995), genetic polymorphisms in the calpastatin gene have not proved practically usable (Lonergan et al., 1995; Green et al., 1996a,b; Green et al., 1994). However, as pointed out in the previous presentation in this symposium, application of best management practices post-mortem results in a toughness rate that is still unacceptable, leaving the only remedy long-term through genetic selection (Tatum, 1999). Collectively, this means that breeders must position themselves on the tenderness issue by collecting objective progeny tenderness data (measured as Warner-Bratzler shear force).

This challenge has been taken very seriously by commodity group leaders in the U.S. and abroad. In 1998, an extensive genetic evaluation project for carcass merit was approved for funding by a consortium of 16 beef cattle breed associations and the beef checkoff (Green et al., 1998). The 42 month study, referred to as the National Carcass Merit Project, was initiated in June of 1998 and will collect complete carcass data (including ribeye shear force) from 11,000 progeny of sires from these breeds. Each breed is testing a minimum of ten of their most widely used sires with a minimum of 50 progeny each. Additional sires will be tested with fewer progeny in a majority of the breeds. The objectives of the project include estimation of EPD for shear force and sensory panel assessments of tenderness (as well as all other carcass traits) and an economic analysis of the costs and benefits associated with this type of information. As the planning for this project developed, a particular focal point of the effort became an evaluation of a set of previously identified and promising 11 DNA marker tests for carcass merit. Since DNA

technology has been portrayed at times to be the savior of the beef industry, more discussion of this area is warranted.

DNA to the Rescue

A number of developments over the past ten years have led to some DNA testing beginning to be made available to industry. Since we have been discussing tenderness and carcass merit traits at length, it might be helpful to show what is happening in this area as an example.

As pointed out previously, the beef cattle industry in the United States has been attempting to improve consumer demand for beef products by improving carcass merit of the cattle population. In particular, the need has been identified to increase the marbling potential of domestic grain-fed U.S. beef. In response to these needs, a beef checkoff funded project was initiated at the Angleton research station of Texas A&M University to identify genes, referred to as “*quantitative trait loci*” (QTLs), affecting variation in marbling ability (Taylor et al., 1996). The project was started in 1990 and required the development of resource families that would be expected to be highly heterozygous for gene loci affecting this trait as well as other measures of carcass merit. Based upon previous research, the scientists chose a design that utilized Brahman x Angus crosses to develop these families due to their divergent performance relative to marbling (Angus high, Brahman low). They first produced reciprocal F_1 crosses between these two breeds. These first crosses were then used to produce full sib families of backcrosses to either Angus or Brahman through multiple ovulation and embryo transfer. A total of 42 full sib families were produced representing 16 sires and 19 dams. Life history data on 613 head of progeny were collected in this project (Taylor et al., 1997).

In analyses conducted in this project, the research team identified a number of possible QTL for several traits including five genes which appear to affect marbling, and an additional seven genes that influence either tenderness as assessed by Warner-Bratzler shear force or sensory taste panel. Additionally, the project has allowed identification of five QTL effects on ribeye area and 5 QTL for dressing percentage. One QTL effect that was detected in the project seemed to influence postweaning growth independent of birth weight variation, a very favorable gene effect. This QTL maps to the same chromosome (bovine chromosome 2) that had been identified to contain the gene causing double-muscling, the so-called myostatin gene (Grobet et al., 1997).

Several things are evident from the experience and results of this project. First, it is clear that these resource families take a great deal of time to develop and collect information from. It is a slow, expensive, and laborious process. Secondly, the reverse genetics approach (i.e. designing the families to allow detection of differences after the fact), is fairly powerful for detecting these effects, but will likely only yield linked markers in chromosomal regions containing large QTL effects. The researchers still must positionally clone and sequence these gene loci before they can determine the genetic cause of these differences and have more direct genotype tests. This positional cloning requires much finer mapping in order to elucidate the gene of interest. Thirdly, because of patenting/licensing of any DNA tests that develop from this

type of research, much vagueness is observed in reporting of research results. Instead of knowing the map locations of the QTL effects presented above, the research group can only say they have identified effects rather than elucidating where those are located and in what linkage groups. Unfortunately this slows down overall progress in the field but is a fact of life in any form of current day biotechnology. Fourthly, because the reverse genetics approach hopes to identify markers to be used in a marker-assisted selection approach, the linkage relationships identified from a particular set of families may not hold up in other populations due to the phase of the linkage relationship. In other words, the markers linked to QTL effects identified in this particular project may not be useful in other families or breed populations. For example, perhaps the effects being found are breed-specific alleles that we already see in measuring differences between breeds, yet are not segregating within those other breeds (i.e. they are fixed). This last issue can possibly result in the direct application of QTL detected through this approach being difficult to apply beyond the resource population of study.

This research provides an excellent example of the process the animal industries will face to make usable technology from this approach. Given that it is unknown how useful the markers identified in that project will be across other families and breeds, a second step must be taken. This is where the National Carcass Merit Project mentioned earlier will play a major role. The project has several objectives, one being to validate the DNA markers identified by the Texas A&M project across the major U.S. beef breeds. The project is designed to collect complete carcass data, including Warner-Bratzler shear force, on 50 progeny from each of 10 widely used reference sires in each of the 16 breeds. Additionally, sensory panel evaluation will be performed on steaks from approximately 3,000 of these progeny. By going through this effort, the question will quickly be answered about whether these markers will be useful in a wide array of germ plasm. Additionally, it is hoped that the researchers will at the same time be able to move closer to positionally cloning the actual QTLs being “marked”. The breeds and numbers of progeny participating in the project are shown in Table 11.

It is interesting to note that there seems to be some redundancy in efforts occurring around the world in this area of technology development. One could easily take the Texas A&M and U.S. national project experience described above and change the names to CRC for Meat Quality and CSIRO Australia and have a very parallel story to tell over the same time period. The same could be done for the University of Saskatchewan and Ag. Canada. In the opinion of the author, more cooperation and less competitiveness in this area might do all of our industries a lot of good. Unfortunately, that does not appear to be the lay of the land for the future.

What Other QTL Have Been Found??

There are a number of QTL effects that have now been identified through research work at several locations. In addition to the work described above, associations have been reported for myostatin (Georges et al., 1998); growth traits (Beever et al., 1992); and carcass attributes, including tenderness (Keele et al., 1999; Stone et al., 1999; Green et al., 1996a,b). Additionally, associations have been reported with the ryanodine receptor gene with the pale, soft and exudative meat quality problem (Milan et al., 1996), markers associated with growth and fatness traits (Andersson et al., 1994; Archibald et al., 1994), and the estrogen receptor gene with litter

size (Rothschild et al., 1998) in swine; the callipyge gene with double-muscling in sheep (Cockett et al., 1994, 1997; Freking et al., 1999); the bovine leukocyte adhesion deficiency condition in Holstein dairy cattle and other markers related to milk production traits (Dentine, 1995; Georges et al., 1995); and the hyper-parakalemic periodic paralysis condition in American Quarter Horses (Spier et al., 1993). It is not a coincidence that many of these associations are with single-gene, simply inherited traits. We are likely to see much of the benefit of DNA marker, or direct gene, testing on these types of qualitative traits.

Table 11. Distribution of progeny across breeds in the National Carcass Genetic Merit Project (Green et al., 1998).

Breed	# DNA Sires @ 50 hd each	# Addl. EBV Sires @ 25 hd each	Total # Sires
Angus	10	20	30
Beefmaster	10	5	15
Brahman	10	5	15
Brangus	10	0	10
Braunvieh	10	0	10
Charolais	10	9	19
Gelbvieh	10	7	17
Hereford	10	23	33
Limousin	10	15	25
Maine-Anjou	10	5	15
Red Angus	10	10	20
Salers	10	0	10
Shorthorn	10	5	15
Simmental	10	15	25
Simbrah	10	5	15
South Devon	10	0	10
Total # Sires	160	124	284
Total # Progeny	8,000	3,100	11,100

EBV sires are to calculate EBV only (no DNA analyses will be performed).

More recently, mapping efforts have been initiated using complex study populations to identify large QTL effects for traits previously untouchable in genetic improvement programs. The two most exciting of these are both located in Nebraska. The first is a project being led by Daniel Pomp and Merlyn Nielsen at the University of Nebraska where they are using lines of mice which have been selected divergently for heat production. Heat loss can be used to estimate maintenance energy requirements of an animal through direct calorimetry. The Nebraska project was initiated in the early 1980s to determine if genetic variation existed for maintenance requirements using this approach. They have been successful in changing the heat production between high and low lines by 50% of the average. Earlier this year, this group reported that in a

QTL search of an F₂ intercross of lines of these mice, they were able to identify two major QTL affecting heat production, with another two putative QTL (Moody et al, 1998). This is exciting in that it indicates that it may be possible through marker-assisted selection approaches to identify animals with improved feed efficiency, perhaps our most difficult economic trait to measure.

A second major QTL effort is underway at the USDA-ARS Meat Animal Research Center. In one aspect of that effort, researchers are attempting to utilize the twinning population where selection has been applied over several generations for twinning rate, to detect QTL for ovulation rate and embryo survival. Initial results in that project have been very promising, with at least one major QTL already identified in the early part of the project (Kappes and Cundiff, personal communication).

Where Will Marker-assisted Selection Be of Greatest Benefit?

The resulting QTL that are identified through the ongoing searches of the developing bovine gene map are likely to be most beneficial for those traits that are difficult and expensive to measure, as detailed above. We can expect the following categories of traits to benefit the most from marker-assisted selection (in order of greatest to least degree of benefit): disease resistance and immunocompetence, carcass quality and palatability attributes, fertility and reproductive efficiency, maintenance requirements (i.e. energetic efficiency), carcass quantity and yield, milk production and maternal ability, and growth performance. This ranking is due to a combination of considerations including: 1) the relative difficulty in collecting performance data, 2) the relative magnitude of the heritability and phenotypic variation observed in the traits, 3) the current existing amount of performance information available, and 4) when performance data becomes available in the life-cycle of the cow herd, (collected at birth, weaning, yearling, maturity?). Most of the rankings above then become self-explanatory.

To be realistic, however, we must realize that QTL will not serve as magic silver bullets. As long as we are relying on markers, rather than the QTL themselves, we are still only crudely defining the genotypes across the larger beef cattle population. Once QTL are finely mapped and direct tests are available, then the accuracy of selection will be markedly improved. Until then, however, the marginal gains that MAS will give us over selection on polygenic breeding values is not as high as one might think. Charles Smith predicted in the late 1970s that gene level information would only provide substantial gains for traits where information is lacking for genetic evaluation. Thus, for traits that are currently widely evaluated in national cattle evaluation programs (eg. growth rate) MAS will not help a great deal. His prediction was quite accurate given that results from a number of simulation studies now show quite clearly that he was correct (Haley and Visscher, 1998). The overall conclusion is that markers combined with EPD information will enhance the accuracy of genetic evaluation. The marginal gain, however, is very dependent on the particular trait, the number of markers available for whole genome scans, the availability of performance information, and the availability of marker information from large numbers of animals.

Lastly, in the current day climate of the beef cattle industry, where everyone is looking for

a quick and magical solution to all that ails us, it is imperative that breeders not overestimate this technology. It is not uncommon to hear statements such as “all we have to do is find the marbling gene and then we will have the problem solved”. This is absolutely not going to happen. We must remember that economically important quantitative traits are controlled by many genes, i.e. they are polygenic. While it is also likely true that there are some of those genes that play a bigger role than others, there still are many genes in the picture. Put simply, to put a measure of performance in front of the word gene and then say that is all we need is much too simplistic.

While marker-assisted selection is a popular new phrase in academic animal breeding circles, what we are more likely to see needed is what could be called “marker-assisted optimum selection”. What this means is that markers identifying QTL of large effect can be used to add to EPD for the same trait. This will result in optimal use of information from both the molecular and phenotypic performance levels. In other words, marker-assisted selection or EPD singularly are not great, but together they markedly increase the accuracy of genetic evaluation. It is important for the breeder to put these last two paragraphs firmly into perspective.

But How Does One Practice Balanced Trait Selection When There Are So Many Important Traits?....

One of the areas being currently debated by some of the thought leaders in the academic beef cattle breeding community would really help producers in this regard. The ideas being batted around relate to how to best combine information on several traits into “selection indexes” for specified breeding objectives. These ideas have been around a long time (since Jay Lush and Lanoy Hazel at Iowa State first proposed them in 1943), but have really become applicable and important as we have developed genetic information on more and more traits in the past ten years. For example, suppose one is looking for a maternal-line bull to produce females for a given production and marketing environment, then these indexes of traits weighted according to their relative amounts of heritable variation, relationships with other traits, and relative economic importance could be very valuable tools.

The dairy and swine industries have already produced indexes for use in their national genetic evaluation programs and I will not be surprised if the beef industry sees rapid development of the same in the next few years. In our case, however, the indexes will need to be somewhat “customized” for a given type of production scenario, similar to the discussion earlier in this paper about the importance of defining and sticking to a particular market target for a producer before deciding what to do genetically. Fortunately, we are seeing tools developed to help in this area such as the recently released Decision Evaluator for the Cattle Industry (DECI) model developed by USDA scientists at the U.S. Meat Animal Research Center (Bourdon, 1998; Jenkins and Williams, 1999). This simulation model is an attempt to provide a tool that will allow a producer to provide a base-line picture of his/her production system so that lots of “what if” questions can be asked. This is a critical area where a great deal of research and development is needed.

Implications

The following are unavoidable conclusions from experiences over the past 30 years:

- 1) Beef is losing market share relative to poultry and pork.
- 2) A large portion of the reason for lost market share is due to higher costs of production. Reproductive efficiency and other aspects of maternal performance in the environment cannot be sacrificed.
- 3) We can genetically alter cattle for end-product performance.
- 4) The most feasible way to approach the end-product non-conformance problem genetically is to use properly designed and implemented crossing systems which match maternal production to environmental feed resources with sire selection based on growth and carcass performance.
- 5) The benefits of heterosis on overall performance of a cow herd (upwards of 25% improvement in weight of calf weaned per cow exposed) cannot be ignored.
- 6) Proper terminal sire selection for growth and carcass performance is unachievable in the absence of end-product EPD and proper maternal sire selection for reproductive and maternal performance is unachievable in the absence of appropriate EPD.
- 7) Single-trait selection has never been and will never be a wise breeding philosophy.

Given these conclusions, we have several possible approaches to be successful in *achieving both cow adaptability and carcass acceptability*. Each of these has merit, and therefore, should be attempted. In priority order, they are:

- 1) Immediately demand that end-product performance data be gathered and utilized in national cattle evaluation programs. This must be done by amassing the necessary progeny data (either carcass or ultrasound) for lean yield and objectively measured meat quality attributes. Additionally, we must implement whole-herd reporting formats for breed performance data collection to enable calculation of EPD for fertility and longevity-related traits.
- 2) Educational plans should be developed, by geo-climatic region, for matching of breed resources to environments.
- 3) Breeders must more willingly evaluate the alternative of using “composite” breeding programs where they are applicable. Breeds need to be working today to determine where their germ plasm fits into the composite puzzle which is inevitably going to become a reality.
- 4) We must develop a high-integrity system of identification on every animal produced in the beef production system. We must then be willing to use this system to provide information feedback and true value discovery/pricing.
- 5) We must use all available resources to identify new DNA-based technologies to assist in making genetic improvement in problematic traits (eg. Maintenance energy cost, disease resistance).
- 6) Seedstock suppliers must adopt the philosophy of being a FULL SERVICE GENETIC PROVIDER to their clientele.

Several of these items are much more politically difficult to achieve than others which are physically more challenging. The political may, or may not, ever happen. This entirely depends

on whether groups and people within the industry are committed to the good of the whole industry or the good of their portion of the industry. The physical challenges, however, are ones over which breeders and scientists have direct control. We can make those happen with the right resources directed in the right directions (eg. carcass EPD, individual animal ID and feedback, and whole-herd reporting). What happens if every commercial bull-buyer starts demanding reproductive/fertility and end-product EPD specifications before he/she will buy a bull? What happens if a feedlotter requires an electronic ID (or better yet a DNA bar code) on every incoming feeder or yearling before they will purchase and/or feed them? What happens if, on the basis of that identification, the industry provides direct pricing (with information feedback) on every animal? What happens if seedstock suppliers develop systems to analyze the needs of their customers followed by production of specification seedstock rather than producing first and then trying to find customers? What happens if animal breeding scientists take a more active role in education and technology adoption? The beef industry would universally benefit in the long term.

Literature Cited

- Allen, D. L., L. Corah, and H. Gardiner. 1987. Evaluating Genetic Influences on Carcass Traits. Proceedings of Annual Mtg. of Beef Improvement Federation, Wichita, KS, pp 55-63.
- American Angus Association. 1998. Spring Sire Evaluation Report, April 1998.
- Andersson, L., C. S. Haley, H. Ellegren, S.A. Knott, M. Johansson, K. Andersson, L. Andersson-Eklund, I. Edfors-Lilja, M. Fredholm, I. Hansson, J. Hakansson, and K. Lundstrom. 1994. Genetic mapping of quantitative trait loci for growth and fatness in the pig. *Science* 263:1771.
- Bourdon, R. M. 1994. Breed roles in hitting a new target. Proceedings of Annual Mtg. of Beef Improvement Federation, Des Moines, IA, pp 115-122.
- Bourdon, R.M. 1998. Shortcomings of current genetic evaluation systems. *J. Anim. Sci.* 76:2308.
- Bourdon, R.M. 1988. Bovine nirvana-from the perspective of a modeler and purebred breeder. *J. Anim. Sci.* 66:1892.
- Brethour J. 1998. Ultrasound backfat and marbling projections using serial ultrasound measures to generate models of marbling and backfat thickness changes in feedlot cattle. *J. Anim. Sci.* (in review).
- Brinks, J.S. 1996.
- Cockett, N.E., S.P. Jackson, T.L. Shay, D. Nielsen, S.S. Moore, M.R. Steele, W. Barendse, R.D. Green, and M. Georges. 1994. Chromosomal localization of the callipyge gene in sheep (*Ovis aries*) using bovine DNA markers. *Proc. National Academy of Science* 91:3019.
- Crouse, J.D., L.V. Cundiff, R.M. Koch, M. Koohmaraie, and S.C. Seideman. 1989. Comparisons of *Bos indicus* and *Bos taurus* inheritance for carcass characteristics and meat palatability. *J. Anim. Sci.* 67:2661.
- Cundiff, L.V. and K.E. Gregory. 1999. What is systematic crossbreeding? Proceedings of Cattlemens College, National Cattlemens Beef Association, Charlotte, NC. February 1999.
- Cundiff, L.V., K.E. Gregory, and R.M. Koch. 1994. Breeding cattle for improved product consistency. In: Proceedings of National Cattlemen's Association Cattlemen's College.,

- Reno, NV, January 1994.
- de Rose, E.P., J.W. Wilton, and L.R. Schaeffer. 1988. Estimation of variance components for traits measured on station-tested beef bulls. *J. Anim. Sci.* 66:626.
- Dickerson, C.E. 1970. Efficiency of animal production – molding the biological components. *J. Anim. Sci.* 30:849.
- Diles, J.J.B., R.D. Green, H.H. Shepard, L.J. Hughes, and G.L. Mathiews. 1996a. Variation within and between groups of genetically identical nuclear transfer Brangus bulls and steers for growth measures over time. *Prof. Anim. Scientist* 12:250.
- Diles, J.J.B., R.D. Green, H.H. Shepard, L.J. Hughes, G.L. Mathiews, and M.F. Miller. 1996h. Relationships between body measurements obtained on yearling Brangus bulls and measures of carcass merit obtained from their steer, clone-mates. *Prof. Anim. Scientist* 12:244.
- Doyle, S.P., R.D. Green, B.L. Golden, C.L. Mathiews, C.R. Comstock, and D.G. LeFever. 1996. Genetic parameter estimates for heifer pregnancy and subsequent rebreeding rate in Angus cattle. *J. Anim. Sci.* 74 (Suppl. I): 117.
- Duello, D.A., D.E. Wilson, and C.H. Rouse. 1993a. The relationship between real-time ultrasound and carcass measurements of 12-13th rib fat thickness and longissimus muscle area in beef cattle. *J. Anim. Sci.* (submitted).
- Duello, D.A., D.E. Wilson, and G.H. Rouse. 1993b. Adjustment procedures and genetic parameter estimates of serial growth and ultrasonically measured carcass traits in performance-tested yearling bulls. *J. Anim. Sci.* (submitted).
- Evans, J.L., B.L. Golden, C.R. Comstock, K.L. Long, R.M. Bourdon, C.H. Mallinckrodt, and R. D. Green. 1996. Genetic parameter estimates for heifer pregnancy rate in Hereford cattle. *J. Anim. Sci.* 74 (Suppl. I):117.
- Evans, J.L., B.L. Golden, D.R.C. Bailey, R.P. Gilbert, and R.D. Green. 1995. Genetic parameter estimates of ultrasound measures of backfat thickness, loin eye muscle area, and gray shading score in Red Angus cattle. *Proc. Western Sect. Amer. Soc. Anim. Sci.* 46:202.
- Freking, B.A., J.W. Keele, S.D. Shackelford, T.L. Wheeler, M. Koohmaraie, M.K. Nielsen, and K.A. Leymaster. 1999. Evaluation of the ovine callipyge locus: III. Genotypic effects on meat quality. *J. Anim. Sci.* (in press).
- George, M.L. 1999. The role of MSA in meeting customer requirements. *Proceedings of Association for the Advancement of Animal Breeding and Genetics, Mandurah, Western Australia.* July 1999.
- Georges, M., L. Grobet, D. Poncelet, L.J. Royo, D. Pirottin, and B. Brouwers. 1998. Positional candidate cloning of the bovine mh locus identifies an allelic series of mutations disrupting the myostatin function and causing double-muscling in cattle. *Proceedings of 6th World Congr. Applied to Livestock Prod.* 26:195.
- Golden, B.L., W.M. Snelling, J.L. Evans, and C.R. Comstock. 1996. Reproduction: The Next Era of Evaluation. *Proceedings of Annual Mtg. of Beef Improvement Federation, Birmingham, AL*, pp 48-56.
- Green, R.D. 1999. DNA + EPDs = Marker-Assisted Optimum Selection. *Proceedings of Cattlemens College, National Cattlemens Beef Association, Charlotte, NC.* February 1999.
- Green, R.D. 1999. Breeding to satisfy customer requirements in the 21st century. *Proceedings of Association for the Advancement of Animal Breeding and Genetics, Mandurah,*

- Western Australia. July 1999.
- Green, R.D. 1999. Utilizing new genetic improvement technologies to improve beef profitability and palatability. Proceedings of Association for the Advancement of Animal Breeding and Genetics. July 1999.
- Green, R.D., M.E. Dikeman, J. Pollak, S. Koontz, J.F. Taylor, S.K. Davis, and J.O. Reagan. 1998. Overview of the National Tenderness EPD and DNA Marker Validation Project. Proceedings of Ann. Mtg. of Beef Improvement Federation, Calgary, ALB., July 1998.
- Green, R.D., N.E. Cockett, J.D. Tatum, D.M. Wulf, D.L. Hancock, and G.C. Smith. 1996a. Association of a TaqI calpastatin polymorphism with postmortem measures of beef tenderness in Charolais- and Limousin-sired steers and heifers. *J. Anim. Sci.* 74 (Suppl. I):113.
- Green, R.D., N.E. Cockett, J.D. Tatum, S.F. O'Connor, D.L. Hancock, and G.C. Smith. 1996b. Association of a TaqI calpastatin polymorphism with postmortem measures of beef tenderness in *Bos taurus* and *Bos taurus*-*Bos indicus* steers and heifers. *J. Anim. Sci.* 74 (Suppl. I):111.
- Green, R.D. 1995. Why do carcass EPD not already exist? Proceedings of 5th Beef Improvement Federation Genetic Prediction Workshop, Kansas City, MO, December 1995. pp69-75.
- Green, R.D., N.E. Cockett, M.F. Miller, D.L. Hancock, C. Bidwell, L.S. Barrett, J.B. Morgan, and J.D. Tatum. 1994. Characterization of TaqI polymorphisms in the bovine calpastatin gene. Proceedings 5th World Congress on Genetics as Applied to Livestock Production 19:450.
- Gregory, K.E., L.V. Cundiff, and R.M. Koch. 1995. Genotypic and phenotypic (co)variances for growth and carcass traits of purebred and composite populations of beef cattle. *J. Anim. Sci.* 73:1920.
- Haley, C.S. and P.M. Visscher. 1998. Strategies to utilize marker-quantitative trait loci associations. *J. Dairy Sci.* 81:85.
- Hamlin, K.E., R.D. Green, L.V. Cundiff, T.L. Wheeler, and M.E. Dikeman. 1995. Real-time ultrasonic measurement of fat thickness and Longissimus muscle area in diverse biological types of beef, steers: II. Relationship between real-time ultrasound measures and carcass retail yield. *J. Anim. Sci.* 73:1725.
- Hassen, A., D.E. Wilson, and G.H. Rouse. 1999. Evaluation of carcass, live, and real-time ultrasound measures in feedlot cattle. I. Assessment of sex and breed effects. *J. Anim. Sci.* 77:273.
- Hoehnbooken, W.D. 1988. Bovine nirvana-from the perspective of an experimentalist. *J. Anim. Sci.* 66:1885.
- Herring, W.O., S.E. Williams, J.K. Bertrand, L.L. Benyshek, and D.C. Miller. 1994. Comparison of live and carcass equations predicting percentage of cutability, retail product weight and trimmable fat in beef cattle. *J. Anim. Sci.* 72:1107.
- Jenkins, T.G. and C.B. Williams. 1999. An overview of the Decision Evaluator for the Cattle Industry model. Proceedings of Cattlemens College, National Cattlemens Beef Association, Charlotte, NC. February 1999.
- Johnson, M.Z., R.R. Schalles, M.E. Dikeman, and B.L. Golden. 1993. Genetic parameter estimates of ultrasound-measured Longissimus muscle area and twelfth-rib fat thickness in Brangus cattle. *J. Anim. Sci.* 71:2623.

- Kappes, S.M., J.W. Keele, R.T. Stone, R. McGraw, T.S. Sonstegard, T.P.L. Smith, N. Lopez-Corrales. 1997. A second generation linkage map of the bovine genome. *Genome Res.* 7:235.
- Keele, J.W., S.D. Shackelford, S.M. Kappes, M. Koohmaraie, and R.T. Stone. 1999. A region on bovine chromosome 15 influences beef longissimus tenderness in steers. *J. Anim. Sci.* 77:1364.
- Koohmaraie, M., J. Killefer, M.D. Bishop, S.D. Shackelford, T.L. Wheeler, and J.R. Arbona. 1995. Calpastatin based methods for predicting meat tenderness. In: A. Ouli, D. Demeyer, and F. Smulders (Ed.). "Expression, Regulation and Role of Proteinases in Muscle Development and Meat Quality". pp 395-412. ECCEAMST (European Consortium for Continuing Education in Advanced Meat Science and Technology), Utrecht, The Netherlands.
- Koots, K.P., J.P. Gibson, C. Smith, and J.W. Wilton. 1994a. Analyses of published genetic parameter estimates for beef production traits. I. Heritability. *Anim. Breeding Abstr.* 62:309.
- Koots, K.P., J.P. Gibson, C. Smith, and J.W. Wilton. 1994b. Analyses of published genetic parameter estimates for beef production traits. 2. Phenotypic and genetic correlations. *Anim. Breeding Abstr.* 62:825.
- Kress, D.D. 1994. Crossbreeding with a New Target. *Proceedings of Annual Mtg. of Beef Improvement Federation*, Des Moines, IA, pp 83-92.
- Kriese, L.A. 1996. Ultrasound: Past pitfalls and present promise? *Proceedings of Annual Mtg. of Beef Improvement Federation*, Birmingham, AL, pp 73-87.
- Lamb, M.A., O.W. Robison, and M.W. Tess. 1990. Genetic parameters for carcass traits in Hereford bulls. *J. Anim. Sci.* 68:64.
- Loneragan, S.M., C.W. Ernst, M.D. Bishop, C.R. Calkins, and M. Koohmaraie. 1995. Relationship of restriction fragment length polymorphisms (RFLP) at the bovine calpastatin locus to calpastatin activity and meat tenderness. *J. Anim. Sci.* 73:3608.
- Macneil, M.D., L.V. Cundiff, C.A. Dinkel, and R.M. Koch. 1984. Genetic correlations among sex-limited traits in beef cattle. *J. Anim. Sci.* 58:1171.
- Marshall, D. 1994. Breed differences and genetic parameters for body composition traits in beef cattle. *J. Anim. Sci.* 72:2745.
- Marston, T.T., J.P. Gleghorn, and L.E. Wankel. 1999. The impact of selecting for marbling on beef cow herds. *Certified Angus Beef*, American Angus Association, St. Joseph, MO.
- Melton, B.E. 1995. Conception to consumption: The economics of genetic improvement. *Proceedings of Annual Mtg. Beef Improvement Federation*, Sheridan, WY. pp 40-87.
- Melton, B.E., E.O. Hady, and R.L. Willham. 1979. Estimation of economic values for selection indices. *An. Prod.* 28:279.
- Moody, D.E., D. Pomp, and M.K. Nielsen. 1998. Isolation of quantitative trait loci (QTL) controlling energy balance in mice. *Proc. of 6th World Congr. on Genetics Applied to Livestock Prod.* 26:300.
- Morgan, J.B., J.W. Savell, D.S. Hale, R.K. Miller, D.B. Griffin, H.R. Cross, and S.D. Shackelford. 1991. National beef tenderness survey. *J. Anim. Sci.* 69:3274.
- NCA. 1992. Improving the consistency and competitiveness of beef: Final report of the national beef quality audit -- 1991.
- O'Connor, S.F., J.D. Tatum, R.D. Green, D.M. Wulf, and G.C. Smith. 1997. Genetic influences

- on beef palatability in *Bos indicus* composite and *Bos taurus* cattle. *J. Anim. Sci.* 75:1822.
- Pollak, E.J. and R.L. Quaas. 1998. Multi-breed genetic evaluations of cattle. *Proc. of 6th World Congr. on Genet. Applied to Livestock Production.* 23:81.
- Robinson, D.L., K. Hammond, and C.A. McDonald. 1993. Live animal measurement of carcass traits: Estimation of genetic parameters for beef cattle. *J. Anim. Sci.* 71:1128.
- Rothschild, M.F. 1998. Identification of quantitative trait loci and interesting candidate genes in the pig: Progress and prospects. *Proceedings of 6th World Congr. on Genetics Applied to Livestock Prodn.* 26:403.
- Schalles, R.R. 1992. Ribeye area EPD's for Brangus cattle. In: *Proceedings of "Looking Under the Hide" Symposium, Brinks Brangus Field Day, October 1992.*
- Shackelford, S.D., M. Koohmaraie, L.V. Cundiff, K.E. Gregory, G.A. Rohrer, and J.W. Savell. 1994. Heritabilities and phenotypic and genetic correlations for bovine postrigor calpastatin activity, intramuscular fat content, Warner-Bratzler shear force, retail product yield and growth rate. *J. Anim. Sci.* 72:857.
- Schiefelbein, D. 1998. Back to the Basics – A Real World Strategy for Improving the Quality and Consistency of Beef. *Proceedings of the Annual Mtg. of Beef Improvement Federation, Calgary, AB, pp 74-89.*
- Shepard, H.H., R.D. Green, B.L. Golden, K.E. Hamlin, T.L. Perkins, and J.B. Diles. 1996. Genetic parameter estimates for live animal ultrasonic measures of carcass merit in Angus cattle. *J. Anim. Sci.* 74:761.
- Sherbeck, J.A., J.D. Tatum, T.G. Field, J.B. Morgan, and G.C. Smith. 1995a. Feedlot performance and palatability traits of Hereford and Hereford x Brahman steers. *J. Anim. Sci.* 73:3613.
- Smith, G.C., J.W. Savell, H.R. Cross, Z.L. Carpenter, C.E. Murphey, G.W. Davis, H.C. Abraham, F.C. Pamsh, and B.W. Berry. 1987. Relationship of USDA quality grades to palatability of cooked beef. *J. Food Quality* 10:269.
- Snelling, W.M., B.L. Golden, and R.M. Bourdon. 1995. Within-herd genetic analyses of stayability of beef females. *J. Anim. Sci.* 73:993.
- Steinkamp, K. 1995. Validation of real-time ultrasound measurements in live Angus cattle to predict body composition. *Proceedings of Annual Mtg. of Beef Improvement Federation, Sheridan, WY. pp 263-265.*
- Stone, R.T., J.W. Keele, S.D. Shackelford, S.M. Kappes, and M. Koohmaraie. 1999. A primary screen of the bovine genome for quantitative trait loci affecting carcass and growth traits. *J. Anim. Sci.* 77:1379.
- Tatum, J.D. 1999. New approaches for improvement in beef tenderness, quality, and consistency. *J. Anim. Sci.* (in press).
- Tatum, J.D., M.H. George, K.E. Belk, and G.C. Smith. 1997. Final Report: Development of a Palatability Assurance "Critical Control Points" (PACCP) Model to Reduce the Incidence of Beef Palatability Problems. NCBA, March 1997.
- Taylor, J.F. and S.K. Davis. 1997. Beef Cattle Genomics: Past, Present, and Future. *Conference Proceedings, Texas A&M University, October 15-17, 1997.*
- Taylor, J.L., S.K. Davis, J.O. Sanders, J.W. Turner, J.W. Savell, R.K. Miller, S.B. Smith, N.E. Cockett, A. Eggen, M.D. Bishop, and S.M. Kappes. The Angleton project: 1996 update. 1996. *Proceedings of the Annual Mtg. of the U. S. Beef Improvement Federation,*

- Birmingham, AL, pp 28-37.
- Taylor, R.E. and T.G. Field. 1999. Beef production and Management Decisions. 3rd Ed. Prentice Hall, Upper Saddle River, NJ.
- Turner, J.W., L.S. Pelton, and H.R. Cross. 1990. Using live animal ultrasound measures of ribeye area and fat thickness in yearling Hereford bulls. J. Anim. Sci. 68:3502.
- Wilson, D.E. 1995. Carcass and live animal evaluation. Proceeding of 5th BIF Genetic Prediction Workshop, Kansas City, MO, December 1995. pp 25-34.
- Wilson, D.E. 1992. Application of ultrasound for genetic improvement. J. Anim. Sci. 70:973.
- Wulf, D.M., J.D. Tatum, R.D. Green, J.B. Morgan, B.L. Golden, and G.C. Smith. 1996. Genetic influences on beef palatability in Charolais- and Limousin-sired steers and heifers. J. Anim. Sci. 74:2394.