

6-2005

Registration of N30-N56, N741, N743, N745, N747, U362, U363, U367, U369-U374, U389-U394, U396-U398, and U500 Sweetclover Genetic Stocks

Kenneth P. Vogel

University of Nebraska - Lincoln, Ken.Vogel@ars.usda.gov

H. J. Gorz

University of Nebraska - Lincoln

F. A. Haskins

University of Nebraska - Lincoln, fhaskins@neb.rr.com

Follow this and additional works at: <http://digitalcommons.unl.edu/agronomyfacpub>

 Part of the [Plant Sciences Commons](#)

Vogel, Kenneth P.; Gorz, H. J.; and Haskins, F. A., "Registration of N30-N56, N741, N743, N745, N747, U362, U363, U367, U369-U374, U389-U394, U396-U398, and U500 Sweetclover Genetic Stocks" (2005). *Agronomy & Horticulture -- Faculty Publications*. 151.

<http://digitalcommons.unl.edu/agronomyfacpub/151>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Registration of N30-N56, N741, N743, N745, N747, U362, U363, U367, U369-U374, U389-U394, U396-U398, and U500 Sweetclover Genetic Stocks

Forty-nine white-flowered sweetclover (*Melilotus alba* Medik.) genetic stocks [N30-N45 (Reg. GS-1-16, PI 549120-549135); N46-N53 (Reg. GS-17-24, PI 557503-PI 557510); N54-N55 (Reg. GS-25-Reg. GS-26, PI 629289-PI 629290); N741, N743, N745, N747 (Reg. GS-27-GS-30, PI 557511-PI 557514); U362, U363, U367 (Reg. GS-31, Reg. GS-32, Reg. GS-33, PI 557515-PI 557517); U369-U374 (Reg. GS-34-GS 39, PI 557518-PI 557523); U389-U394 (Reg. GS-40-GS 45, PI 557524-PI 557529); U396-U398 (Reg. GS-46-GS 48, PI 557530-PI 557532); U500 (Reg. GS-49, PI 557533)] (Table 1); and N56 (Reg. no. GS-50, PI 634019), a yellow-flowered sweetclover [*Melilotus officinalis* (L.) Lam.] genetic stock, were developed jointly by USDA-ARS and the Agricultural Research Division, Institute of Agriculture and Natural Resources, University of Nebraska-Lincoln and were jointly released in May 2004. The genetic stocks, which contain unique combinations of genes and traits, were developed over more than three decades of cooperative sweetclover genetic research.

The 49 *M. alba* genetic stocks include a set of 16 lines, N30 through N45, which represent all possible homozygous combinations of four allelic pairs, *Y/y*, *C/c*, *Cu/cu*, and *B/b* (Table 1). The *Y/y* alleles affect seed color, and the *C/c* alleles are concerned with both seed and seedling color (Gorz et al., 1975). The *Cu/cu* and *B/b* genes affect coumarin (more accurately, *o*-hydroxycinnamic acid β -D-glucoside) content (Gorz and Haskins, 1969) and β -glucosidase activity (Haskins and Gorz, 1965), respectively. The development of these 16 lines involved both annual and biennial forms of *M. alba*, and the greenhouse conditions under which seed of these lines were produced did not permit distinguishing between these forms. Both forms may be present in these lines.

Lines N46 through N49, and N741, N743, N745, and N747, are two sets of four lines, each set representing all possible homozygous combinations of the *Cu/cu* (coumarin content) and *B/b* (β -glucosidase activity) alleles. N46 through N49 are annuals. They were derived from an initial cross of *cucubb* biennial plants \times *CuCuBB* plants of PI 165554, a small, annual, autogamous introduction from India, followed by six successive backcrosses of *cucubb* segregates to the *CuCuBB* annual parent. N741, N743, N745, and N747 are biennial lines. They are F_{21} generation lines derived from an initial *cucubb* \times *CuCuBB* cross followed by self-pollination of a single doubly heterozygous plant in each generation from F_1 to F_{17} . The four homozygous genotypes were isolated in F_{18} .

N50 through N53 are biennial lines representing all possible homozygous combinations of the *Y/y* and *C/c* allelic pairs. As indicated above, the *C/c* genes influence seedling color, and both *Y/y* and *C/c* affect seed color. These four lines are the F_6 generation from a single F_1 plant that was obtained from a cross of the N1 strain (*yyCC* genotype) \times a line designated JF-1 (*YYcc* genotype). N54 and N55 are biennial lines that are homozygous for susceptibility and resistance, respectively, to stem canker (gooseneck) disease [caused by *Ascochyta caulicola* (Laub.)].

U389 is an annual line that was derived from a single plant of the introduction, PI 165554, mentioned above. All of the other U-numbered lines were developed following treatment of U389 seed with ethyl methanesulfonate. Although not always identified as such, U389 was the "wild-type" (+/+) line used in the referenced studies involving the lines that resulted from ethyl methanesulfonate treatment.

The normal parent (U389) and the chlorophyll-deficient mutants (U369, U371, U372, U373, U374, U396, U397, U398)

were used by Markwell and coworkers (Bevins et al., 1993, 1992; Markwell and Chelgen, 1988; Markwell et al., 1986, 1985a, 1985b; Yang et al., 1990) and Nakitani and Baliga (1985) in their biochemical research. U389 also was used by Kneen and LaRue (1988) to create a series of non-nodulating mutants for studying the process of nitrogen fixation in legumes.

N56, a biennial strain of *M. officinalis*, was developed by crossing N27 (a large-seeded, high-coumarin, early-maturing *M. officinalis*) to N1 (a finestem, small-seeded, low-coumarin, late-maturing *M. alba*) with one backcross of finestem, low-coumarin F_2 segregates to N27 followed by a second backcross to N29 (a low-coumarin strain of *M. officinalis*). N56 combines finestem growth habit and low coumarin content of *M. alba* with the large-seeded trait and early maturity of *M. officinalis*.

Registration of these genetic stocks supplements three previously released and registered biennial, yellow-flowered, sweetclover germplasms, N27, N28, and N29 (Gorz et al., 1992a, 1992b). Seed of all lines has been deposited in the National Plant Germplasm System. Requests for any of the 49 *M. alba* lines and N56 *M. officinalis* should be to the National Plant Germplasm System (<http://www.ars-grin.gov/npgs.orders.html>; verified 9 March 2005). Seed should be scarified before planting. It is requested that appropriate recognition be made if these genetic stocks contribute to research or the development of a new breeding line or cultivar.

K.P. VOGEL,* H.J. GORZ, AND F.A. HASKINS

References

- Bevins, M.A., S. Madhavan, and J. Markwell. 1993. Two sweetclover (*Melilotus alba* Desr.) mutants temperature sensitive for chlorophyll expression. *Plant Physiol.* 103:1123-1131.
- Bevins, M., C. Yang, and J. Markwell. 1992. Characterization of a chlorophyll-deficient mutant of sweetclover (*Melilotus alba*). *Plant Physiol. Biochem.* 30:327-331.
- Gengenbach, B.G., H.J. Gorz, and F.A. Haskins. 1970. Genetic studies of induced mutants in *Melilotus alba*. II. Inheritance and complementation of chlorophyll-deficient mutants. *Crop Sci.* 10:154-156.
- Gengenbach, B.G., F.A. Haskins, and H.J. Gorz. 1969. Genetic studies of induced mutants in *Melilotus alba*. I. Short-internode dwarf, curled leaf, multifoliolate leaf, and cotyledonary branching. *Crop Sci.* 9:607-610.
- Gorz, H.J. 1955. Inheritance of reaction to *Ascochyta caulicola* in sweetclover. *Agron. J.* 47:379-383.
- Gorz, H.J., and F.A. Haskins. 1969. Absence of dominance of the *Cu* gene in influencing *o*-hydroxycinnamic acid content in *Melilotus alba*. *Crop Sci.* 9:79-81.
- Gorz, H.J., F.A. Haskins, G.R. Manglitz, R.R. Smith, and K.P. Vogel. 1992a. Registration of N27 sweetclover germplasm. *Crop Sci.* 32:509.
- Gorz, H.J., F.A. Haskins, G.R. Manglitz, R.R. Smith, and K.P. Vogel. 1992b. Registration of N28 and N29 sweetclover germplasms. *Crop Sci.* 32:510.
- Gorz, H.J., J.E. Specht, and F.A. Haskins. 1975. Inheritance of seed and seedling color in sweetclover. *Crop Sci.* 15:235-238.
- Haskins, F.A., and H.J. Gorz. 1965. Absence of dominance of the *B* gene in influencing β -glucosidase activity in *Melilotus alba*. *Genetics* 51:733-738.
- Kleinohs, A., H.J. Gorz, and F.A. Haskins. 1968. Mutation induction in *Melilotus alba annua* by chemical mutagens. *Crop Sci.* 8:631-632.
- Kneen, B.E., and T.A. LaRue. 1988. Induced symbiosis mutants of pea (*Pisum sativum*) and sweetclover (*Melilotus alba annua*). *Plant Sci.* 58:177-182.
- Markwell, J.P., and T.S. Chelgren. 1988. Chlorophyll expression varies with developmental state in the temperature-sensitive *ch₁* mutation of *Melilotus alba*. *Plant Physiol.* 87:172-175.
- Markwell, J.P., S.J. Danko, H. Bauwe, J. Osterman, H.J. Gorz, and F.A. Haskins. 1986. A temperature-sensitive chlorophyll *b*-deficient mutant of sweetclover (*Melilotus alba*). *Plant Physiol.* 81:329-334.
- Markwell, J.P., A.N. Webber, S.J. Danko, and N.R. Baker. 1985a.

Table 1. Phenotypes and genotypes of 49 *Melilotus alba* genetic stocks.

| Reg. no. | PI no. | Line no. | Seed color [†] | Phenotype | | β-glucosidase activity | Genotype | Reference |
|--------------|--------|----------|---|----------------|----------|------------------------|---|---|
| | | | | Seedling color | Coumarin | | | |
| GS-1 | 549120 | N30 | SG | green | low | low | <i>yyccucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-2 | 549121 | N31 | SG | green | low | high | <i>yyccucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-3 | 549122 | N32 | SG | green | high | low | <i>yyccCuCubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-4 | 549123 | N33 | SG | green | high | high | <i>yyCCuCuBB</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-5 | 549124 | N34 | DG | red | low | low | <i>yyCCucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-6 | 549125 | N35 | DG | red | low | high | <i>yyCCucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-7 | 549126 | N36 | DG | red | high | low | <i>yyCCucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-8 | 549127 | N37 | DG | red | high | high | <i>yyCCCuCuBB</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-9 | 549128 | N38 | LY | green | low | low | <i>YYccucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-10 | 549129 | N39 | LY | green | low | high | <i>YYccucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-11 | 549130 | N40 | LY | green | high | low | <i>YYccCuCubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-12 | 549131 | N41 | LY | green | high | high | <i>YYccCuCuBB</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-13 | 549132 | N42 | MY | red | low | low | <i>YYCCucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-14 | 549133 | N43 | MY | red | low | high | <i>YYCCucubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-15 | 549134 | N44 | MY | red | high | low | <i>YYCCCuCubb</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-16 | 549135 | N45 | MY | red | high | high | <i>YYCCCuCuBB</i> | Gorz and Haskins, 1969; Gorz et al., 1975; Haskins and Gorz, 1965; Specht et al., 1976 |
| GS-17 | 557503 | N46 | | | low | low | <i>cucubb</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-18 | 557504 | N47 | | | high | high | <i>cucubb</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-19 | 557505 | N48 | | | low | low | <i>CuCubb</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-20 | 557506 | N49 | | | high | high | <i>CuCuBB</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-21 | 557507 | N50 | MY | red | | | <i>YYCC</i> | Gorz et al., 1975; Specht et al., 1976 |
| GS-22 | 557508 | N51 | LY | green | | | <i>YYcc</i> | Gorz et al., 1975; Specht et al., 1976 |
| GS-23 | 557509 | N52 | DG | red | | | <i>yyCC</i> | Gorz et al., 1975; Specht et al., 1976 |
| GS-24 | 557510 | N53 | SG | green | | | <i>yycc</i> | Gorz et al., 1975; Specht et al., 1976 |
| Other traits | | | | | | | | |
| GS-25 | 629289 | N54 | susceptibility to stem canker (goose-neck) [‡] | | | | <i>eeGG</i> | Gorz, 1955 |
| GS-26 | 629290 | N55 | resistance to stem canker (goose-neck) [‡] | | | | <i>EEGG</i> | Gorz, 1955 |
| GS-27 | 557511 | N741 | | low | | | <i>cucubb</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-28 | 557512 | N743 | | low | | | <i>cucubb</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-29 | 557513 | N745 | | high | | | <i>CuCubb</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-30 | 557514 | N747 | | high | | | <i>CuCuBB</i> | Gorz and Haskins, 1969; Haskins and Gorz, 1965 |
| GS-31 | 557515 | U362 | folded leaflet | | | | <i>ff</i> | Kleinbofs et al., 1968; Ronnenkamp et al., 1973 |
| GS-32 | 557516 | U363 | elongated stem | | | | <i>el</i> | Kleinbofs et al., 1968; Ronnenkamp et al., 1973 |
| GS-33 | 557517 | U367 | short-petiole dwarf | | | | <i>d₁d₂</i> | Kleinbofs et al., 1968; Ronnenkamp et al., 1973 |
| GS-34 | 557518 | U369 | chlorophyll deficient | | | | <i>chl₂chl₃</i> | Kleinbofs et al., 1968; Ronnenkamp et al., 1975; Specht et al., 1975; Yang et al., 1990 |
| GS-35 | 557519 | U370 | chlorophyll deficient | | | | <i>chl₃chl₉</i> | Kleinbofs et al., 1968; Ronnenkamp et al., 1975; Specht et al., 1975; Yang et al., 1990 |
| GS-36 | 557520 | U371 | chlorophyll deficient | | | | | Bevins et al., 1993; Kleinbofs et al., 1968; Ronnenkamp et al., 1975; Specht et al., 1975; Yang et al., 1990 |
| GS-37 | 557521 | U372 | chlorophyll deficient | | | | <i>chl₁chl₁₁</i> | Bevins et al., 1993; Kleinbofs et al., 1968; Ronnenkamp et al., 1975; Specht et al., 1975; Yang et al., 1990 |
| GS-38 | 557522 | U373 | chlorophyll deficient | | | | <i>chl₂chl₂</i> | Kleinbofs et al., 1968; Ronnenkamp et al., 1975; Specht et al., 1975; Yang et al., 1990 |
| GS-39 | 557523 | U374 | chlorophyll deficient | | | | <i>chl₂chl₅</i> | Gengenbach et al., 1969; Markwell et al., 1985a; Nakitani and Baliga, 1985; Ronnenkamp et al., 1975; Specht et al., 1975; Yang et al., 1990 |
| GS-40 | 557524 | U389 | normal parental line | | | | + / + | Bevins et al., 1993; Gengenbach et al., 1970, 1969; Kneen and LaRue, 1988; Markwell et al., 1986; Specht et al., 1975 |
| GS-41 | 557525 | U390 | short-internode dwarf | | | | <i>dwdw</i> | Gengenbach et al., 1969; Kleinbofs et al., 1968 |
| GS-42 | 557526 | U391 | multifoliate leaf | | | | <i>MJMf</i> | Gengenbach et al., 1969; Kleinbofs et al., 1968 |
| GS-43 | 557527 | U392 | curled leaf | | | | <i>clcl</i> | Gengenbach et al., 1969; Kleinbofs et al., 1968 |
| GS-44 | 557528 | U393 | cotyledonary branching | | | | <i>cbcb</i> | Gengenbach et al., 1969; Kleinbofs et al., 1968 |
| GS-45 | 557529 | U394 | chlorophyll deficient | | | | <i>chl₄</i> | Bevins et al., 1982; Gengenbach et al., 1970; Markwell et al., 1986, 1985b; Specht et al., 1975; Yang et al., 1990 |
| GS-46 | 557530 | U396 | chlorophyll deficient | | | | <i>chl₆</i> | Gengenbach et al., 1970; Kleinbofs et al., 1968; Specht et al., 1975; Yang et al., 1990 |
| GS-47 | 557531 | U397 | chlorophyll deficient | | | | <i>chl₇</i> | Gengenbach et al., 1970; Kleinbofs et al., 1968; Specht et al., 1975; Yang et al., 1990 |
| GS-48 | 557532 | U398 | chlorophyll deficient, dark veins | | | | <i>chl₂chl₅chl₈chl₉</i> | Gengenbach et al., 1970; Kleinbofs et al., 1975; Markwell et al., 1985b; Ronnenkamp et al., 1975; Specht et al., 1975 |
| GS-49 | 557533 | U500 | chlorophyll deficient, short-internode dwarf | | | | <i>chl₂chl₅dwdw</i> | Gengenbach et al., 1970, 1969 |

† Seed colors: SG-silver green; DG-dark green; LY-light yellow; MY-medium yellow.

‡ Caused by *Ascochyta caulicola* (Laub.).

- Fluorescence emission and thylakoid protein kinase activities of three higher plant mutants deficient in chlorophyll *b*. *Biochem. Biophys. Acta.* 808:156–163.
- Markwell, J.P., A.N. Webber, and B. Lake. 1985b. Mutants of sweetclover (*Melilotus alba*) lacking chlorophyll *b*. *Plant Physiol.* 77:948–951.
- Nakitani, H.J., and V. Baliga. 1985. A clover mutant lacking the chlorophyll *a* and *b*-containing protein antenna complexes. *Biochem. Biophys. Res. Commun.* 131:182–189.
- Ronnenkamp, R.R., H.J. Gorz, and F.A. Haskins. 1975. Genetic studies of induced mutants in *Melilotus alba*. IV. Inheritance and complementation of six additional chlorophyll-deficient mutants. *Crop Sci.* 15:187–188.
- Ronnenkamp, R.R., F.A. Haskins, and H.J. Gorz. 1973. Genetic studies of induced mutants in *Melilotus alba*. III. Folded leaflet, elongated stem, and short-petiole dwarf. *Crop Sci.* 13:320–321.
- Specht, J.E., H.J. Gorz, and F.A. Haskins. 1976. Genetic regulation of flavonoid content in seeds and seedlings of *Melilotus alba*. *Phytochemistry* 15:133–134.
- Specht, J.E., F.A. Haskins, and H.J. Gorz. 1975. Contents of chlorophylls *a* and *b* in chlorophyll-deficient mutants of sweetclover. *Crop Sci.* 15:851–853.
- Yang, C., J.C. Osterman, and J. Markwell. 1990. Temperature sensitivity as a general phenomenon in a collection of chlorophyll-deficient mutants of sweetclover (*Melilotus alba*). *Biochem. Genet.* 28:31–40.
-
- K.P. Vogel, H.J. Gorz, and F.A. Haskins, USDA-ARS and Department of Agronomy and Horticulture, 344 Keim Hall, P.O. Box 830937, Univ. of Nebraska, Lincoln, NE 68583-0937. Joint contribution of the USDA-ARS and University of Nebraska-Lincoln. Journal Series Paper No. 14852. Registration by CSSA. Accepted 28 Feb. 2005. *Corresponding author (kpv@unlserve.unl.edu).

doi:10.2135/cropsci2004.0709

Published in *Crop Sci.* 45:1675–1678 (2005).

© Crop Science Society of America

677 S. Segoe Rd., Madison, WI 53711 USA