University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

U.S. Environmental Protection Agency Papers

U.S. Environmental Protection Agency

2010

Assessment of a robust model protocol with accelerated throughput for a human recombinant full length estrogen receptor- α binding assay: Protocol optimization and intralaboratory assay performance as initial steps towards validation

Alexius Freyberger

Bayer Schering Pharma AG, alexius.freyberger@bayerhealthcare.com

Vickie Wilson

U.S. EPA

Marc Weimer

German Cancer Research Center

Shirlee Tan

U.S. EPA

Hoai-Son Tran

Bayer Schering Pharma AG

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usepapapers>

Freyberger, Alexius; Wilson, Vickie; Weimer, Marc; Tan, Shirlee; Tran, Hoai-Son; and Ahr, Hans-Jürgen, "Assessment of a robust model protocol with accelerated throughput for a human recombinant full length estrogen receptor- α binding assay: Protocol optimization and intralaboratory assay performance as initial steps towards validation" (2010). *U.S. Environmental Protection Agency Papers*. 165.

<https://digitalcommons.unl.edu/usepapapers/165>

This Article is brought to you for free and open access by the U.S. Environmental Protection Agency at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in U.S. Environmental Protection Agency Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Alexius Freyberger, Vickie Wilson, Marc Weimer, Shirlee Tan, Hoai-Son Tran, and Hans-Jürgen Ahr

Assessment of a robust model protocol with accelerated throughput for a human recombinant full length estrogen receptor- α binding assay: Protocol optimization and intralaboratory assay performance as initial steps towards validation

Alexius Freyberger^{a,*}, Vickie Wilson^b, Marc Weimer^c, Shirlee Tan^d, Hoai-Son Tran^a, Hans-Jürgen Ahr^a

^a Bayer Schering Pharma AG, BSP-GDD-GED-GTOX Special Toxicology, Bldg 514, Aprather Weg 18, D-42096 Wuppertal, Germany

^b U.S. EPA, Office of Research and Development, NHEERL, Toxicity Assessment Division, Reproductive Toxicology Branch, Research Triangle Park, NC, USA

^c German Cancer Research Center, Biostatistics-C060, Im Neuenheimer Feld 280, D-69120 Heidelberg, Germany

^d U.S. EPA, Office of Science Coordination and Policy, OPPTS, 1200 Pennsylvania Ave., NW, Washington, DC 20460, USA

ARTICLE INFO

Article history:

Received 28 August 2009

Received in revised form

18 December 2009

Accepted 5 January 2010

Available online 13 January 2010

Keywords:

Full length human recombinant estrogen receptor α

Binding assay

Screening for endocrine active compounds

ABSTRACT

Despite about two decades of research in the field of endocrine active compounds, still no validated human recombinant (hr) estrogen receptor- α (ER α) binding assay is available, although hr-ER α is available from several sources. In a joint effort, US EPA and Bayer Schering Pharma with funding from the EU-sponsored 6th framework project, ReProTect, developed a model protocol for such a binding assay. Important features of this assay are the use of a full length hr-ER α and performance in a 96-well plate format. A full length hr-ER α was chosen, as it was considered to provide the most accurate and human-relevant results, whereas truncated receptors could perform differently. Besides three reference compounds [17 β -estradiol, norethynodrel, dibutylphthalate] nine test compounds with different affinities for the ER α [diethylstilbestrol (DES), ethynylestradiol, meso-hexestrol, equol, genistein, *o,p'*-DDT, nonylphenol, *n*-butylparaben, and corticosterone] were used to explore the performance of the assay. Three independent experiments per compound were performed on different days, and dilutions of test compounds from deep-frozen stocks, solutions of radiolabeled ligand and receptor preparation were freshly prepared for each experiment. The ER α binding properties of reference and test compounds were well detected. As expected dibutylphthalate and corticosterone were non-binders in this assay. In terms of the relative ranking of binding affinities, there was good agreement with published data obtained from experiments using a human recombinant ER α ligand binding domain. Irrespective of the chemical nature of the compound, individual IC₅₀-values for a given compound varied by not more than a factor of 2.5. Our data demonstrate that the assay was robust and reliably ranked compounds with strong, weak, and no affinity for the ER α with high accuracy. It avoids the manipulation and use of animals, i.e., the preparation of uterine cytosol as receptor source from ovariectomized rats, as a recombinant protein is used and thus contributes to the 3R concept (reduce, replace, and refine). Furthermore, in contrast to other assays, this assay could be adjusted to an intermediate/high throughput format. On the whole, this assay is a promising candidate for further validation.

© 2010 Elsevier Inc. All rights reserved.

1. Introduction

There is concern that man-made and natural compounds may interfere with the endocrine system and thus may affect wildlife and humans and/or their progeny. Initial studies focused on interactions with estrogen receptor (ER)-mediated signaling and date back to the 1980s [see 1]. These studies have intensified over the years and have been used to characterize a number of chemicals and natural products as compounds with weak estrogenic proper-

ties. More recently, interactions with other receptors such as the androgen receptor have gained attention [2–4]. Both the recommendations to the US Environmental Protection Agency (EPA) by their Endocrine Disruptor Screening and Testing Advisory Committee (EDSTAC) [5] and the OECD conceptual framework for the testing and assessment of endocrine disrupting chemicals [6] recognized receptor binding assays as important tools to study interactions with sex hormone receptors. These assays, therefore, represent important components of the US EPA tier 1 screening battery and level 2 of the OECD's conceptual framework. Interestingly, despite two decades of research in the field of endocrine active compounds and the availability of human recombinant ER from several sources, no validated human recombinant ER bind-

* Corresponding author. Tel.: +49 202 36 8832; fax: +49 202 36 4137.

E-mail address: alexius.freyberger@bayerhealthcare.com (A. Freyberger).

ing assay is available. In a joint effort, US EPA and Bayer Schering Pharma (BSP) (with funding from the European Union-sponsored 6th framework project, ReProTect), developed a model protocol. Additional advisement to this activity was given from an international expert group working on the validation of recombinant ER and AR binding assays under the auspices of the OECD's Validation Management Group for Non-Animal Testing (VMG-NA). Important features of the protocol for competitive binding studies were the use of (a) a commercially available full length human recombinant (hr) estrogen receptor- α (ER α), (b) [3 H]- β -estradiol (1 nM) as ligand, (c) an amount of ER α resulting in a specific binding corresponding to approximately $20 \pm 5\%$ relative to the added [3 H]-ligand, and (d) a 96-well plate format that allows feasibility for up to nine chemicals to be tested at a time (not including the reference chemicals). A full length hr-ER α was chosen, as it was considered to provide the most accurate and human-relevant results, whereas a truncated receptor could perform in a different manner. A commercially available full length hr-ER α was considered helpful for people not experienced in the isolation of receptors, not having the equipment to grow (human) cells prior to isolation, or that do not want to use animals for the preparation of receptors. A set of reference (17 β -estradiol, a strong binder; norethynodrel, a weak binder; and dibutylphthalate, a non-binder) and test compounds (diethylstilbestrol (DES), ethynylestradiol, meso-hexestrol, equol, genistein, *o,p'*-DDT, nonylphenol, *n*-butylparaben, and corticosterone) that had been agreed upon by a Chemical Advisory Board (CAB) jointly appointed by the hr-ER α binding assay validation working group of the OECD's VMG-NA (see above) was used to explore the performance of the assay by evaluating a variety of chemical classes and differing affinities for the receptor. Results were subjected to an independent statistical analysis. The outcome of this effort is reported herein.

2. Materials and methods

2.1. Estrogen receptor- α

Full length human recombinant estrogen receptor- α was obtained from Invitrogen Corporation (Carlsbad, CA, USA) (order no. P2187, lot no. 342161). Functional receptor concentration according to the manufacturer was 2088 pmol/mL with a specific activity of 8700 pmol/mg. The receptor preparation was slowly thawed on ice, and aliquots (20 pmol receptor) were placed in microvials, rapidly frozen in liquid nitrogen and kept deep-frozen at -80°C . Aliquots were thawed only once.

2.2. Chemicals

17 β -Estradiol (CAS 50-28-2) 98%, 17 α -ethynylestradiol (57-63-6) 98%, diethylstilbestrol (DES, 56-53-1) 99%, *m*-hexestrol (84-16-2) 98%, dibutylphthalate, DL-dithiothreitol (DTT) >99%, leupeptin semi-sulfate, bovine serum albumin (BSA), and dextran-coated charcoal was supplied by Sigma-Aldrich (Taufkirchen, Germany). Nonylphenol (84852-15-3) 96.9% and corticosterone (50-22-6) 98.5% were obtained from Fluka (Buchs, Switzerland) through Sigma-Aldrich. Racemic equol (531-95-3) 98.2% was delivered by Apin Chemicals (Abingdon, UK), 1,1,1-trichloro-2,2-bis(*o,p'*-chlorophenyl)ethane (*o,p'*-DDT, 789-02-6) 99.2% was purchased from Supelco (Bellefonte, PA, USA) through Sigma-Aldrich. Norethynodrel (68-23-5, United States Pharmacopial Convention) 100%, genistein (446-72-0, American Custom Chemical Corp.) 99.16%, and *n*-butylparaben (94-26-8, ChemService) 98.2% were kindly provided by U.S. EPA. Tris(hydroxymethyl)aminomethane *p.a.*, anhydrous glycerol, ethanol *p.a.* and DMSO *p.a.* were supplied by E. Merck (Darmstadt, Germany). Ultima Flo APTM and Ultima Gold scintillation cocktails were products of Canberra-Packard (Frankfurt, Germany). [3 H]-17 β -estradiol (estradiol [2,4,6,7,16,17- 3 H(N)], specific activity: 4.07 TBq/mmol, 37 MBq/mL) in ethanol with a radiochemical purity of >97% was supplied by PerkinElmer (Rodgau-Jügesheim, Germany).

2.3. Determination of receptor binding

Receptor binding experiments were performed in 96-well microtiter plates in at least triplicate incubations. Assay buffer (0.2 mM leupeptin, 2 mM DTT, 10 g BSA and 100 mL glycerol made up to 1 L with 10 mM Tris-HCl; pH 7.5) was freshly prepared for each experiment prior to incubation. Stock solutions of reference compounds 17 β -estradiol (10 mM), norethynodrel (10 mM), and dibutylphthalate (100 mM) and test compounds (100 mM) were prepared in ethanol, if 100 mM could

not be achieved, DMSO was used, also for the determination of concurrent total and non-specific binding. Aliquots of stock solutions were kept deep-frozen at -80°C , and each aliquot was used only once and discarded at the end of an individual experiment. Serial dilutions of stocks in solvent were made, then diluted into assay buffer 1:50 as the final step. [3 H]-ligand solution (4 nM if not otherwise indicated resulting in a final concentration of 1 nM) was prepared in assay buffer. Similarly ER α was dissolved in assay buffer. To study receptor binding, assay buffer (80 μL) containing test compound or solvent (2%) was mixed with 40 μL [3 H]-ligand solution for 10–15 min at $2-8^\circ\text{C}$. Then, 40 μL of ER α solution was added and the whole mixture was incubated at $2-8^\circ\text{C}$ overnight under slight (appr. 250 rpm) continuous shaking. Following incubation overnight, 80 μL of a 5% dextran-coated charcoal suspension (the optimal percentage had been empirically determined) in cold assay buffer was added. After mixing in the cold at $2-8^\circ\text{C}$ for 10 min, charcoal was sedimented by centrifugation at 4000 rpm for 5 min in a cooling centrifuge at 4°C . Finally, 50 μL aliquots of the clear supernatant containing the ER α -ligand complex were placed in another 96-well plate and mixed with 200 μL Ultima Flo APTM scintillation cocktail (Canberra-Packard, Frankfurt, Germany) and radioactivity was determined by LSC using a LSC microplate reader (1450 MicrobetaTM Trilux, Wallac, Freiburg, Germany).

In all studies, the actual amount of ligand was determined by liquid scintillation counting (LSC) in a β -counter (1900 TR Counter, Canberra-Packard, Frankfurt, Germany) with quench correction. The maximally tolerated deviation from the theoretical amount was 6%. If the tolerance was exceeded, depending on the deviation, small volumes of buffer or small amounts of radiolabeled ligand were added to meet the limits.

The amount of receptor as specified in the model protocol should result in a specific binding roughly corresponding to $20 \pm 5\%$ relative to the added labeled ligand under conditions of competitive binding, however, slight deviations would be acceptable if not occurring regularly. Furthermore, non-specific binding under these conditions should not exceed 35% of total binding. To determine the amount of receptor necessary experimentally, final concentrations of 0.25, 0.5, 0.75, and 1.0 nM ER α were incubated with 1 nM [3 H]-17 β -estradiol, and non-specific binding was determined in parallel in the presence of 1 μM radioinert ligand. For each condition six replicates were performed.

Saturation of the receptor was studied by incubating the receptor with increasing final concentrations (0.03, 0.06, 0.08, 0.10, 0.30, 0.60, 1.0, 3.0 nM) of [3 H]-17 β -estradiol. Non-specific binding was determined in parallel in the presence of a 1000-fold molar excess of unlabeled 17 β -estradiol. All incubations were performed in triplicate.

Studies on competitive binding were performed using a ligand concentration of 1 nM [3 H]-17 β -estradiol and had to be spread across several 96-well plates. On the first plate, ER α was incubated with the reference compounds 17 β -estradiol (0.010, 0.10, 0.30, 1.0, 3.0, 10, 100, 1000 nM), norethynodrel (3.0, 30, 100, 300, 1000, 3000, 10,000, 30,000 nM), and dibutylphthalate (0.1, 1.0, 10, 100, 1000, 10×10^3 , 100×10^3 , 1000×10^3 nM); furthermore, vehicle controls (no competitor), buffer controls (no competitor, no solvent) and incubations for non-specific binding in the presence of 1 μM unlabeled 17 β -estradiol were performed. On the other plates, the receptor was incubated with test compounds up to 1 mM (0.1, 1.0, 10, 100, 1000, 10×10^3 , 100×10^3 , 1000×10^3 nM), if possible.

In the course of this investigation, for each reference/test compound four independent competitive binding experiments (runs A–D) were performed. For solvent controls, buffer controls, and non-specific binding six replicates were performed each, incubations containing reference/test compounds were studied in triplicate. Run D is the repetition of run C which was invalidated. The statistical analysis for run C had consistently indicated a putative data error for all 12 compounds under investigation, and in the light of a ligand concentration of 1 nM the IC₅₀-value for estradiol of 0.08 nM was implausibly low.

2.4. Data handling

Using the data analysis program, Prism 5.02 (GraphPad Software, Inc., San Diego, CA), a user defined analysis titled "One site – Total + Non-Specific (NS) Binding, accounting for ligand depletion" was performed and entered into Prism templates for all saturation binding runs. More information on the equation used is available from the book by Motulsky and Christopoulos (2003), "Fitting models to biological data using linear and non-linear regression: a practical guide to curve fitting" GraphPad Software, Inc., San Diego, CA. Available online at: <http://www.graphpad.com/manuals/prism4/RegressionBook.pdf> – page 210.

For competitive binding experiments, prior to fitting a dose-response model and estimation of IC₅₀-values mean non-specific binding was subtracted from binding observed for solvent and buffer controls or in the presence of reference/test compounds to achieve the specific binding. After subtraction specific binding in the presence of reference/test compounds was divided by the mean specific binding of the solvent control. The three-parametric log-logistic function

$$f(x) = \frac{\theta_1}{1 + \exp(\theta_2(\log(x) - \theta_3))}$$

was fitted to the transformed data using the `drcm` function of R-package `drc` [7,8]. Parameter θ_1 is the upper asymptote of the response range and corresponds to what is called "Top" in GraphPad Prism, parameter θ_2 is the slope parameter, and parameter θ_3 corresponds to the log of the IC₅₀. Note that there is no estimation

Fig. 1. Determination of the necessary receptor concentration–linear relationship between specific binding and receptor concentration. Under conditions corresponding to competitive binding increasing ER α concentrations were incubated with 1 nM [3 H]-17 β -estradiol. Non-specific binding was assessed in parallel in the presence of 1 μ M unlabeled 17 β -estradiol. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. ER α -bound radiolabel in the supernatant was determined by liquid scintillation counting. Specific binding was calculated as the difference of total and non-specific binding. For each condition six replicates were performed. Means $\pm$ S.D. are given.

of the lower asymptote of the response range. The “Bottom” was assumed to be 0 after correction for non-specific binding in case of full displacement of ligand from the receptor. In the context of receptor binding assays, the IC $_{50}$ -value represents the concentration of a test compound displacing 50% of receptor bound ligand. IC $_{50}$ -values and 95% confidence intervals were calculated using the ED function of package drc. This software program was also used to generate graphs of competitive binding experiments.

3. Results

3.1. Determination of the necessary amount of receptor

The model protocol requested an amount of receptor resulting in a specific binding corresponding to 20 $\pm$ 5% of the added ligand under conditions of competitive binding (i.e., 1 nM [3 H]-17 β -estradiol). When increasing amounts of estrogen receptor- α were incubated with 1 nM [3 H]-17 β -estradiol in the absence and presence of unlabeled 1 μ M 17 β -estradiol (to assess non-specific binding), a linear relationship between total and specific binding and receptor concentration was obtained. Non-specific binding was similar irrespective of the receptor concentration (Fig. 1). The optimal range used for the following studies was at a receptor concentration of 0.5 nM, concomitant specific binding corresponding to 23.1% of the added ligand (Table 1).

3.2. Saturation of estrogen receptor- α binding

To characterize the functionality of the receptor, saturation experiments were performed. Receptor was incubated with increasing concentrations of [3 H]-17 β -estradiol, non-specific binding was assessed in the presence of 1000-fold molar excess of

One Site - Total+NSB accounting for ligand depletion

Fig. 2. (A) Saturation binding data of the human recombinant full length estrogen receptor- α utilizing [3 H]-17 β -estradiol. ER α (nominally 0.5 nM) was incubated with increasing final concentrations (0.03, 0.06, 0.08, 0.10, 0.30, 0.60, 1.0, 3.0 nM) of [3 H]-17 β -estradiol (1 nM = 39,072 dpm). Non-specific binding was determined in parallel in the presence of a 1000-fold molar excess of unlabeled 17 β -estradiol. For each condition three replicates were performed. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. Receptor-bound radiolabeled ligand in the supernatant was determined by liquid scintillation counting. Specific binding was calculated as the difference of total and non-specific binding. Graphs were generated using the non-linear regression software program Prism version 5.02. The figure shows a representative experiment. (B) Scatchard analysis of the human recombinant full length estrogen receptor- α utilizing [3 H]-17 β -estradiol. Data for total and non-specific binding shown in (A) were analysed by Prism version 5.02 taking into account ligand depletion. The figure shows a representative experiment.

unlabeled 17 β -estradiol. Non-specific binding increased linearly and moderately in relation to the [3 H]-17 β -estradiol concentration, whereas total binding increased strongly and almost linearly up to 0.6 nM [3 H]-17 β -estradiol. At higher concentrations of [3 H]-17 β -estradiol total binding was still rising, but at a slower rate. Specific binding increased strongly and almost linearly up to 0.6 nM [3 H]-17 β -estradiol, but at higher concentrations approached a plateau. These findings correspond well with a fully functional receptor showing saturable binding at higher ligand concentrations (Fig. 2). Results of the Scatchard analysis of the saturation experiments are given in Table 2. The observed mean K $_d$ of 0.3734 nM is in the reported range for the human ER α (see also Section 4), and the B $_{max}$ of 0.3439 nM corresponds well to the added receptor concentration of 0.5 nM.

Table 1
Determination of the necessary receptor concentration.

Receptor concentration (nominal) [nM]	Mean specific binding (total incubation) [dpm]	Specific binding (relative to the nominally added ligand) [%]
0.25	4,874	12.5
0.50	9,029	23.1
0.75	12,635	32.3
1.00	16,151	41.3

The assay was performed under conditions corresponding to competitive binding, i.e., in the presence of 1 nM [3 H]-17 β -estradiol (nominally 39,072 dpm) and increasing ER α concentrations as indicated. Non-specific binding was assessed in parallel incubations containing 1 μ M unlabeled 17 β -estradiol. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. Receptor-bound radiolabeled ligand in the supernatant was determined by liquid scintillation counting. Specific binding was calculated as the difference of total and non-specific binding. For each condition six replicates were performed.

Table 2

Scatchard analysis of saturation experiments.

Experiment No.	Kd [nM]	Bmax [nM]
1	0.3822	0.3533
2	0.3851	0.2992
3	0.3530	0.3791
Mean $\pm$ S.D.	0.3734 $\pm$ 0.0178	0.3439 $\pm$ 0.0408

ER α was incubated with increasing final concentrations (0.03, 0.06, 0.08, 0.10, 0.30, 0.60, 1.0, 3.0 nM) of [3 H]-17 β -estradiol. Non-specific binding was determined in parallel in the presence of a 1000-fold molar excess of unlabeled 17 β -estradiol. For each condition three replicates were performed. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. Receptor-bound radiolabeled ligand in the supernatant was determined by liquid scintillation counting. Three independent experiments were run. Scatchard analysis was performed using Graph Pad Prism 5.02 taking into account ligand depletion.

3.3. Competition experiments

General binding characteristics in the absence of competitor are summarized in Table 3. In the presence of optimal ligand concentrations (99–101% of the nominal value) little variability for

total, non-specific and specific bindings was observed across the experiments. Specific binding for vehicle controls corresponded to 22.6–29.8% of the theoretically added ligand, the corresponding coefficient of variation across the six experiments was 12.2%. Similar figures were obtained for specific binding for buffer controls (total binding in the absence of solvent minus non-specific binding), however, there seemed to be a trend to higher values for specific binding in DMSO controls compared to specific binding in buffer controls. Mean non-specific binding corresponded to 15% of total binding, accordingly, total binding mostly represented specific binding to the receptor.

Standard solvent was ethanol, and with the exception of 17 β -estradiol and norethynodrel, all compounds were considered for testing up to 1 mM, an aim that could not always be achieved. A final concentration of 1 mM resulted in strong turbidity for dibutylphthalate, DES, and ethynylestradiol, and in slight turbidity for m-hexestrol and nonylphenol. Genistein and corticosterone could only be tested up to 100 μ M (final conc.) due to their limited solubility in ethanol, whereas 100 μ M *o,p'*-DDT still resulted in turbidity. In order to potentially improve their solubility, genistein, *o,p'*-DDT, and corticosterone together with the set of reference

Table 3

Receptor binding characteristics in competition experiments in the absence of competitor.

Experiment	Solvent	Ligand concentration [% of nominal]	Binding [% of nominally added ligand]				
			Total		Non-specific	Specific	
			Buffer	Solvent		Buffer	Solvent
A	Ethanol	100.6	30.75	30.33	4.35	26.40	25.98
B	Ethanol	99.9	26.88	27.86	4.02	22.86	23.85
D	Ethanol	100.2	34.76	35.01	5.26	29.50	29.75
E	DMSO	99.1	24.21	26.33	3.71	20.50	22.62
F	DMSO	101.3	23.43	27.70	4.55	18.88	23.15
G	DMSO	99.3	29.18	33.77	4.34	24.84	29.43

ER α was incubated with 1.0 nM of [3 H]-17 β -estradiol in the presence or absence of 1% solvent as indicated. Non-specific binding was determined in parallel in the presence of 1 μ M unlabeled 17 β -estradiol. For each condition six replicates were performed. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. Receptor-bound radiolabeled ligand in the supernatant was determined by liquid scintillation counting. Three independent experiments were run for each solvent. Specific binding was calculated as the difference of total and non-specific binding. Corresponding mean values are given.

Fig. 3. Competitive binding to the human recombinant full length estrogen receptor- α of reference and test compounds using ethanol as solvent. Receptor (nominally 0.5 nM) was incubated with 1 nM [3 H]-17 β -estradiol and increasing concentrations of compounds as indicated. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. Receptor-bound radiolabeled ligand in the supernatant was determined by liquid scintillation counting. Binding was corrected for non-specific binding (NSB) and expressed relative to the corresponding solvent control. For the solvent control and for NSB in the presence of 1 μ M 17 β -estradiol six replicates were performed, any other condition was studied in triplicate. Data correspond to run D. For the sake of clarity, only the curve fit, but no individual data points are given.

Fig. 4. Competitive binding to the human recombinant full length estrogen receptor- α of reference and test compounds using DMSO as solvent. Receptor (nominally 0.5 nM) was incubated with 1 nM [3 H]-17 β -estradiol and increasing concentrations of compounds as indicated. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. Receptor-bound radiolabeled ligand in the supernatant was determined by liquid scintillation counting. Binding was corrected for non-specific binding (NSB) and expressed relative to the corresponding solvent control. Data correspond to run D. For the solvent control and for NSB in the presence of 1 μ M 17 β -estradiol six replicates were performed, any other condition was studied in triplicate. Data correspond to run F. For the sake of clarity, only the curve fit, but no individual data points are given.

compounds were also tested using DMSO as solvent. Corticosterone and genistein could be dissolved in DMSO at 100 mM, but the resulting final concentration of 1 mM resulted in slight turbidity for genistein. Similarly, final concentrations of 1 mM dibutylphthalate and 100 μ M *o,p'*-DDT resulted in strong and slight (*o,p'*-DDT) turbidity. Accordingly, only for corticosterone the use of DMSO finally resulted in an improved solubility.

ER α binding data for all compounds dissolved in ethanol expressed as IC₅₀-values are summarized in Table 4. Irrespective of limited solubility in ethanol or in the final assay, IC₅₀-values could be observed for all reference and test compounds except for dibutylphthalate and corticosterone that showed no affinity to the ER α . A compilation of representative displacement curves using ethanol as solvent is shown in Fig. 3. Corresponding relative binding affinities (IC₅₀ of 17 β -estradiol divided by the IC₅₀ of the test chemical and multiplied by 100) are given in Table 5. Genistein, *o,p'*-DDT, and corticosterone together with the set of reference compounds were also tested using DMSO as solvent. A compilation of representative displacement curves using DMSO as solvent is given in Fig. 4. Corresponding IC₅₀-values and relative binding affinities are presented in Tables 6 and 7.

Overall, the following rank order of affinity to the ER α could be established:

17 β -Estradiol > ethynylestradiol > DES > m-hexestrol > equol > norethynodrel > genistein $\approx$ nonylphenol > *o,p'*-DDT > butylparaben

Dibutylphthalate and corticosterone were characterized as non-binders. The observed rank order of affinity to the ER α was in line with our expectations.

4. Discussion

The value of receptor binding assays both as a mechanistic tool to characterize receptor mediated endocrine activity, but also as an important screening assay for endocrine active compounds, is well recognized. Interestingly, after two decades of research in the field of endocrine active compounds, still no validated human recombinant (full length) ER α binding assay is available. Aim of this work was to evaluate a model protocol as a first step towards a validation of a human recombinant full length ER α binding assay.

The model protocol worked well and the general requirements were fulfilled: non-specific binding in competition experiments was clearly below the maximally tolerated 35% of total binding and specific binding roughly corresponded to 20 $\pm$ 5% relative to the added [3 H]-17 β -estradiol for solvent controls. The upper limit was occasionally slightly exceeded, but not on a regular basis and thus was acceptable. Saturation experiments demonstrated that at a ligand concentration of 1 nM binding the receptor was still not saturated. This indicates that on the one hand competitors could readily displace [3 H]-17 β -estradiol from the ER α , and on the other hand the ligand concentration was high enough to provide a readily measurable signal. Scatchard analysis of the saturation experiments provided a mean K_d of 0.3734 nM which is well within the range of reported K_d values for full length human recombinant ER α or ER isolated from human cells, whereas truncated recombinant ER α and possibly rat uterine ER may have higher K_d values (see Table 8). The observed B_{max} of 0.3439 nM corresponds well to the added 0.5 nM receptor protein (determined by means of the hydroxyapatite methodology according to the manufacturer's information) and indicates equivalence of the charcoal and hydroxyapatite methodologies.

When competitive binding was studied, irrespective of the chemical nature of the compound and the solvent used, for a given compound individual IC₅₀-values varied by not more than a factor of 2.5 from one another. Variability of binding affinities relative to 17 β -estradiol (RBA) was in the same range, only for equol (2.9) and genistein (3.3) slightly higher values were observed. In general IC₅₀-values for runs A and B were more comparable to each other than data from run D. However, there was no uniform trend, and IC₅₀-values obtained in run D were either higher, lower, or comparable (n-butylparaben) to those in the other runs. No difference in the quality of the data sets for run D was observed and IC₅₀-values for run D maximally differed from those of runs A and B by a factor of 2.2. This difference – in the absence of a trend – was considered as random. Mean coefficients of variation (CVs) across all compounds were 25% (ethanol as solvent) and 13% (DMSO) for IC₅₀-values and 24% (ethanol) and 13% (DMSO) for relative binding affinities. These low values, especially in light of the low number of independent experiments, indicate that the assay is capable of accurately assessing receptor affinities with only small variability.

Following an ICCVAM/NICEATM recommendation [9] ethanol was used as solvent whenever possible. Accordingly, only few compounds, namely the reference compounds 17 β -estradiol, norethynodrel, and dibutylphthalate as well as genistein, *o,p'*-DDT, and corticosterone were tested using both solvents. The use of DMSO seemed to provide lower variability of the IC₅₀/RBA values and possibly slightly lower IC₅₀-values compared to the use of ethanol. However, in the light of the low number of experiments with DMSO and the small number of compounds studied, it is currently not possible to consider one solvent more appropriate than the other. Basically, both solvents can be used with the assay, and these options can be helpful to achieve optimal dissolution of test compounds.

Optimal interactions of a ligand with its receptor requires that the free ligand concentration is similar to the added concentration, i.e., total binding as the sum of specific and non-specific

Table 4IC₅₀-values achieved for reference and test compounds using ethanol as solvent.

Compound	Experiment	IC ₅₀ -value [nM]	Confidence interval [nM]		Coefficient of variation [%]
			Lower	Upper	
17β-Estradiol	A	1.297	1.1800	1.425	16.2
	B	1.301	1.1880	1.425	
	D	0.965	0.8814	1.057	
	Mean ± S.D.	1.188 ± 0.193			
Norethynodrel	A	451.2	408.2	498.6	30.3
	B	410.8	366.1	460.8	
	D	702.3	623.9	790.6	
	Mean ± S.D.	521.4 ± 157.9			
Dibutylphthalate	A	No affinity ^a			–
	B	No affinity			
	D	No affinity			
	Mean ± S.D.	–			
17α-Ethynylestradiol	A	1.939	1.673	2.246	31.7
	B	2.508	1.930	3.259	
	D	1.294	1.199	1.397	
	Mean ± S.D.	1.914 ± 0.607			
DES	A	7.961	6.980	9.081	18.8
	B	7.723	6.611	9.021	
	D	5.550	4.931	6.247	
	Mean ± S.D.	7.078 ± 1.329			
m-Hexestrol	A	34.62	30.58	39.18	38.7
	B	36.07	20.39	63.81	
	D	16.02	13.16	19.50	
	Mean ± S.D.	28.90 ± 11.18			
Equol, racemic	A	93.78	75.03	117.2	47.5
	B	91.09	67.00	123.8	
	D	197.2	180.0	216.0	
	Mean ± S.D.	127.4 ± 60.50			
Genistein	A	422.6	350.2	509.9	Not calculated
	B	– ^b	–	–	
	D	1049	788.8	1394	
	Mean	735.8			
Nonylphenol	A	868.8	628.7	1201	10.1
	B	1024	692.6	1514	
	D	859.8	589.2	1255	
	Mean ± S.D.	917.5 ± 92.3			
o,p'-DDT	A	– ^b	–	–	Not calculated
	B	1333	950.0	1872	
	D	2399	1937	2970	
	Mean	1866			
n-Butylparaben	A	228.4 × 10 ²	166.6 × 10 ²	313.1 × 10 ²	5.8
	B	254.0 × 10 ²	183.0 × 10 ²	352.7 × 10 ²	
	D	232.0 × 10 ²	162.0 × 10 ²	332.4 × 10 ²	
	Mean ± S.D.	238.1 × 10 ² ± 13.85 × 10 ²			
Corticosterone	A	No affinity ^a			–
	B	No affinity			
	D	No affinity			
	Mean ± S.D.	–			

ERα (nominally 0.5 nM) was incubated with 1 nM [³H]-17β-estradiol and increasing concentrations of the indicated compounds. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. ERα-bound radiolabel in the supernatant was determined by liquid scintillation counting. Binding was corrected for non-specific binding (NSB) and expressed relative to the corresponding solvent control. For the solvent control and NSB in the presence of 1 μM 17β-estradiol six replicates were performed, any other condition was studied in triplicate. IC₅₀-values were calculated by means of the software R-package *drc* using a three-parametric log-logistic function. Results from three independent experiments A, B, and D are given.

^a Dibutylphthalate and corticosterone were tested up to 100 μM.

^b No IC₅₀-value was calculated due to unclear dose response (displacement curves were comparable to corresponding experiments, but contained one or two aberrant data points at low concentrations).

binding should not exceed 10%. However, assay miniaturization in order to allow intermediate or high throughput, increases the risk of ligand depletion (where the actual free ligand concentration is significantly lower than the added ligand concentration), since receptor concentrations are rather high in these assays in order to achieve adequate signal sizes [10]. In competition experiments, ligand depletion results in a shift of competition curves/IC₅₀-values to the right, i.e., to higher concentrations [11]. With regard to the

experiments reported herein, total binding in vehicle controls of competition experiments corresponded to 26–35% of the added ligand (see Table 3). Thus, in all experiments slight ligand depletion had occurred. Modeling of ligand depletion [11] indicated that the impact on competition experiments of a depletion up to 38% was small. Accordingly, competition experiments reported herein was sufficiently accurate in terms of ligand depletion and no corrective action is needed. The use of relative binding affinities with all com-

Table 5

Relative binding affinities for reference and test compounds using ethanol as solvent.

Compound	Experiment No.	Relative binding affinity [%]	Coefficient of variation [%]
17 β -Estradiol		100	–
Norethynodrel	A	0.2875	38.9
	B	0.3167	
	D	0.1374	
	Mean $\pm$ S.D.	0.2472 $\pm$ 0.0962	
Dibutylphthalate	A	No affinity	–
	B	No affinity	
	D	No affinity	
	Mean $\pm$ S.D.	–	
17 α -Ethinylestradiol	A	66.89	17.9
	B	51.87	
	D	74.57	
	Mean $\pm$ S.D.	64.45 $\pm$ 11.55	
DES	A	16.29	3.3
	B	16.85	
	D	17.39	
	Mean $\pm$ S.D.	16.84 $\pm$ 0.55	
m-Hexestrol	A	3.746	30.4
	B	3.607	
	D	6.024	
	Mean $\pm$ S.D.	4.459 $\pm$ 1.357	
Equol, racemic	A	1.383	48.2
	B	1.428	
	D	0.489	
	Mean $\pm$ S.D.	1.100 $\pm$ 0.530	
Genistein	A	0.3069	Not calculated
	B	–	
	D	0.0920	
	Mean	0.1995	
Nonylphenol	A	0.1493	14.4
	B	0.1271	
	D	0.1122	
	Mean $\pm$ S.D.	0.1295 $\pm$ 0.0186	
o,p'-DDT	A	–	Not calculated
	B	0.0976	
	D	0.0402	
	Mean	0.0689	
n-Butylparaben	A	0.00568	15.4
	B	0.00512	
	D	0.00416	
	Mean $\pm$ S.D.	0.00499 $\pm$ 0.00077	
Corticosterone	A	No affinity	–
	B	No affinity	
	D	No affinity	
	Mean $\pm$ S.D.	–	

Relative binding affinities were calculated from data given in Table 4.

pounds being tested under the same conditions should even further minimize a small impact of ligand depletion.

An important step in the evaluation of an assay is the comparison of assay data with data from similar assays observed for the same test compounds. In Table 9, results from the investigation reported herein are compared to ER α binding data from several sources. With regard to data generated for full length human recombinant ER α [12,13] and for a human recombinant ER α ligand binding domain as receptor source (by the Chemicals Evaluation and Research Center (CERI) in Japan with funding by the Japanese Ministry of Economy, Trade and Industry [14]), a good to very good fit for the binding relative to estradiol (RBA) was observed. For compounds with weak affinity to the ER α the concurrence was excellent, and also for compounds with stronger affinity, the ranking was comparable, although in absolute terms in our hands the relative binding values for DES, ethinylestradiol and m-hexestrol were lower mainly compared to the CERI data. A similar constellation was observed when comparing our RBAs with those observed

for rat uterine cytosol as receptor source [15,16]. It remains to be elucidated further whether these lower RBAs are characteristic of the human recombinant full length ER α as opposed to a human truncated or rat receptor or are related to other reasons such as the solvent used or assay buffer composition (e.g., in aqueous media DES can be converted to its much weaker cis-isomer). If this was the case for a large portion of DES, the lower RBA of DES could readily be explained [17]. The CERI assay utilized DMSO as the solvent in all experiments, while this assay used EtOH as the solvent for DES. Finally, as reported here and by others [14,15], dibutylphthalate and corticosterone did not show affinity for the ER α .

On the whole, our data demonstrate that the assay was robust and reliably ranked compounds with strong, weak, and without affinity for the ER α with high accuracy. Using a human recombinant full length estrogen receptor- α , it avoids the manipulation and use of animals, i.e., the preparation of uterine cytosol from ovariectomized rats, and thus contributes to the 3R concept. Furthermore, in contrast to other assays, our assay could be adjusted

Table 6IC₅₀-values achieved for reference and test compounds using dimethylsulfoxide as solvent.

Compound	Experiment No.	IC ₅₀ -value [nM]	Confidence interval [nM]		Coefficient of variation [%]
			Lower	Upper	
17β-Estradiol	E	0.9857	0.8443	1.151	13.4
	F	0.8193	0.7280	0.9220	
	G	0.7654	0.6624	0.8845	
	Mean ± S.D.	0.8568 ± 0.1148			
Norethynodrel	E	424.5	384.5	468.8	10.0
	F	353.4	308.9	404.4	
	G	365.5	298.6	447.4	
	Mean ± S.D.	381.1 ± 38.0			
Dibutylphthalate	E	No affinity ^a			–
	F	No affinity			
	G	No affinity			
	Mean ± S.D.	–			
Genistein	E	744.5	611.3	906.6	10.5
	F	776.4	678.6	888.4	
	G	633.1	520.1	770.7	
	Mean ± S.D.	718.0 ± 75.2			
o,p'-DDT	E	1405	1081	1825	19.0
	F	1770	1339	2340	
	G	1225	923.6	1625	
	Mean ± S.D.	1467 ± 278			
Corticosterone	E	No affinity ^a			–
	F	No affinity			
	G	No affinity			
	Mean ± S.D.	–			

ERα (nominally 0.5 nM) was incubated with 1 nM [³H]-17β-estradiol and increasing concentrations of the indicated compounds. Following incubation overnight, unbound radiolabeled ligand was removed by absorption to dextran-bound charcoal (DBC) and sedimentation of DBC by centrifugation. ERα-bound radiolabel in the supernatant was determined by liquid scintillation counting. Binding was corrected for non-specific binding (NSB) and expressed relative to the corresponding solvent control. For the solvent control and NSB in the presence of 1 μM 17β-estradiol six replicates were performed, any other condition was studied in triplicate. IC₅₀-values were calculated by means of the software R-package *drc* using a three-parametric log-logistic function. Results from three independent experiments E, F, and G are given.

^a Dibutylphthalate was tested up to 100 μM, corticosterone up to 1 mM.

to an intermediate/high throughput format. In terms of the ECVAM modular approach of validation, the data set presented herein corresponds to the first two modules, namely test definition and within-laboratory variability [18]. Further testing of the protocol in collaboration with OECD in additional laboratories is ongoing and will provide an in-depth evaluation of transferability and inter-laboratory variability. The assay may be used to screen for

endocrine-modulating compounds with affinity for the ERα, to further investigate equivocal results from other *in vitro* assays such as transactivation assays, and it may also be used as a mechanistic tool in order to characterize effects on the endocrine system observed in animal studies. Whether ERα binding studies can reliably predict the outcome of corresponding *in vivo* screening assays such as the uterotrophic assay remains to be explored.

Table 7

Relative binding affinities for reference and test compounds using dimethylsulfoxide as solvent.

Compound	Experiment No.	Relative binding affinity [%]	Coefficient of variation [%]
17β-Estradiol		100	–
Norethynodrel	E	0.2322	5.8
	F	0.2318	
	G	0.2094	
	Mean ± S.D.	0.2245 ± 0.0131	
Dibutylphthalate	E	No affinity	–
	F	No affinity	
	G	No affinity	
	Mean ± S.D.	–	
Genistein	E	0.1324	11.3
	F	0.1055	
	G	0.1209	
	Mean ± S.D.	0.1196 ± 0.0135	
o,p'-DDT	E	0.0702	20.5
	F	0.0463	
	G	0.0625	
	Mean ± S.D.	0.0596 ± 0.0122	
Corticosterone	E	No affinity	–
	F	No affinity	
	G	No affinity	
	Mean ± S.D.	–	

Relative binding affinities were calculated from data given in Table 5.

Table 8

Kd values for estrogen receptors isolated from various sources.

Estrogen receptor	Expression system	Receptor preparation	Methodology ^a	Kd [nM] ^b	Reference
Full length human recombinant ER α	Insect cells	Purified protein	Charcoal	0.37	This investigation
Full length human recombinant ER α	Insect cells	Extract	Charcoal	0.22	[12]
Human ER (mainly α -subtype) from MCF-7 cells	–	Cytosol	Charcoal	0.13	[19]
Human ER (mainly α -subtype) from MCF-7 cells	–	Cell extract	No information	0.36	[20]
Full length human recombinant ER α	Yeast	Yeast extract	No information	0.35	[20]
Human recombinant ER α –ligand binding domain	Yeast	Yeast extract	No information	1.00	[20]
Human recombinant ER α –ligand binding domain	Bacteria	Bacteria extract	No information	1.49	[20]
Full length quail recombinant ER α	Insect cells	Extract	Charcoal	0.58	[12]
Full length alligator recombinant ER α	Insect cells	Extract	Charcoal	0.44	[12]
Full length salamander recombinant ER α	Insect cells	Extract	Charcoal	0.28	[12]
Full length fathead minnow recombinant ER α	Insect cells	Extract	Charcoal	0.58	[12]
Rat ER (mainly α -subtype) from rat uterus, laboratory X	–	Cytosol	Hydroxyapatite	0.15–1.18	[16]
Rat ER (mainly α -subtype) from rat uterus, laboratory Y	–	Cytosol	Hydroxyapatite	0.04–0.24	[16]
Rat ER (mainly α -subtype) from rat uterus, laboratory Z	–	Cytosol	Hydroxyapatite	0.8–1.5	[16]

^a Methodology used to separate receptor-bound from unbound ligand.^b [³H]-17 β -estradiol was used as ligand for Kd determinations.**Table 9**Comparison of relative ER α binding affinities to published data.

Compound	Investigation					
	This investigation (full length human recombinant) RBA [%]	Rider et al. [12] (full length human recombinant) RBA [%]	Kuiper et al. [13] (full length human recombinant) RBA [%]	METI/CERI [14] (human recombinant ligand binding domain) RBA [%]	Blair et al. [15] (rat uterus cytosol) RBA [%]	EPA validation [16] (rat uterus cytosol) RBA [%]
17 β -Estradiol	100	100	100	100	100	100, 100, 100 ^a
Ethinylestradiol	64.5	91.2	Not done	142	190	165, 268, 113
DES	16.8	Not done	236	88.9	400	590, 284, equivocal ^b
m-Hexestrol	4.46	Not done	Not done	37.6	300	345, 2090, 270
Norethynodrel	0.247/0.225 ^{DM}	Not done	Not done	0.282	0.204	0.135, 0.147, 0.130
Butylparaben	0.00499	Not done	Not done	0.018	Not done	0.00416, 0.0105, 0.00086
Equol	1.10	Not done	Not done	Not done	Not done	0.0812, 0.0542, 0.039
Genistein	0.199/0.120 ^{DM}	Not done	4/0.7	0.12	Not done	0.971, 4.81, 0.402
Nonylphenol	0.130	Not done	0.05	0.108	0.019–0.037	0.125, 0.101, equivocal
<i>o,p'</i> -DDT	0.0689/0.0596 ^{DM}	Not done	0.01	Not done	0.001	Negative, 14,400, negative
Dibutylphthalate	No affinity	Not done	Not done	No IC ₅₀ , RBA<0.001	No affinity	Not done
Corticosterone	No affinity	No affinity	Not done	Not done	No affinity	0.00103, 2.04, equivocal

For our data, solvent is ethanol, if not otherwise indicated.

^a Data from three laboratories (X, Y, and Z).^b Equivocal: displacement of labeled ligand was observed, however, no characteristic binding curve was observed.

Conflict of interest

There are no conflicts of interest.

Acknowledgements

The authors wish to thank the OECD estrogen receptor binding group for helpful discussions. The donation of norethynodrel, genistein, and n-butylparaben by the US Environmental Protection Agency is gratefully acknowledged. This work was supported in part by funding to Bayer Schering Pharma AG from the EU-sponsored project ReProTect (LSHB-CT-2004-503257). The statistical analysis (M.W.) was performed as contract work for ECVAM (contract no. CCR.IHCP.C432976.XO).

Disclaimer: This manuscript has been reviewed by the U.S. EPA and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the agency, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

References

- [1] McLachlan JA, Korach KS, Newbold RR, Degen GH. Diethylstilbestrol and other estrogens in the environment. *Fundam Appl Toxicol* 1984;4:686–91.
- [2] Kelce WR, Stone CR, Laws SC, Gray LE, Kemppainen JA, Wilson EM. Persistent DDT metabolite *p,p'*-DDE is a potent androgen receptor antagonist. *Nature* 1995;375:581–5.
- [3] Lambright CR, Ostby J, Bobseine K, Wilson V, Hotchkiss AK, Mann PC, et al. Cellular and molecular mechanisms of action of linurone: an antiandrogenic herbicide that produces reproductive malformations in male rats. *Toxicol Sci* 2000;56:389–99.
- [4] Wilson VS, Lambright C, Ostby J, Gray Jr LE. In vitro and in vivo effects of 17 β -trenbolone: a feedlot effluent contaminant. *Toxicol Sci* 2002;70:202–11.
- [5] EDSTAC. 1998. Final report. Available via EPA. <http://www.epa.gov/scipoly/oscpendo/history/finalrpt.htm>.
- [6] OECD, Organisation for Economic Cooperation and Development. Draft summary report of the sixth meeting of the task force on Endocrine Disrupters Testing and Assessment (EDTA 6); 2003. Paris.
- [7] Ritz C, Streibig JC. Bioassay analysis using R. *J Stat Software* 2005;12(5):1–22, <http://www.bioassay.dk>.
- [8] R Development Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2009. ISBN 3-900051-07-0, <http://www.R-project.org>.
- [9] ICCVAM/NICEATM. Expert panel evaluation of the validation status of in vitro test methods for detecting endocrine disruptors: estrogen and androgen receptor binding and transcriptional activation assays. Expert panel final report; 2002. Available via ICCVAM/NICEATM <http://iccvam.niehs.nih.gov/>.
- [10] Scaramellini Carter CM, Leighton-Davies JR, Charlton SJ. Miniaturized receptor binding assays: complications arising from ligand depletion. *J Biomol Screening* 2007;12:255–66.
- [11] Hulme EC, Birdsall NJM. Strategy and tactics in receptor-binding studies. In: Hulme EC, editor. *Receptor–ligand interactions. A practical approach*. Oxford, New York, Tokyo: IRL Press at Oxford University Press; 1992. p. 63–176.

- [12] Rider CV, Hartig PC, Cardon MC, Wilson VS. Development of a competitive binding assay system with recombinant estrogen receptors from multiple species. *Toxicol Lett* 2009;184:85–9.
- [13] Kuiper GGJM, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, et al. Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor β . *Endocrinology* 1998;139:4252–63.
- [14] Ministry of Economy, Trade and Industry, Japan. Risk assessment of endocrine disrupters. Current status of development of testing methods for endocrine disrupters, Appendix 1. 2002 (revised 2003) <http://www.meti.go.jp/english/report/data/g020205ae.html>.
- [15] Blair RM, Fang H, Branham WS, Hass BS, Dial SL, Moland CL, et al. The estrogen receptor relative binding affinities of 188 natural and xenochemicals: structural diversity of ligands. *Toxicol Sci* 2000;54:138–53.
- [16] U.S. Environmental Protection Agency. Endocrine Disruptor Screening Program (EDSP)—Estrogen Receptor Binding Assay Using Rat Uterine Cytosol (ER-RUC) Peer Review Material;2009 <http://www.epa.gov/scipoly/oscpendo/pubs/assayvalidation/estrogen.html>.
- [17] Katzenellenbogen JA, Carlson KE, Katzenellenbogen BS. Facile geometric isomerization of phenolic non-steroidal estrogens and antiestrogens: limitations to the interpretation of experiments characterizing the activity of individual isomers. *J Steroid Biochem* 1985;22:589–96.
- [18] Hartung T, Bremer S, Casati S, Coecke S, Corvi R, Fortaner S, et al. A modular approach to the ECVAM principles on test validity. *ATLA* 2004;32:467–72.
- [19] Eckert RL, Katzenellenbogen BS. Physical properties of estrogen receptor complexes in MCF-7 human breast cancer cells. *J Biol Chem* 1982;257:8840–6.
- [20] Harris Wooge C, Nilsson GM, Heierson A, McDonnell DP, Katzenellenbogen BS. Structural requirements for high affinity ligand binding of estrogen receptors: a comparative analysis of truncated and full length estrogen receptors expressed in bacteria, yeast, and mammalian cells. *Mol Endocrinol* 1992;6:861–9.