

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Transactions of the Nebraska Academy of
Sciences and Affiliated Societies

Nebraska Academy of Sciences

1989

Range Extensions for Some Nebraska Bryophytes

Linda L. Spessard
Kearney State College

Follow this and additional works at: <https://digitalcommons.unl.edu/tnas>



Part of the [Life Sciences Commons](#)

Spessard, Linda L., "Range Extensions for Some Nebraska Bryophytes" (1989). *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*. 174.
<https://digitalcommons.unl.edu/tnas/174>

This Article is brought to you for free and open access by the Nebraska Academy of Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Transactions of the Nebraska Academy of Sciences and Affiliated Societies by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Range Extensions for Some Nebraska Bryophytes

Linda L. Spessard
Department of Biology
Kearney State College
Kearney, Nebraska 68849-0531

Range extensions are presented for seven species of mosses in Nebraska: *Amblystegium serpens*, *Brachythecium oxycladon*, *Bryum caespiticium*, *B. capillare*, *Ephemerum cassinervum*, *Orthotrichum pumilum*, and *O. stellatum*. Six new county records are listed.

INTRODUCTION

The list of mosses given here has been made during the past several years. The collections have been identified, catalogued, and placed in the Kearney State College (KSC) bryophyte herbarium. Most of the reports are from central and western counties where little bryological work has been done. A literature search shows that no reports have been made from Adams County. Churchill (1982), Koch (1975), and Spessard (1984) reported collections from Hitchcock County, Spessard (1982) from Loup County, and Churchill (1976) from Webster County.

Nomenclature follows Crum et al. (1973), and new county reports are identified by an asterisk (*) preceding the name of the county. Collectors and collections numbers are given for each voucher specimen.

MOSSES

Amblystegium serpens (Hedw.) BSG

HITCHCOCK COUNTY: 1.6 km W of Trenton, on soil; "G" (Grigg). A highly variable species, *A. serpens* is the most widespread moss in Nebraska. It occurs in almost every known habitat and seems to adapt to most environments. It grows on a number of substrates: several soil types, rotting logs, tree bark, and moist rocks. The distribution of *A. serpens* in Nebraska appears to be statewide. Reported from Culbertson City Park by Spessard, 1984.

Brachythecium oxycladon (Brid.) Jaeg. & Sauerb.

*ADAMS COUNTY: northeast of Hastings city limit, 907 N. Burlington, 1.6 km w of post office; on soil; 107 (Forney). Occurs less frequently than *B. salebrosum* (Web. & Mohr) BSG, with which it is most likely to be confused. With practice it is possible to distinguish the two by using the size of the alar cells; *B. oxycladon* has small and quadrate alar cells, while those of *B. salebrosum* are larger, more lax, and appear similar in shape to the lower leaf cells.

Bryum caespiticium Hedw.

*HITCHCOCK COUNTY: 701 W. "E" St., Trenton; on sandy soil; "E" (Grigg). A difficult taxon to identify when not fruiting, it is sometimes confused with *B. lisae* De Not. var. *cuspidatum* (BSG) Marg. Crum (1973) indicates that *B. caespiticium* has a "silky appearance" than *B. lisae* var. *cuspidatum*. The distribution in Nebraska appears to be statewide.

Bryum capillare Hedw.

*HITCHCOCK COUNTY: Elm Creek at Trenton Dam, 0.4 km N of confluence with Republican River; "D" (Grigg). Found sporadically across the state, *B. capillare* seems to be found in moister habitats than other species of *Bryum*.

Ephemerum cassinervum (Schwaegr.) Hampe

*HITCHCOCK COUNTY: confluence of Elm Creek and Republican River at Trenton Dam, on sandy soil; "F" (Grigg). A rare moss in Nebraska, with previous reports from Lancaster and York counties (Churchill, 1977). Whether this moss is rare in Nebraska or is rarely found because of its ephemeral nature is unknown.

Orthotrichum pumilum Sw.

*WEBSTER COUNTY: 0.2 km S and 3.6 km W of Guide Rock; 27 (Friskopp). *Orthotrichum pumilum* is the most widespread of the corticolous mosses in Nebraska, with a statewide distribution. When collected, it is often fruiting even when environmental conditions are not necessarily conducive to fruiting of bryophytes.

Orthotrichum stellatum Brid.

*LOUP COUNTY: Taylor City Park, on tree bark; 6899 (Spessard). The only other report of this moss from Nebraska was made by Churchill (1977) from Sioux County.

ACKNOWLEDGMENTS

The author wishes to acknowledge J. R. Forney, Dudley Friskopp, and Roger Grigg for their willingness to collect mosses for the Kearney State College bryophyte herbarium.

LITERATURE CITED

- Churchill, S. P. 1976. Contributions toward a moss flora of Nebraska. *Bryologist* 79: 241–242.
- _____. 1977. Contributions toward a moss flora of Nebraska II. *Bryologist* 80: 160–162.
- _____. 1982. Mosses of the Great Plains VIII. Additions. *Bryologist* 85: 218–221.
- Crum, H. A. 1973. *Mosses of the Great Lakes forest*. Ann Arbor, University of Michigan Herbarium: 404 pp.
- _____, W. C. Steere, and L. E. Anderson. 1973. A new list of mosses of North America, north of Mexico. *Bryologist* 76: 85–130.
- Koch, R. G. 1971. Pottiaceae in Nebraska. *Bryologist* 74: 206–207.
- Spessard, L. L. 1982. A bryological survey of north-central Nebraska. *Transactions of the Nebraska Academy of Sciences* 10: 17–20.
- _____. 1984. A bryological survey of southwestern Nebraska. *Transactions of the Nebraska Academy of Sciences* 12: 19–22.