University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

US Department of Energy Publications | US Department of Energy

1-1-2011

Persistence Of Uranium Groundwater Plumes: Contrasting Mechanisms At Two Contaminated Doe Sites

John M. Zachara
Pacific Northwest National Laboratory, john.zachara@pnl.gov

James A. Davis
Lawrence Berkeley National Laboratory

Philip E. Long
Pacific Northwest National Laboratory

Kenneth H. Williams
Lawrence Berkeley National Laboratory

Mark Freshley
Northwest National Laboratory

See next page for additional authors

Follow this and additional works at: http://digitalcommons.unl.edu/usdoepub

Part of the Bioresource and Agricultural Engineering Commons

Zachara, John M.; Davis, James A.; Long, Philip E.; Williams, Kenneth H.; Freshley, Mark; and McKinley, James P., "Persistence Of Uranium Groundwater Plumes: Contrasting Mechanisms At Two Contaminated Doe Sites" (2011). *US Department of Energy Publications.* Paper 177.
http://digitalcommons.unl.edu/usdoepub/177

This Article is brought to you for free and open access by the US Department of Energy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in US Department of Energy Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors
John M. Zachara, James A. Davis, Philip E. Long, Kenneth H. Williams, Mark Freshley, and James P. McKinley

This article is available at DigitalCommons@University of Nebraska - Lincoln: http://digitalcommons.unl.edu/usdoepub/177

2011 GSA Annual Meeting in Minneapolis (9–12 October 2011)
Paper No. 88-7

Presentation Time: 10:10 AM-10:30 AM

PERSISTENCE OF URANIUM GROUNDWATER PLUMES: CONTRASTING MECHANISMS AT TWO CONTAMINATED DOE SITES

ZACHARA, John[1], DAVIS, James A.[2], LONG, Philip E.[3], WILLIAMS, Kenneth H.[2], FRESHLEY, Mark[1], and MCKINLEY, James P.[4], (1) Fundamental Sciences Directorate, Northwest National Laboratory, PO Box 999; MS K8-96, Richland, WA 99354, john.zachara@pnnl.gov, (2) Earth Sciences Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Rd. MS-90-1116, Berkeley, CA 94720, (3) Energy and Environment Directorate, Pacific Northwest National Laboratory, PO Box 999, Richland, WA 99354, (4) Pacific Northwest National Laboratory, 902 Battelle Boulevard, P.O. Box 999, MSIN K6-96, Richland, WA 99352

We examine subsurface uranium (U) plumes at two U.S. Department of Energy sites located near large river systems. Following excavation of contaminated materials, both sites were projected to naturally flush remnant uranium contamination to levels below regulatory limits (e.g., 30 µg/L), within 10 years at the Hanford 300 Area (Columbia River) and 12 years at the Rifle site (Colorado River). The observed rate of attenuation was much lower than expected at both sites. Current understanding suggests that the two sites have common, but also differing mechanisms controlling plume persistence. At Hanford, the persistent source is adsorbed U(VI) in the deep vadose zone, released to the aquifer during water table excursions associated with spring high-water in the Columbia River. The release of U(VI) from the vadose zone and its transport within the oxic, coarse-textured aquifer sediments is dominated by kinetically-limited surface complexation. While the U sources sustaining the plume at the Rifle site are multiple, there is evidence that slow oxidation of naturally reduced contaminant U(IV) in the saturated zone, as well as a continuous influx of natural U(VI) from up-gradient sources, is important. Equilibrium surface complexation controls the U(VI) migration rate in the sub-oxic Rifle groundwater. Microbiologic activity is diverse at both sites. Strains of *Geobacter* and other metal reducing bacteria are present, typically at low natural abundance, capable of enzymatic U(VI) reduction in localized zones of accumulated detrital organic carbon, or after organic carbon amendment. Major differences between the sites include the geochemical nature of residual contaminant U; the nature of mineral adsorbents and the rates of kinetic processes (biotic and abiotic) influencing U(VI) solid-liquid distribution; the presence of detrital organic matter and the spatial heterogeneity in microbially-driven redox properties; and the magnitude of groundwater hydrologic dynamics controlled by river-stage fluctuations and aquifer properties (e.g., hydraulic gradient and conductivity). The comparative analysis of these sites provides important guidance to the

characterization, modeling, and remediation of groundwater contaminant plumes influenced by surface water interaction that are common worldwide.

2011 GSA Annual Meeting in Minneapolis (9–12 October 2011)
General Information for this Meeting

Session No. 88
Uranium in Groundwater: Biogeochemical Controls on Mobility and Persistence
Minneapolis Convention Center: Room 211AB
8:00 AM-12:00 PM, Monday, 10 October 2011

Geological Society of America *Abstracts with Programs*, Vol. 43, No. 5, p. 230

© Copyright 2011 The Geological Society of America (GSA), all rights reserved. Permission is hereby granted to the author(s) of this abstract to reproduce and distribute it freely, for noncommercial purposes. Permission is hereby granted to any individual scientist to download a single copy of this electronic file and reproduce up to 20 paper copies for noncommercial purposes advancing science and education, including classroom use, providing all reproductions include the complete content shown here, including the author information. All other forms of reproduction and/or transmittal are prohibited without written permission from GSA Copyright Permissions.

This article is a U.S. government work, and is not subject to copyright in the United States.