

11-1974

OER for Mixed Neutrons and γ Rays

Robert Katz

University of Nebraska-Lincoln, rkatz2@unl.edu

S. C. Sharma

University of Nebraska-Lincoln, sharma@uta.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/physickatz>



Part of the [Physics Commons](#)

Katz, Robert and Sharma, S. C., "OER for Mixed Neutrons and γ Rays" (1974). *Robert Katz Publications*. 189.
<http://digitalcommons.unl.edu/physickatz/189>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Robert Katz Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Correspondence

THE EDITOR – SIR,

OER for Mixed Neutrons and γ Rays

A formula for the OER in a mixed field of neutrons and gamma rays (Hall, 1972a; 1972b) is given as

$$O_{\text{mix}} = O_n O_\gamma [1 + p_n (R_n - 1)] / [O_\gamma R_n p_n + O_n (1 - p_n)]$$

where R_n is the RBE of neutrons relative to γ rays under aerated conditions, O_{mix} is the OER for the mixture of neutrons and γ rays, O_n and O_γ are the OERs for neutrons and γ rays, respectively, and p_n is the proportion of absorbed dose due to neutrons. Recent calculations of the survival, RBE, and OER of kidney cells in a mixed field of 14 MeV neutrons and γ rays (Katz and Sharma, 1974) made for comparison with experimental findings with Chinese hamster ovary cells (Railton *et al.*, 1974) were in good agreement with the Hall formula. In consequence, we have explored the matter more fully, calculating O_{mix} for neutrons of different energies (0.5, 1.0, 2.0, 14 MeV, and mean energy 7.5 MeV having the energy spectrum of the Hammersmith Hospital generator), onto different cells (HeLa, leukemia, human kidney), at a range of survival levels (1, 10, 20, 50, 90 per cent). Our results, calculated from track structure theory (Katz and Sharma, 1973) and secondary particle-energy spectra in tissue from these neutrons (Caswell and Coyne, 1973; Dennis, 1972) are within 5 per cent of the values of O_{mix} found from the Hall formula, when that formula is supplied with calculated values of R_n , O_n , and O_γ , at appropriate survival levels. Our calculations suggest that the Hall formula may be used with confidence for neutrons of different energies and for pions.

Yours, etc.,

Robert Katz

S. C. Sharma

Physics Department, University of Nebraska–Lincoln
Lincoln, Nebraska 68508 U.S.A.

References

- Caswell, R. S., and Coyne, J. J., 1973. *Proceedings of the Fourth Symposium on Microdosimetry, Verbania-Pallanza, Italy* (Euratom).
- Dennis, J. A., 1972. Private communication.
- Hall, E. J., 1972a. A determination of the oxygen enhancement ratio for ^{252}Cf using cultured mammalian cells. *British Journal of Radiology*, **45**, 284-288.
- Hall, E. J., 1972b. *Ibid.*, **45**, 628.
- Katz, R., and Sharma, S. C., 1973. Response of cells to fast neutrons, stopped pions, and heavy ion beams. *Nuclear Instruments and Methods*, **111**, 93-116.
- Katz, R., and Sharma, S. C., 1974. Cellular survival in a mixed radiation environment. *International Journal of Radiation Biology*, in press.
- Railton, R., Porter, D., Lawson, R. C., and Hannan, W. J., 1974. The oxygen enhancement ratio and relative biological effectiveness for combined irradiations of Chinese hamster cells by neutrons and gamma-rays. *International Journal of Radiation Biology*, **25**, 121-127.